

A .gov website belongs to an official government organization in the United States. A lock ( ) or https:// means you've safely connected to the .gov website. Share sensitive information only on official, secure websites.

Standard Precautions are designed to both protect dental health care personnel (DHCP) and prevent DHCP from spreading infections among patients. Standard Precautions are the minimum infection prevention practices that apply to all patient care, regardless of suspected or confirmed infection status of the patient, in any setting where health care is delivered. Standard Precautions include:

Each element of Standard Precautions is described in the following sections. Education and training are critical elements of Standard Precautions, because they help DHCP make appropriate decisions and comply with recommended practices. When Standard Precautions alone cannot prevent transmission, they are supplemented with Transmission-Based Precautions. This second tier of infection prevention is used when patients have diseases that can spread through contact, droplet or airborne routes (e.g., skin contact, sneezing, coughing), and are always used in addition to Standard Precautions. Dental settings are not typically designed to carry out all of the Transmission-Based Precautions (e.g., Airborne Precautions for patients with suspected tuberculosis, measles, or chickenpox) that are recommended for hospital and other ambulatory care settings. Patients, however, do not usually seek routine dental outpatient care when acutely ill with diseases requiring Transmission-Based Precautions. Nonetheless, DHCP should develop and carry out systems for early detection and management of potentially infectious patients at initial points of entry to the dental setting. To the extent possible, this includes rescheduling non-urgent dental care until the patient is no longer infectious or referral to a dental setting with appropriate infection prevention precautions when urgent dental treatment is needed. Hand hygiene is the most important measure to prevent the spread of infections among patients and DHCP. Education and training programs should thoroughly address indications and techniques for hand hygiene practices before performing routine and

oral surgical procedures. For routine dental examinations and nonsurgical procedures, use water and plain soap (hand washing), or antimicrobial soap (hand antisepsis) specific for health care settings, or use an alcohol-based hand rub. Although alcohol-based hand rubs are effective for hand hygiene in health care settings, soap and water should be used when hands are visibly soiled (e.g., dirt, blood, body fluids). For surgical procedures, A perform a surgical hand scrub before putting on sterile surgeon's gloves. For all types of hand hygiene products, follow the product manufacturer's label for instructions. Complete guidance on how and when hand hygiene should be performed, including recommendations regarding surgical hand antisepsis and artificial nails, can be found in the Guideline for Hand Hygiene in Health-Care Settings. Personal protective equipment (PPE) refers to wearable equipment that is designed to protect DHCP from exposure to or contact with infectious agents. PPE that is appropriate for various types of patient interactions and effectively covers personal clothing and skin likely to be soiled with blood, saliva, or other potentially infectious materials (OPIM) should be available. These include gloves, face masks, protective eye wear, face shields, and protective clothing (e.g., reusable or disposable gown, jacket, laboratory coat). Examples of appropriate use of PPE for adherence to Standard Precautions include: DHCP should be trained to select and put on appropriate PPE and remove PPE so that the chance for skin or clothing contamination is reduced. Hand hygiene is always the final step after removing and disposing of PPE. Training should also stress preventing further spread of contamination while wearing PPE by: The application of Standard Precautions and guidance on appropriate selection and an example of putting on and removal of personal protective equipment is described in detail in the 2007 Guideline for Isolation Precautions. Respiratory hygiene/cough etiquette infection prevention measures are designed to limit the transmission of respiratory pathogens spread by droplet or airborne routes. The strategies target primarily patients and individuals accompanying patients to the

dental setting who might have undiagnosed transmissible respiratory infections, but also apply to anyone (including DHCP) with signs of illness including cough, congestion, runny nose, or increased production of respiratory secretions. DHCP should be educated on preventing the spread of respiratory pathogens when in contact with symptomatic persons. Respiratory hygiene/cough etiquette measures were added to Standard Precautions in 2007. Additional information related to respiratory hygiene/cough etiquette can be found in the 2007 Guideline for Isolation Precautions. Recommendations for preventing the spread of influenza are available at: [Infection Control in Health Care Facilities | CDC](#). Most percutaneous injuries (e.g., needlestick, cut with a sharp object) among DHCP involve burs, needles, and other sharp instruments. Implementation of the OSHA Bloodborne Pathogens Standard has helped to protect DHCP from blood exposure and sharps injuries. However, sharps injuries continue to occur and pose the risk of bloodborne pathogen transmission to DHCP and patients. Most exposures in dentistry are preventable; therefore, each dental practice should have policies and procedures available addressing sharps safety. DHCP should be aware of the risk of injury whenever sharps are exposed. When using or working around sharp devices, DHCP should take precautions while using sharps, during cleanup, and during disposal. Engineering and work-practice controls are the primary methods to reduce exposures to blood and OPIM from sharp instruments and needles. Whenever possible, engineering controls should be used as the primary method to reduce exposures to bloodborne pathogens. Engineering controls remove or isolate a hazard in the workplace and are frequently technology based (e.g., self-sheathing anesthetic needles, safety scalpels, and needleless IV ports). Employers should involve those DHCP who are directly responsible for patient care (e.g., dentists, hygienists, dental assistants) in identifying, evaluating, and selecting devices with engineered safety features at least annually and as they become available. Other examples of engineering controls include sharps containers and needle recapping devices. When engineering

controls are not available or appropriate, work-practice controls should be used. Work-practice controls are behavior-based and are intended to reduce the risk of blood exposure by changing the way DHCP perform tasks, such as using a one-handed scoop technique for recapping needles between uses and before disposal. Other work-practice controls include not bending or breaking needles before disposal, not passing a syringe with an unsheathed needle by hand, removing burs before disassembling the handpiece from the dental unit, and using instruments in place of fingers for tissue retraction or palpation during suturing and administration of anesthesia. All used disposable syringes and needles, scalpel blades, and other sharp items should be placed in appropriate puncture-resistant containers located close to the area where they are used. Sharps containers should be disposed of according to state and local regulated medical waste rules. For more information about sharps safety, see the Guidelines for Infection Control in Dental Health-Care Settings—2003, the CDC Workbook for Designing, Implementing, and Evaluating a Sharps Injury Prevention Program, and the CDC Sample Screening and Device Evaluation Forms for Dentistry. Safe injection practices are intended to prevent transmission of infectious diseases between one patient and another, or between a patient and DHCP during preparation and administration of parenteral (e.g., intravenous or intramuscular injection) medications. Safe injection practices are a set of measures DHCP should follow to perform injections in the safest possible manner for the protection of patients. DHCP most frequently handle parenteral medications when administering local anesthesia, during which needles and cartridges containing local anesthetics are used for one patient only and the dental cartridge syringe is cleaned and heat sterilized between patients. Other safe practices described here primarily apply to use of parenteral medications combined with fluid infusion systems, such as for patients undergoing conscious sedation. Unsafe practices that have led to patient harm include (1) use of a single syringe—with or without the same needle—to administer medication to multiple

patients, (2) reinsertion of a used syringe—with or without the same needle—into a medication vial or solution container (e.g., saline bag) to obtain additional medication for a single patient and then using that vial or solution container for subsequent patients, and (3) preparation of medications in close proximity to contaminated supplies or equipment. Safe injection practices were covered in the Special Considerations section (Aseptic Technique for Parenteral Medications) of the 2003 CDC dental guidelines. However, because of reports of transmission of infectious diseases by inappropriate handling of injectable medications, CDC now considers safe injection practices to be a formal element of Standard Precautions. Complete guidance on safe injection practices can be found in the 2007 Guideline for Isolation Precautions. Additional materials, including a list of frequently asked questions from providers and a patient notification toolkit, are also available. CDC protects patients and providers by developing recommendations that guide infection prevention and control practices wherever dental care is delivered. Languages Language Assistance Languages Language Assistance

