

Probability & Stats - 2 lectures

Suggestions
on WhatsApp

Remedial - Session
clear confusions

Assumption: gone through regular classes


Next class onwards

- any left-over topics
- regular - topics

} → primary instruction

{ TTS → time being

{ MWF → 4-5 classes



~~Topics:~~ (lots of examples & problems)

- 1. Joint & conditional probability
- 2. Independent events
- 3. Mutually exclusive & exhaustive events
- 4. Bayes Theorem & examples
- 5. Descriptive Stats → ~~Open~~

~~operations~~

1. chat window → end of each topic

↳ context

2. Out of context → Q&A (@ end)

OnMule (Audio) →

3. participate in problem-solving → chat

~~read eqns:~~

$$P(M_1 \text{ and } S) = P(M_1 \cap S) = p(M_1, S)$$

Joint Probability

~~posting (1000)~~ U: Universe of all covid patients


$$\begin{aligned} p(M_1 \cap S) &= 0.95 \\ &= 95/1000 \end{aligned}$$

(100) M_1 : all patients administered medicine - U = 1000

(700) S : patient survived

$$P(M_1) = 0.1 = 10\% = 10/1000$$

$$P(S) = 0.7 = 700/1000$$



100
700

$$M_1 \cap S = 95$$

Joint Prob



2 or more events ^{jointly -} occurring together

{ Set-based Venn-diagrams → A&C
↳ helpful

InterviewBit Software Services x Live | DSML Advanced : Pro x dsml-course/Probability Theor x dsml-course/Probability and St x Popular Interview Question fro x New Tab x +

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

GEO MRTT

Srikanth Varma Chekuri (You)

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

Notify me about Nothing

Saurav Kumar To: everyone 9:15 pm

Pin a message +

keshav kumar To: Me 9:16 pm

intersection of m1 and S

Tai Rakesh To: Me 9:16 pm

$p(m_1) + p(s) - p(m \cap s)$

Prabhakar Patra To: Me 9:16 pm

.07

Tai Rakesh To: Me 9:19 pm

0.095

Adyashree Mahapatra To: Me 9:19 pm

0.095

Tai Rakesh To: Me 9:19 pm

95/1000

Adyashree Mahapatra To: Me 9:19 pm

95/1000

Deepak Mittal To: Me 9:20 pm

$P(M) + P(S) - P(M \cup S)$

Srujanika To: Me 9:20 pm

0

keshav kumar To: Me 9:20 pm

$0.1 * 0.95$

Start Doubt Session Yes No

To: Everyone Enable/Disable Chat

Type message

8 / 8

You are sharing your screen now

Stop Sharing

Srikanth Varma Chekuri (You) (Screen)

00:20:47

8:22 People

Chat

Questions

GEOMRTT

Srikanth Varma Chekuri (You)

Notify me about Nothing

Saurav Kumar To: everyone 9:15 pm

Pin a message +

keshav kumar To: Me 9:16 pm

intersection of m1 and S

Tai Rakesh To: Me 9:16 pm

$p(m_1) + p(s) - p(m \cap s)$

Prabhakar Patra To: Me 9:16 pm

.07

Tai Rakesh To: Me 9:19 pm

0.095

Adyashree Mahapatra To: Me 9:19 pm

0.095

Tai Rakesh To: Me 9:19 pm

95/1000

Adyashree Mahapatra To: Me 9:19 pm

95/1000

Deepak Mittal To: Me 9:20 pm

$P(M) + P(S) - P(M \cup S)$

Srujanika To: Me 9:20 pm

0

keshav kumar To: Me 9:20 pm

$0.1 * 0.95$

Start Doubt Session Yes No

To: Everyone Enable/Disable Chat

Type message

8 / 8

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

GEOMLTT

(Q) Throw a dice $\{1, 2, 3, 4, 5, 6\}$ ← Space

A: even number on a throw $\rightarrow \{2, 4, 6\}$

B: prime number on a throw $\rightarrow \{2, 3, 5\}$

You are sharing your screen now

Stop Sharing

$P(A \text{ and } B) = \frac{1}{6}$

Srikanth Varma Chekuri (You) (Screen)

00:23:45

9 / 9

80 People

Chat

Notify me about Nothing

Pin a message +

1/6

Adyashree Mahapatra To: Me 9:22 pm

1/6

Harini To: Me 9:22 pm

1/6

Saurav Kumar To: Me 9:22 pm

1/6

Harini To: Me 9:22 pm

2 is the only prime and even number

keshav kumar To: Me 9:22 pm

1/2 * 1/3

Tai Rakesh To: Me 9:22 pm

2 is the common element out of 6

Srinivas MV To: Me 9:23 pm

1/6 because only 2 is both prime and even

Prabhakar Patra To: Me 9:23 pm

1/6

Adyashree Mahapatra To: Me 9:23 pm

A: even number: 2, 4, 6
B: prime: 2, 3, 5
A and B: 2

Start Doubt Session

To: Everyone

Enable/Disable Chat

Type message

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

(Q) Throw a dice $\rightarrow \{1, 2, 3, 4, 5, 6\}$

A : even number $\{2, 4, 6\}$

B : prime number $\{2, 3, 5\}$

C : NOT $\{2, 3, 4\}$ $\{1, 5, 6\}$

$P(A \text{ and } B \text{ and } C)$

$$= \frac{0}{6} = 0$$

You are sharing your screen now
Stop Sharing

Srikanth Varma Chekuri (You) (Screen)

00:26:20

86

24 People

Chat

Notify me about Nothing

Pin a message +

mohan To: Me 9:25 pm

Pooja Ayanile To: Me 9:25 pm

Prabhakar Patra To: Me 9:25 pm

Aayushi Jain To: Me 9:25 pm

keshav kumar To: Me 9:25 pm

Harini To: Me 9:25 pm

0 - nothing is common in the sets A: {2,4,6}
B: {2,3,5}
C: {1,5,6}

Pooja Ayanile To: Me 9:25 pm
nothing is satisfy all a,b,c

Srujanika To: Me 9:25 pm

Srujanika To: Me 9:26 pm
no

Start Doubt Session Yes No

To: Everyone Enable/Disable Chat

Type message

10 / 10



$$P(S|M_1) =$$


Same covid example:

M_1 : patient admin M_1

S : patient survived

(Q) probability of survival
a patient was admin. M_1 given that
 $\underbrace{M_1}_{\text{conditioned}} \rightarrow P(S|M_1)$

Conditional prob:



as
Notation

$$P(S|M_1)$$

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

NOT $\frac{5}{100}$

$P(S|M_1) = \frac{95}{100}$

$\frac{P(M_1 \cap S)}{P(M_1)}$; $U = 1000$

95 out of 100 patients who have been admin... MI survived

Visually

You are sharing your screen now

Stop Sharing

95/100 as 95 people are survived out of 100

5 out of 100 survived with M1 so 5 %

$p(s|and\ m1)/p(m1)=95/100=0.95$

95 out of 100 patients given medicine are survived.

got it

To: Everyone | Enable/Disable Chat

Type message

Srikanth Varma Chekuri (You)

Chat

Notify me about: Nothing

Pin a message

Srinivas MV To: Me 9:29 pm

Tai Rakesh To: Me 9:31 pm

mohan To: Me 9:31 pm

SAVARA KALYAN To: Everyone 9:31 pm

.95

Srujanika To: Me 9:31 pm

Srinivas MV To: Me 9:31 pm

Adyashree Mahapatra To: Me 9:32 pm

Srinivas MV To: Me 9:33 pm

Start Doubt Session

00:34:41

12 / 12

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

$P(S|M_1) = \frac{P(S \cap M_1)}{P(M_1)}$

if $P(M_1) \neq 0$

You are sharing your screen now

Stop Sharing

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

Pin a message

Srinivas MV To: Me 9:31 pm
5 out of 100 survived with M1 so 5 %

Srujanika To: Me 9:31 pm
 $p(\text{sand } m_1)/p(m_1) = 95/100 = 0.95$

Adyashree Mahapatra To: Me 9:32 pm
95 out of 100 patients given medicine are survived.

9:33 pm 3

Srinivas MV To: Me 9:33 pm
got it

Tai Rakesh To: Me 9:35 pm
that means someone has take medicine

Shashank Upadhyay To: Me 9:36 pm
so $P(M_1|S)$ will be
 $P(M_1 \cap S)/P(S)$

Tai Rakesh To: Me 9:36 pm
from the example, that means someone has to take medicine

Tai Rakesh To: Me 9:36 pm
yes

Start Doubt Session

To: Everyone Enable/Disable Chat

Type message

GEOMLTT Srikanth Varma Chekuri (You) (Screen)

00:36:22

13 / 14

Some one has to take the medicine 1

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

$p(M_1|S) = \frac{95}{100} = \frac{p(M_1 \cap S)}{p(S)}$ if $p(S) \neq 0$

You are sharing your screen now
Stop Sharing

U (1000)

M₁ S

100 95 100

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

Pin a message +

Srujana Kata To: Me 9:31 pm p(sand m1)/p(m1)=95/100=0.95

Adyashree Mahapatra To: Me 9:32 pm 95 out of 100 patients given medicine are survived.

9:33 pm 3

Srinivas MV To: Me 9:33 pm got it

Tai Rakesh To: Me 9:35 pm that means someone has take medicine

Shashank Upadhyay To: Me 9:36 pm so $p(M_1|S)$ will be $p(M_1 \cap S)/p(S)$

Tai Rakesh To: Me 9:36 pm from the example, that means someone has to take medicine

Tai Rakesh To: Me 9:36 pm yes

keshav kumar To: Me 9:37 pm 95/100

Start Doubt Session Yes No

To: Everyone Enable/Disable Chat

Type message

00:38:05 14 / 15

(Q) Two distinct dice

(Q) prob that sum of two dice equals 7

given that D_1 has shown value 3

[define events properly]

$P(E|F)$

D_1 D_2

$E \cap F$

You are sharing your screen now
Stop Sharing

Srikanth Varma Chekuri (You) (Screen)
00:42:42

Scalera.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

80

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

Pin a message

when D_1 is 3, only option of D_2 is 4. so out of all possibilities of combinations it is 1/36

Adyashree Mahapatra To: Me 9:41 pm

Dice is 3 is fixed. So, probability of getting 4 to make sum is 1/6.

keshav kumar To: Me 9:41 pm

D_1 has 3 already. that makes sample space as all variants on D_2 . and outcome is just 4 on D_2

Srujanika Kata To: Me 9:41 pm

only combination for 3 to get 7 is 4, options on D_2 is 6 so 1/6

Saurav Kumar To: Me 9:41 pm

because the first dice is fixed, so the second dice probability is 1/6

Shashank Upadhyay To: Me 9:41 pm

prob of getting 3 on first dice is 1/6 and sum should be 7 so dice 2 should have 4 for ans = (1/6)(1/6)

Rahul Singh To: Me 9:42 pm

as in event1 = 1/6 as {3} out of {1..6}, event2 = {4} out of {1..6}

Start Doubt Session

To: Everyone Enable/Disable Chat

Type message

15 / 16

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

E: SUM of 2 dice is 7

F: D₁ results in 3

$P(E|F) = \frac{1}{6}$

(intuitive) ✓

You are sharing your screen now
Stop Sharing

D₁ D₂

3 4

1 → 4
2 → 5
3 → 6
4 → 7 ✓
5 → 8
6 → 9

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

Advaithree Mahantra To: Me 9:41 pm Pin a message +
to make sum is 1/6.

keshav kumar To: Me 9:41 pm dice 1 has 3 already. that makes sample space as all variants on dice 2. and outcome is just 4 on dice 2

Srujanika Kata To: Me 9:41 pm only combination for 3 to get 7 is 4 , options on dice 2 is 6 so 1/6

Saurav Kumar To: Me 9:41 pm because the first dice is fixed, so the second dice probability is 1/6

Shashank Upadhyay To: Me 9:41 pm prob of getting 3 on frst dice is 1/6 and sum should be 7 so dice 2 should have 4 for ans= (1/6)(1/6)

Rahul Singh To: Me 9:42 pm as in event1 = 1/6 as {3} out of {1..6} , event2 = {4} out of {1..6}

9:43 pm 5

Srinivas MV To: Me 9:43 pm if D1 is always 3, it is 1/6

Start Doubt Session

To: Everyone Yes No

Enable/Disable Chat

Type message

GEOMLRTT Srikanth Varma Chekuri (You) (Screen)

00:46:24

16 / 18

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

Formula

$$P(E|F) = \frac{P(E \cap F)}{P(F)}$$

SUM = 7
D₂ = 4
and D₁ = 3

$$= \frac{1/36}{1/4} = \boxed{\frac{1}{6}}$$

You are sharing your screen now

Stop Sharing

D₁ D₂

$\left\{ (1,1), (1,2), (1,3), \dots \right.$
 $\left. (2,1), (2,2), \dots \right.$

Srikanth Varma Chekuri (You) (Screen)

00:46:30

80% 90% 100%

17 / 18



Start Doubt Session

To: Everyone

Enable/Disable Chat

Type message

25 People

Chat

Notify me about: Nothing

Pin a message

to make sum is 1/6.

keshav kumar To: Me 9:41 pm

dice 1 has 3 already. that makes sample space as all variants on dice 2. and outcome is just 4 on dice 2

Srujanika To: Me 9:41 pm

only combination for 3 to get 7 is 4 , options on dice 2 is 6 so 1/6

Saurav Kumar To: Me 9:41 pm

because the first dice is fixed, so the second dice probability is 1/6

Shashank Upadhyay To: Me 9:41 pm

prob of getting 3 on frst dice is 1/6 and sum should be 7 so dice 2 should have 4 for ans= (1/6)(1/6)

Rahul Singh To: Me 9:42 pm


as in event1 = 1/6 as {3} out of {1..6} , event2 = {4} out of {1..6}

9:43 pm

Srinivas MV To: Me 9:43 pm

if D1 is always 3, it is 1/6

Yes No



$$P(S|M_1) = 0.7 = P(S)$$

↓
patient survival $\equiv (S)$

did not depend on M_1


↓
 $M_1 \text{ & } S_2$ are independent events

Common-
Confusion

Independent
events

$M_1 \cap S \neq \emptyset$
need not
empty set

$$P(M_1 \cap S) \neq 0$$



$$P(S|M_1) = P(S)$$

Mutually exclusive
events

e.g: $S : \text{Survived}$
 $D : \text{dead}$

$$S \cap D = \emptyset$$

$$P(S \cap D) = 0$$



InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

$P(S|M_1) = P(S)$

M1 does not Impact
the survival of
the patient

You are sharing your screen now

Stop Sharing

Srikanth Varma Chekuri (You) (Screen)

00:56:31

80 25 People

Chat

Notify me about Nothing

Pin a message +

Rahul Singh To: Me 9:48 pm

Shashank Upadhyay To: Me 9:48 pm

Srinivas MV To: Me 9:48 pm

700/1000

Harini To: Me 9:49 pm

700/1000

Saurav Kumar To: Me 9:49 pm

700/1000

keshav kumar To: Everyone 9:51 pm

can we say, medicine does not cause survival or will that be a wrong statement? in strictest sense.

Harini To: Me 9:54 pm

So all MEE are independent events but not all independent events are MEE?

9:55 pm 4

9:56 pm 1

Start Doubt Session

To: Everyone Enable/Disable Chat

Type message

21 / 21

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

GEOMLTT

Mut. excl. events

$S \cap D = \emptyset$

Indep - $P(A|B) = P(A)$

$P(S|D) \neq P(S)$

$P(S \cap D) = 0$

$P(D)$

You are sharing your screen now

Stop Sharing

Srikanth Varma Chekuri (You) (Screen)

01:04:48

22 / 24

80 People

Chat

Notify me about Nothing

Pin a message + need not be but can be ? 9:59 pm

1

keshav kumar To: Everyone 10:02 pm

ok

Safwan To: Everyone 10:02 pm

in this case isn't it there has to be 2 dice or 2 throw

Safwan To: Everyone 10:02 pm

?

10:02 pm

1

Safwan To: Everyone 10:02 pm

but given B means

Safwan To: Everyone 10:02 pm

already already happened?

Shashank Upadhyay To: Me 10:03 pm

i got confused between MEE and independent event

can you please explain again if possible

10:04 pm

1

Start Doubt Session

To: Everyone

Enable/Disable Chat

Type message

Yes No

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

dice

even = {A: even number {2, 4, 6}, B: prime number {2, 3, 5}}

You are sharing your screen now
Stop Sharing

$P(A) = 3$

$P(A|B) = \frac{P(A \cap B)}{P(B)}$

$= \frac{1}{6}$

$\frac{1}{2}$

$= \frac{1}{3}$

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

Pin a message

Aayushi Jain To: Me 9:59 pm need not be but can be ? 9:59 pm 1

keshav kumar To: Everyone 10:02 pm ok

Safwan To: Everyone 10:02 pm in this case isn't it there has to be 2 dice or 2 throw

Safwan To: Everyone 10:02 pm ? 10:02 pm 1

Safwan To: Everyone 10:02 pm but given B means

Safwan To: Everyone 10:02 pm already already happened?

Shashank Upadhyay To: Me 10:03 pm i got confused between MEE and independent event can you please explain again if possible

Start Doubt Session

Yes No

To: Everyone Enable/Disable Chat

Type message

01:03:28 23 / 24

~~DSML Advanced : Probability & Statistics Remedial -1 | Lecture~~

$P(S \cup D \cup C) = P(U) = \frac{1000}{1000} = 1$ Mutually exclusive and exhaustive events

✓ S: survived
D: dead
C: medical care currently

$U = 1000$

$S \cup D \cup C$

$S \cap D = \emptyset$

$S \cap C = \emptyset$

$D \cap C = \emptyset$

$SUDUC = U$

Geometric representation:

Chat Log:

- 1 Saurav Kumar To: Me 10:09 pm 0
- 0 Praveen Galagali To: Everyone 10:09 pm 1?
- 1 Aayushi Jain To: Me 10:09 pm 1
- 1 Tai Rakesh To: Me 10:09 pm $p(s)+d+c$
- 1 Saurav Kumar To: Me 10:09 pm 1
- 0 Saurav Kumar To: Me 10:09 pm typo
- 1 Tai Rakesh To: Me 10:09 pm 1 total max probability
- 0 Praveen Galagali To: Everyone 10:10 pm one of the event will occur for sure
- 0 Deepak Mittal To: Me 10:10 pm event D intersection event C should be phi in the previous page

Start Doubt Session

To: Everyone Enable/Disable Chat

Type message

01:11:19 24 / 25

personal
✓

$$\left\{ \begin{array}{l} P(A|B) = P(A) \\ \quad \quad \quad \end{array} \right.$$

→ independence
derived

$$P(A \cap B) = P(A|B) P(B)$$

$$\left[\because P(A|B) = \frac{P(A \cap B)}{P(B)} \right]$$

✓ $P(A \cap B) = P(A) P(B)$ → independence

Q

n

coins that are tossed

E_i : event that ith coin is a head = $\frac{1}{2}$

$$\frac{1}{2} \quad \frac{1}{2} \quad \frac{1}{2} \quad \frac{1}{2} \quad \dots \frac{1}{2}$$

prob of all n-coins turning heads?

$$= \frac{1}{2^n}$$

① One case out of 2^n = $\frac{1}{2^n}$

$$\rightarrow ② P(\underline{E_1} \cap \underline{E_2} \cap \underline{E_3} \cap \dots \cap \underline{E_n}) = \underline{\frac{1}{2}} \underline{\frac{1}{2}} \underline{\frac{1}{2}} \dots \underline{\frac{1}{2}} = \frac{1}{2^n}$$

(Q) Given: M_1 and S are independent events $\Rightarrow P(S|M_1) = P(S)$

Can we say: M_1^C and S are also independent?

NOT M_1

$P(S|M_1^C) = P(S)$

$$\frac{630}{900} = 0.7 = \frac{700}{1000}$$

You are sharing your screen now
Stop Sharing

Praveen Galagali To: Me 10:22 pm
 $P(S) = 70/1000$ but $P(M_1 \text{ complement}) = 630/1000$. SO they are not equal?

Adyashree Mahapatra To: Me 10:22 pm
 $P(S) - P(M_1 \text{ and } S) / P(U - m_1) = 630/900$

Srinivas MV To: Me 10:22 pm
in the same diagram, if M_1 grows, S has to grow the same way to maintain independence. So, other way around would mean compliment is also independent

Praveen Galagali To: Me 10:22 pm
700*

keshav kumar To: Me 10:22 pm
 $m' \text{ and } s = (700-70)/900$

Tai Rakesh To: Everyone 10:23 pm
630

Tai Rakesh To: Everyone 10:23 pm
900

Adyashree Mahapatra To: Me 10:23 pm
0.7

Start Doubt Session

To: Everyone Enable/Disable Chat

Type message

01:24:04 27 / 27

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

(Q) Expt: Toss a coin and throw a dice = Yes

$\frac{1}{2}$ E: the coin is H

$\frac{1}{6}$ F: the dice is 3

You are sharing your screen now
Stop Sharing

Q1: are E & F indep?

Q2: are E & F mutually exclusive

if $P(E|F) = P(E)$

$$= \frac{P(E \cap F)}{P(F)} = \frac{\frac{1}{2} \cdot \frac{1}{4}}{\frac{1}{6}} = \frac{\frac{1}{8}}{\frac{1}{6}} = \frac{1}{2}$$

$E \cap F = \emptyset$

Srikanth Varma Chekuri (You) (Screen)
01:32:48

86 People

Chat

Notify me about Nothing

Q1 user Q2 user

Pin a message +

not mutually exclusive

mohan To: Me 10:27 pm yes, no

Adyashree Mahapatra To: Me 10:28 pm $P(H \text{ and } 3)/P(3) = 0$ which is not equal to $P(H) = 1/2$

Saurav Kumar To: Me 10:28 pm $1/2 * 1/6 = 1/12$

Prabhakar Patra To: Me 10:29 pm 1/12

Adyashree Mahapatra To: Me 10:31 pm but we can take $P(E \text{ and } F) = P(E) * P(F)$ only when we assume that they are independent. So, how we take this value to prove independence? 10:31 pm 5

Shashank Upadhyay To: Me 10:32 pm if E1: dice show even number E2: dice show odd number so can we say then E1 and E2 are MEE

Start Doubt Session

To: Everyone Enable/Disable Chat

Type message

28 / 29

InterviewBit Software Services x Live | DSML Advanced : Pro x dsml-course/Probability Theor x dsml-course/Probability and St x Popular Interview Question fro x New Tab x +

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

event e does not impact occurrence of event

Pin a message +

Safwan To: Everyone 10:27 pm

Q2 no mutually exclusive

Srinivas MV To: Me 10:27 pm

Q1:yes : Q2: no

Tim Sabu To: Me 10:27 pm

not mutually exclusive

mohan To: Me 10:27 pm

yes, no

Adyashree Mahapatra To: Me 10:28 pm

$P(H \text{ and } 3)/P(3) = 0$ which is not equal to
 $P(H) = 1/2$

Saurav Kumar To: Me 10:28 pm

$1/2 * 1/6 = 1/12$

Prabhakar Patra To: Me 10:29 pm

$1/12$

Adyashree Mahapatra To: Me 10:31 pm

but we can take $P(E \text{ and } F) = P(E) * P(F)$ only when we assume that they are independent. So, how we take this value to prove independence?

10:31 pm

Start Doubt Session

GEO MRTT

Srikanth Varma Chekuri (You) (Screen)

01:31:18

29 / 29

People

Chat

Questions

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

1,2,3,4,5,6

E1: dice shows even num

E2: dice shows odd num

$E_1 \cap E_2 = \emptyset$

You are sharing your screen now

Stop Sharing

mutually exclusive

Srikanth Varma Chekuri (You) (Screen)

01:34:00

30 / 30

8:00

GEOMRTT

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

Pin a message

P(H and 3)/P(3) = 0 which is not equal to P(H)=1/2

Saurav Kumar To: Me 10:28 pm
1/2*1/6=1/12

Prabhakar Patra To: Me 10:29 pm
1/12

Adyashree Mahapatra To: Me 10:31 pm
but we can take P(E and F) = P(E) *P(F) only when we assume that they are independent. So, how we take this value to prove independence?

10:31 pm
5

Shashank Upadhyay To: Me 10:32 pm
if
E1: dice show even number
E2: dice show odd number
so can we say
then E1 and E2 are MEE

10:33 pm
1

Tai Rakesh To: Everyone 10:33 pm
are all mutual exclusive events are independent

Start Doubt Session

To: Everyone Enable/Disable Chat

Type message

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

-if $A \& B$ are MEE

then $A \& B$ are indep

given $A \cap B = \emptyset$

$P(A|B) = P(A)$

$P(A \cap B) = P(A) P(B)$

LHS : 0
need not be 0
always

You are sharing your screen now
Stop Sharing

Srikanth Varma Chekuri (You) (Screen)
01:37:03

80 GEOMRTT Srikanth Varma Chekuri (You)

Chat Notify me about Nothing Pin a message + Saurav Kumar To: Me 10:28 pm 1/2 * 1/6 = 1/12 Prabhakar Patra To: Me 10:29 pm 1/12 Adyashree Mahapatra To: Me 10:31 pm but we can take $P(E \text{ and } F) = P(E) * P(F)$ only when we assume that they are independent. So, how we take this value to prove independence? 10:31 pm 5 Shashank Upadhyay To: Me 10:32 pm if E1: dice show even number E2: dice show odd number so can we say then E1 and E2 are MEE 10:33 pm 1 Tai Rakesh To: Everyone 10:33 pm are all mutual exclusive events are independent Shashank Upadhyay To: Me 10:34 pm yes Start Doubt Session Yes No To: Everyone Enable/Disable Chat Type message

(Q) A and B are two events ; which of the following are true

MSQ

~~a~~ $P(A \cap B) = P(A) P(B)$ → indep $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

~~b~~ $P(A \cup B) = P(A) + P(B)$ → MET $A \cap B = \emptyset$ 

~~c~~ $P(A|B) = P(A \cap B) / P(B)$ \times

~~d~~ $P(A \cup B) \leq P(A) + P(B)$ ✓

$A \cup B = A + B - A \cap B$

Scalera.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

Pin a message

SAVARA KALYAN To: Me 10:39 pm d

Tai Rakesh To: Everyone 10:40 pm d

Harini To: Me 10:40 pm only d since we don't know anything else about a and b

Saurav Kumar To: Me 10:40 pm d

Srujanika To: Me 10:40 pm none

keshav kumar To: Me 10:41 pm d alone

Shashank Upadhyay To: Me 10:41 pm d

Srujanika To: Me 10:44 pm how can union less than union B 10:44 pm 4

Srujanika To: Me 10:44 pm typo

Start Doubt Session Yes  No 

To: Everyone Enable/Disable Chat

Type message

01:44:31 32 / 32

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

GEMRTT

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

Pin a message +

TarKakesh To: Everyone 10:40 pm d

Harini To: Me 10:40 pm only d since we don't know anything else about a and b

Saurav Kumar To: Me 10:40 pm d

Srujanika To: Me 10:40 pm none

keshav kumar To: Me 10:41 pm d alone

Shashank Upadhyay To: Me 10:41 pm d

Srujanika To: Me 10:44 pm how can union less than Aunion B 10:44 pm 4

Srujanika To: Me 10:44 pm typo

Srujanika To: Me 10:44 pm how can union less tha a +B


Start Doubt Session Yes No

To: Everyone Enable/Disable Chat

Type message

01:46:57 34 / 34

A \cup B = A + B - A \cap B



You are sharing your screen now

Stop Sharing

InterviewBit Software Services | Live | DSML Advanced : Pr... | dsml-course/Probability Theor... | dsml-course/Probability and St... | Popular Interview Question fro... | New Tab

scaler.com/meetings/i/dsml-advanced-probability-statistics-remedial-1/live

DSML Advanced : Probability & Statistics Remedial -1 | Lecture

(Q)

given $P(A) = 1 \Rightarrow U = A$; $P(B) = 1/2$

$P(A|B)$

a 1

b $P(B|A)$

You are sharing your screen now
Stop Sharing

$P(B) = 1/2$

$U = A$

Srikanth Varma Chekuri (You)

Chat

Notify me about Nothing

Pin a message

Tai Rakesh To: Me 10:48 pm
 $p(b/a)=1/2**$

Srujanika To: Me 10:49 pm
not possible until u mention relation between A and B

Safwan To: Everyone 10:49 pm
a = 1 and b = 1/2 as its independent events

Tai Rakesh To: Me 10:50 pm
 $p(A/B)=1$ A occurs compulsory irrespective of B
 $p(B/a)=1/2$ A occurs compulsory all the time

keshav kumar To: Me 10:50 pm
a=1, b = 1/2 . in case a, b has already occurred and event a will always occur hence a is 1.

Harini To: Me 10:50 pm
since $p(A) = 1$ it means it's a sure event. So $P(A/B)$ is also 1 and $p(B/A) = 1/2$

keshav kumar To: Me 10:51 pm
in case of b, event a has already occurred and its sure event hence occurrence of event B is independent of event A

Start Doubt Session



Yes No

To: Everyone Enable/Disable Chat

Type message

01:51:35

35 / 35



Bayes Thm

$$p(\tilde{A}|B) = \frac{p(\tilde{B}|A) p(\bar{A})}{p(B)}$$

if $p(B) \neq 0$




cond. prob

Proof:



$$P(\tilde{A}|\tilde{B}) = \frac{\cancel{P(A \cap B)}}{P(B)} = \frac{\cancel{P(B \cap A)}}{P(B)} = \frac{\cancel{P(B|A) P(A)}}{P(B)}$$

If $P(B) \neq 0$

Real-world



Let's assume ✓ IV. of the population has covid



✓ What is the prob that actually has covid?


have covid
rapid RT-PCR +ve

$$P(C|+ve) = ?$$


Bayes Thm

$$\frac{P(+ve|C) P(C)}{P(+ve)}$$

$$\rightarrow ? = 0.108$$



$\tilde{P}(+ve) = ?$



$$\left\{ \begin{aligned} P(\tilde{+ve}) &= P(\tilde{+ve} \cap C) + P(\tilde{+ve} \cap NC) \\ &= P(+ve|C)p(C) + P(+ve|NC)p(NC) \\ &= 0.9 \times 0.01 + 0.1 \times 0.99 = 0.108 \end{aligned} \right.$$

You have left the meeting

We get frequent requests for your notes!

Notes written by you helps in understanding the topic better. You can upload the notes in two simple steps mentioned below

- 1 Scan the QR code with your iPad
Scanner should be present in the top menu on your iPad
 - 2 Upload Notes on the generated link
All notes uploaded will be visible in the saved version of this session



OR

Drag and drop files or click here to upload

Files Uploaded from your computer appear here