

Seismic Quantities

pySYD pipeline

Information that we can obtain

- In this study they look for Solar-like oscillations in a star's power spectrum ($\leq 20 \mu\text{Hz}$)
- They are characterized by two global seismic quantities:

ν_{max} the frequency of the maximum acoustic power,

and

$\Delta\nu$ the large frequency spacing between adjacent overtone oscillation modes.

$$\nu_{max} \propto g T_{eff}^{-1/2}$$

$$\Delta\nu \propto \rho^{-1/2}$$

$$\left(\frac{M}{M_{\odot}}\right) \approx \left(\frac{\nu_{max}}{\nu_{max,\odot}}\right)^3 \left(\frac{\Delta\nu}{\Delta\nu_{\odot}}\right)^{-4} \left(\frac{T_{eff}}{T_{eff,\odot}}\right)^{3/2} \quad (1)$$

$$\left(\frac{M}{M_{\odot}}\right) \approx \left(\frac{\Delta\nu}{\Delta\nu_{\odot}}\right)^2 \left(\frac{L}{L_{\odot}}\right)^{3/2} \left(\frac{T_{eff}}{T_{eff,\odot}}\right)^{-6} \quad (2)$$

$$\left(\frac{M}{M_{\odot}}\right) \approx \left(\frac{\nu_{max}}{\nu_{max,\odot}}\right) \left(\frac{L}{L_{\odot}}\right) \left(\frac{T_{eff}}{T_{eff,\odot}}\right)^{-7/2} \quad (3)$$

$$\left(\frac{M}{M_{\odot}}\right) \approx \left(\frac{\nu_{max}}{\nu_{max,\odot}}\right)^{12/5} \left(\frac{\Delta\nu}{\Delta\nu_{\odot}}\right)^{-14/5} \left(\frac{L}{L_{\odot}}\right)^{3/10} \quad (4)$$

Seismic Quantities

pySYD pipeline

Information that we can obtain

- In this study they look for Solar-like oscillations in a star's power spectrum ($\leq 20 \mu\text{Hz}$)
- They characterized by two globaleismic quantities:

ν_{max} the frequency of the maximum acoustic power,

and

$\Delta\nu$ the large frequency spacing between adjacent overtone oscillation modes.

$$\nu_{max} \propto g T_{eff}^{-1/2}$$

$$\Delta\nu \propto \rho^{-1/2}$$

