

Prebiotic effect of dietary polyphenols: A systematic review

Aline Medeiros Alves-Santos, Clara Sandra Araújo Sugizaki, Glaucia Carielo Lima, Maria Margareth Veloso Naves*

School of Nutrition, Federal University of Goiás (UFG), Goiânia, Goiás, Brazil

ARTICLE INFO

Keywords:

Gut microbiota
Health
Beneficial bacteria
Butyric acid
Phenolic compounds
Metabolites

ABSTRACT

Prebiotics are substrates selectively metabolized by hindgut microorganisms conferring health benefits. Recent studies suggest polyphenols as candidate to prebiotics. Thus, this systematic review aimed to investigate the prebiotic effect of dietary polyphenols in preclinical and clinical studies. Animal studies demonstrated that the consumption of polyphenols, especially catechins, anthocyanins and proanthocyanidins, increases the abundance of *Lactobacillus*, *Bifidobacterium*, *Akkermansia*, *Roseburia*, and *Faecalibacterium* spp. Moreover, polyphenols supplementation increased the production of short-chain fatty acids (SCFA), including butyrate. The included clinical trials showed an increased abundance of *Lactobacillus acidophilus*, *Bifidobacterium* and *Faecalibacterium* spp., and a reduction in plasma lipopolysaccharide-binding protein after the consumption of anthocyanins and ellagic acid. In conclusion, there is strong evidence in preclinical studies that dietary polyphenols can stimulate both the growth of microorganisms identified as prebiotic targets and an increase in the production of SCFA. Therefore, clinical trials are warranted to investigate the prebiotic effect of dietary polyphenols on humans.

1. Introduction

Prebiotics are defined as substrates selectively utilized by the host's microorganisms resulting in benefits for metabolic health, gastrointestinal system, bone health and mental health. Some dietary fibers, especially resistant oligosaccharides (inulin, fructo-oligosaccharides and galacto-oligosaccharides), are well-recognized in literature as prebiotics (Gibson et al., 2017). Besides dietary fibers, recent studies have shown the interaction between polyphenols and the gut microbiota, suggesting them as candidate compounds to prebiotics (Sanders, Merenstein, Reid, Gibson, & Rastall, 2019; Short et al., 2018; Singh, Cabral, Kumar, Ganguly, & Pandey, 2019).

Polyphenols are secondary metabolites of plants, characterized by aromatic rings bearing one or more hydroxyl groups in their chemical structure, ranging from that of a simple phenolic molecule to that of a complex high-molecular mass polymer (Mojzer et al., 2016). These compounds have low bioavailability and extensive metabolism in the large intestine, favoring interactions with intestinal microorganisms (Bian, Wei, Zhao, & Li, 2020). Actually, there is a bidirectional interaction, in which polyphenols modulate the gut microbiota and, conversely, microorganisms can modulate the activity of the phenolic compounds. This interaction can regulate the metabolism and the bioavailability of polyphenols, converting them into metabolites, which may have different effects on the host health (Singh et al., 2019).

Dietary polyphenols are associated with a reduced risk of cardio-metabolic diseases when regularly consumed (Noad et al., 2016). Studies show that polyphenols have antioxidant, anti-inflammatory (Chai, Davis, Zhang, Zha, & Kirschner, 2019), anti-obesogenic, antilipidemic (Fang et al., 2019) and anti-diabetic (Paquette et al., 2017) activities. However, the role of dietary polyphenols in health largely depends on their metabolism, absorption and bioavailability processes which are, in turn, related to the gut microbiota modulation, in terms of composition and functionality. Although polyphenols are currently recognized as modulators of the gut microbiota composition, there is still no conclusive evidence of their prebiotic effect (Cueva, Silva, Pinillos, Bartolomé, & Moreno-Arribas, 2020). The prebiotic effect of each polyphenol can be influenced by the food source and the chemical structure of the compound, along with the individual differences in the gut microbiota composition (Serreli & Deiana, 2019). Therefore, establishing a relationship among the polyphenols consumption, the growth of microorganisms recognized as prebiotic targets, the metabolites generated and the effect on health is somewhat of a challenge. This systematic review aimed to investigate the available evidence of the prebiotic effect of dietary polyphenols.

* Corresponding author at: School of Nutrition, UFG, Rua 227, Setor Leste Universitário, 74605-080 Goiânia, Goiás, Brazil.

E-mail addresses: alinemedeiros@ufg.br (A.M. Alves-Santos), maria_margareth_veloso@ufg.br (M.M.V. Naves).

<https://doi.org/10.1016/j.jff.2020.104169>

Received 29 June 2020; Received in revised form 17 August 2020; Accepted 19 August 2020

Available online 21 September 2020

1756-4646/ © 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1
Detailed search terms in the databases (August 2, 2019).

Database	Search term	N° of items found (total = 4025)
PubMed	(polyphenols[MeSH Terms] or flavonoids [MeSH Terms] or tannins [MeSH Terms] or lignans [MeSH Terms] or stilbenes [MeSH Terms] or curcumin [MeSH Terms] or "phenolic acids" [Title/Abstract]) AND (microbiota [MeSH Terms] or "human microbiome" [MeSH Terms] or "gut microbiota" [Title/Abstract] or "gut microbiome" [Title/Abstract] or prebiotic [MeSH Terms] or firmicutes [Title/Abstract] or bacteroidetes [MeSH Terms])	796
Scopus	(TITLE-ABS-KEY ((polyphenols OR flavonoids OR tannins OR lignans OR stilbenes OR curcumin OR "phenolic acids"))) AND TITLE-ABS-KEY ((microbiota OR "human microbiome" OR "gut microbiota" OR "gut microbiome" OR prebiotic OR firmicutes OR bacteroidetes)))	1633
Web of Science	#1 TS=(polyphenols OR flavonoids OR tannins OR lignans OR stilbenes OR curcumin OR "phenolic acids") AND #2 TS=(microbiota OR "human microbiome" OR "gut microbiota" OR "gut microbiome" OR prebiotic OR firmicutes OR bacteroidetes)	1596

2. Material and methods

2.1. Search strategy and study selection

In accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Moher et al., 2015), a systematic literature search was carried out through the PubMed, Scopus and Web of Science databases up to August 02, 2019, using terms related to prebiotic, gut microbiota and polyphenols (Table 1). After the removal of duplicates using a bibliographic reference manager, two authors independently reviewed the titles and abstracts of each paper identified in the search. This procedure was performed using a systematic review software. The selected studies were retrieved for full-text analysis and eligible articles were identified. Afterwards, they were reanalyzed for details of the study design and their outcome to select the papers included in this review. Any disagreements in this regard were solved by the means of discussion with a third author.

2.2. Inclusion and exclusion criteria

The following criteria were applied for inclusion: (1) a randomized controlled study design using animals and/or humans; (2) intervention with either isolated polyphenol or food extract; (3) phenolic compounds profile of the food extract available; (4) outcomes related to changes in the gut microbiota composition, with stimulation of microorganisms recognized as prebiotic targets (*Lactobacillus*, *Bifidobacterium*, *Roseburia*, *Eubacterium*, and *Faecalibacterium* spp.) (Gibson et al., 2017), including the *Akkermansia* spp. (Sanders et al., 2019), and/or an increase in the production of short-chain fatty acids (SCFA), including butyrate; and (5) publishing in English. Studies with a lower than eight number of animals per group, acute studies, drug-induced pathologies, gut microbiota composition assessment by culture plate methods, and without the microorganism genera identification were excluded.

2.3. Data extraction

The included studies were reviewed and the following data were abstracted: sample source, exposure dosage, animal model or study population (subjects and sample size), intervention period, diet type, outcomes related to gut microbiota and host health, paper's first author and date of publication.

2.4. Quality assessment

The quality assessment of animal studies was conducted using the SYRCLE's risk of bias tool (Hooijmans et al., 2014). Cochrane Collaboration's tool (Higgins et al., 2011) was used for the quality assessment of clinical trials.

3. Results

3.1. Description of studies

The initial search strategy yielded 4025 articles, out of which 1685 were excluded as duplicates. After analysis of titles and abstracts, 2044 records were excluded, including *in vitro* studies, review articles, conference abstracts, letters, protocols, editorials, and unavailable full-texts, and 296 were selected for full-text review. Among these, 242 studies were excluded according to the exclusion criteria. Thus, full-text review of 54 eligible studies were made, excluding 30 articles for different reasons: there was no growth of microorganisms recognized as prebiotic targets (Casanova-Martí et al., 2018; Cheng, Chen, Zhang, et al., 2019; Cires et al., 2019; Collins et al., 2016; Cowan et al., 2014; González-Sarriás et al., 2017; Griffin et al., 2017; Guo, Tang, et al., 2018; Li, Liu, Liu, Liao, & Zou, 2019; Most, Penders, Lucchesi, Goossens, & Blaak, 2017; Porras et al., 2017; Remely et al., 2017; Shen, Wan, Wang, & Jiang, 2019; Sung et al., 2017; Tan et al., 2018; Unno & Osakabe, 2018; Wang et al., 2019; Yu et al., 2019; Yuan et al., 2018; Zhang, Dong, et al., Zhang, Zhang, et al., 2018; Zhang, Wu, Li, Xin, & Liu, 2019; Zhou, Zhang, Arikawa, & Chen, 2019); there was no increase in the production of butyrate (Giné et al., 2019; Grzelak-Błaszczak et al., 2018; Zhou, Tang, Shen, & Wang, 2018); and extracts contained other compounds such as oligosaccharides, polysaccharides or dietary fiber (Chiu et al., 2017; Gao et al., 2018; Garcia-Mazcorro et al., 2018; Romo-Vaquero et al., 2014). Finally, 24 articles (22 animal studies and 2 clinical trials) fulfilled the eligibility criteria and were included in this systematic review (Fig. 1). The animal studies were organized into two sections, according to polyphenol classes and considering the majority compound of the polyphenols profile available in the studies: flavonoids and other polyphenols (lignans, phenolic acids, stilbenes and vanillin) (Table 2). Data on the polyphenol class and subclass were extracted from the Phenol-Explorer database (www.phenol-explorer.eu).

3.2. Prebiotic effect of dietary polyphenols

3.2.1. Animal studies

3.2.1.1. Flavonoids. The flavonoids were the most investigated phenolic compounds in relation to the effects on the gut microbiota composition and the benefits to the host health. Prebiotic effect of anthocyanins, a subclass of the flavonoids, was evaluated in three animal studies (Anhê et al., 2018; Li, Wu, et al., 2019; Van Hul et al., 2018). In a study with Arctic berry extracts (Anhê et al., 2018), and another with cinnamon bark and grape pomace extracts (Van Hul et al., 2018), the most abundant polyphenols present in the extracts were anthocyanins and proanthocyanidins. However, the authors associated the main outcomes with the presence of the proanthocyanidins. The proanthocyanidins or procyanidins effects on the gut microbiota were investigated in other three studies. A marked increase in the abundance of *Akkermansia* was observed after the consumption of polymeric




Fig. 1. Flowchart of the systematic review.

procyanidins from apple (Masumoto et al., 2016), and cranberry extract rich in proanthocyanidins (Anhê et al., 2015, 2017). Moreover, body weight gain reduction, insulin sensitivity improvement, upregulation of genes involved in lipid catabolism, and downregulation of pro-inflammatory genes in the liver were the health benefits associated with the consumption of proanthocyanidins (Anhê et al., 2015, 2017; Masumoto et al., 2016).

The prebiotic effect of the flavanols (epigallocatechin gallate, epigallocatechin, epicatechin gallate, epicatechin), a flavonoid subclass, was characterized in seven animal studies. The main food sources investigated were green tea, black tea, oolong tea, and Pu-erh tea. The

intervention doses were very different among studies, and intervention periods ranged from 28 to 196 days (Dey et al., 2019; Henning et al., 2018; Liu et al., 2019; Lu et al., 2019; Ma et al., 2019; Wang et al., 2018; Xia et al., 2019). An increase in the abundance of *Akkermansia* and *Bifidobacterium* was observed in mice fed a high-fat diet after the consumption of green tea polyphenols (Dey et al., 2019; Ma et al., 2019; Wang et al., 2018). Supplementation of the oolong tea extract and decaffeinated black tea extract increased the butyrate production (Liu et al., 2019; Henning et al., 2018), while the Pu-erh tea extract consumption stimulated the *Akkermansia* and *Roseburia* growth (Lu et al., 2019; Xia et al., 2019).

Table 2
Characteristics of included studies on prebiotic effect of dietary polyphenols (n = 24).

Compound (food)	Dose	Animal/Human	Study design Period (days)	Diet	Main outcome	Reference
					Gut microbiota	
Animal studies						
Flavonoids						
Anthocyanins (bilberry extract)	10, 20 or 40 mg extract/kg bw/day (IG)	Sprague-Dawley rats (n = 10/group)	70	STD	↑ <i>Lactobacillus</i> ↑ cecal SCFA	Li, Wu, et al. (2019)
Anthocyanins and proanthocyanidins (Arctic berry extracts)	200 mg extract/kg bw/day (water)	Male C57BL/6J mice (n = 12/group)	56	HFHS	↓ <i>Firmicutes/Bacteroidetes</i> ratio ↑ <i>Akkermansia muciniphila</i> (NS) cecal SCFA	Anhê et al. (2018)
Anthocyanins (grape pomace extract) and proanthocyanidin A (cinnamon bark extract)	grape pomace: 2 g extract/kg diet cinnamon bark: 8.2 g extract/ kg diet	Male C57BL/6J mice (n = 14/group)	56	HFD	↑ <i>Roseburia</i> (NS) cecal SCFA	Van Hul et al. (2018)
Catechins (green tea extract)	2% of extract (diet)	Male C57BL/6J mice (n = 10/group)	56	HFD	↑ <i>Bifidobacterium</i> and <i>Akkermansia muciniphila</i> ↓ <i>Firmicutes/Bacteroidetes</i> ratio	Dey et al. (2019)
Catechins (green tea polyphenols)	0.05, 0.2 or 0.8% of tea polyphenols (diet)	HFA C57BL/6J mice (n = 8/group)	56	HFHS	↑ fecal acetate and butyrate ↓ <i>Firmicutes/Bacteroidetes</i> ratio	Wang et al. (2018)
Catechins (green tea polyphenols)	100, 200 or 400 mg of green tea polyphenols/kg bw/day (IG)	SPF C57BL/6 mice (n = 8/group)	84	HFD	↑ <i>Bifidobacterium</i> , <i>Akkermansia</i> , <i>Roseburia</i> (the highest dose)	Ma et al. (2019)
Catechins and caffeine (green tea, black tea and oolong tea water extracts)	1% tea extract (diet)	Male C57BL/6J mice (n = 12/group)	196	HFD	↑ fecal SCFA (oolong tea) ↓ plasma LPS	Liu et al. (2019)
Catechins (decaffeinated green and black tea polyphenol extracts)	0.5 g extract/100 g diet	Male C57BL/6J Mice (n = 12/group)	28	HFHS	↑ fecal propionate and butyrate (black tea) ↓ <i>Firmicutes</i> ↑ <i>Bacteroidetes</i>	Henning et al. (2018)
Catechins (ripened Pu-erh tea extract)	0.1, 0.2, or 0.4% of extract (water)	Male C57BL/6N mice (n = 10/group)	56	HFD	↑ <i>Roseburia</i> and <i>Akkermansia</i> ↓ <i>Firmicutes/Bacteroidetes</i> ratio ↓ serum LPS	Lu et al. (2019)
Catechins and caffeine (aqueous raw and ripe Pu-erh tea extracts)	0.15 or 0.4 mg extract/kg bw/ day (diet)	Male Wistar rats (n = 18/group)	77	HFHS	↑ <i>Akkermansia muciniphila</i> ↑ <i>Firmicutes</i> ↓ <i>Bacteroidetes</i>	Xia et al. (2019)
Genistein	3 mg/kg bw/day (diet)	Male C57BL6 mice (n = 8/group)	180	HFD	↑ <i>Faecalibacterium</i> ↓ serum LPS	López et al. (2018)
Hydroxysafflor yellow A	200 mg/kg bw/day (IG)	SPF C57BL/6J mice (n = 8/group)	84	HFD	↑ <i>Akkermansia</i> ↑ cecal SCFA	Liu et al. (2018)
Isoflavones (soy extract)	150 or 450 mg extract/kg bw/ day (diet)	Male Sprague Dawley rats (n = 16/group)	91	HFD	↓ <i>Firmicutes/Bacteroidetes</i> ratio ↑ <i>Faecalibacterium</i> ↑ fecal butyrate and ↓ serum LPS	Luo et al. (2019)
Naringenin (<i>S. chinensis</i> pollen extract)	7.86 or 15.72 g extract/kg bw/ day (IG)	Male C57BL/6 mice (n = 12/group)	112	HFD	↓ <i>Firmicutes/Bacteroidetes</i> ratio ↑ <i>Lactobacillus</i>	Cheng, Chen, Liu, Zhao, and Cao (2019)
(continued on next page)						

(continued on next page)

Table 2 (continued)

Compound (food)	Dose	Animal/Human	Study design		Main outcome	Reference
			Period (days)	Diet		
Polymeric and oligomeric procyanidin (apple)	0.5% procyanidins (diet)	C57BL/6J male mice (n = 10/group)	140	HFHS	↑ <i>Akkermansia</i> (eight times) ↓ <i>Firmicutes/Bacteroidetes</i> ratio ↓ serum LPS	Masumoto et al. (2016)
Proanthocyanidin (cranberry extract)	200 mg extract/kg bw/day (water)	Male C57BL/6J mice (n = 8–11/group)	147	HFHS	↑ <i>Akkermansia muciniphila</i> ↓ <i>Firmicutes/Bacteroidetes</i> ratio	Anhê et al. (2017)
Proanthocyanidin (cranberry extract)	200 mg extract/kg bw/day (IG)	Male C57BL/6J mice (n = 12/group)	56	HFHS	↑ <i>Akkermansia</i> spp. ↓ plasma LPS	Anhê et al. (2015)
<i>Other polyphenols</i>						
Lignans - syringaresinol	10 or 50 mg/kg bw/day (diet)	Male C57BL/6 mice (n = 12/group)	70	STD	↑ <i>Lactobacillus</i> and <i>Bifidobacterium</i> ↓ serum LPS-binding protein	Cho et al. (2016)
Phenolic acids - dicaffeoylquinic acids (<i>Ilex kudingcha</i>)	3.3 or 10 mg/mouse (IG)	SPF male C57BL/6 mice (n = 8/group)	56	HFD	↑ <i>Bifidobacterium</i> and <i>Akkermansia</i> ↓ serum LPS	Xie et al. (2019)
Stilbenes - pterostilbene	15 mg/kg bw/day (IG)	Zucker (<i>f</i> u/ <i>f</i> a) rats (n = 10/group)	42	STD	↑ <i>Akkermansia muciniphila</i> ↓ <i>Firmicutes</i>	Exeberria et al. (2017)
Stilbenes - resveratrol	200 mg/kg bw/day (diet)	Male Kunming mice (n = 8/group)	84	HFD	↑ <i>Lactobacillus</i> and <i>Bifidobacterium</i> ↓ <i>Firmicutes</i> ↑ <i>Bacteroidetes</i>	Qiao et al. (2014)
Other polyphenols - vanillin	0.1% diet	Male C57BL/6J mice (n = 7–8/group)	98	HFD	↑ <i>Firmicutes</i> ↓ <i>Bacteroidetes</i> ↑ cecal SCFA	Guo, Han, Zhan, You, and Huang (2018)
<i>Clinical trials</i>						
Anthocyanins (wild blueberry drink)	25 g/250 mL água	20 healthy male individuals	crossover, placebo controlled (6 weeks)		↑ <i>Lactobacillus acidophilus</i> ↑ <i>Bifidobacterium</i> spp.	Vendrame et al. (2011)
Punicalagins and ellagic acid (pomegranate extract)	0.45 g or 1.8 g extract (capsules)	49 overweight-obese subjects (mild hyperlipidemia)	double-blind, crossover, placebo controlled (9 weeks)		↑ <i>Faecalibacterium</i> ↓ plasma LBP	González-Sarrías et al. (2018)

bw: body weight, SCFA: short-chain fatty acids, STD: standard, chow and normal diet, HFD: high-fat diet, HFHS: high-fat/high-sucrose diet, IG: intragastric, LPS: lipopolysaccharides, HFA: human flora-associated, SPF: specific-pathogen-free, (NS): not significant or not affected, NR: not reported, LBP: lipopolysaccharide-binding protein, ↑ and ↓: increase and decrease in the specific parameter, respectively.

The intervention with soy isoflavone extract containing 51% daidzin, 30% glycitin and 9% genistein increased the abundance of *Faecalibacterium* and the fecal butyrate content in obese rats, and improved intestinal barrier function through increasing the expressions of zonula occludens 1 (ZO-1), occludin and mucin 2 (Muc-2) genes (Luo et al., 2019). An increase of the *Faecalibacterium* genus was also observed after the consumption of genistein (López et al., 2018).

The effects of other flavonoid subclasses were investigated in two studies, in which the intervention samples were hydroxysafflor yellow A (HSYA) (Liu et al., 2018) and *Schisandra chinensis* pollen extract (Cheng et al., 2019). The HSYA supplementation, a compound with a mono-chalcone glycoside structure, increased the abundance of *Akkermansia*, and the production of acetate, propionate, and butyrate. Moreover, the number of goblet cells and the expression of tight junction proteins ZO-1 were increased after intragastric supplementation with HSYA (Liu et al., 2018). Regarding the *S. chinensis* pollen extract, the most abundant phenolic compound was naringenin (1.89 mg/g), a flavanone. Treatment with pollen extract increased *Lactobacillus* in obese mice with a dose-effect relationship (Cheng, Chen, Liu, et al., 2019).

3.2.1.2. Other polyphenols. Prebiotic effect of other phenolic compounds, like lignans, phenolic acids, stilbenes and vanillin, was evaluated in five animal studies (Cho et al., 2016; Etxeberria et al., 2017; Guo et al., 2018; Qiao et al., 2014; Xie et al., 2019). In summary, there was an increase in the abundance of *Lactobacillus*, *Bifidobacterium* and *Akkermansia*, and a reduction in serum concentrations of pro-inflammatory cytokines and expression of lipid synthesis related genes.

The consumption of syringaresinol, a lignan present in oilseeds, cereal brans and berry seeds, enhanced the population of *Lactobacillus* and *Bifidobacterium*, and reduced serum lipopolysaccharide-binding protein (LBP) concentration (Cho et al., 2016). Supplementation with aqueous kudingcha extract, composed by 3,4-di-O-caffeoylquinic acids (3,4-diCQAs – 26.9%), 3,5-diCQAs (42.3%) and 4,5-diCQAs (30.8%), increased the abundance of *Bifidobacterium* and *Akkermansia*, and decreased the concentrations of serum interleukin 6 (IL-6), tumor necrosis factor- α (TNF- α) and lipopolysaccharides (LPS) and the hepatic expression of lipid synthesis related genes (Xie et al., 2019). The supplementation with resveratrol, the largest representative of the stilbenes class, increased the abundance of *Lactobacillus* and *Bifidobacterium*, and decreased the genes expression related to fatty acids synthesis, adipogenesis and lipogenesis in mice fed a high-fat diet (Qiao et al., 2014). Animals treated with a standard diet supplemented with pterostilbene, a dimethoxy resveratrol derivative, showed an increase in the abundance of *Akkermansia muciniphila*, and an improvement in insulin sensitivity (Etxeberria et al., 2017). An increase in the production of acetate, propionate and butyrate and a decrease in the high concentrations of inflammatory factors (LPS, IL-6, and TNF- α) were observed in mice fed a high-fat diet after the supplementation with vanillin (Guo, Han, et al., 2018).

3.2.2. Clinical trials

The effect of anthocyanins was also evaluated in a crossover clinical trial, in which the consumption of a wild blueberry drink (375 mg of anthocyanins) by healthy adults increased *Bifidobacterium* spp. and *Lactobacillus acidophilus* (Vendrame et al., 2011). In another clinical trial with overweight-obese individuals, the consumption of pomegranate extract (1.8 g/day, 656 mg of phenolics) rich in hydroxybenzoic acids (punicalagins and free ellagic acid) increased *Faecalibacterium* and decreased the concentration of plasma LBP (González-Sarriás et al., 2018).

3.3. Methodological quality assessment of studies

The animal studies were generally classified as low or unclear regarding the risk of bias (Table S1). None of the included studies

satisfied all areas established by the SYRCLE's tool for methodological quality assessment of animal studies. Data on selection bias (baseline characteristics and allocation concealment), performance bias (random housing and blinding), and detection bias (random outcome assessment and blinding) were unavailable in all studies. The clinical trials (González-Sarriás et al., 2018; Vendrame et al., 2011), evaluated by Cochrane Collaboration's tool, were at low-risk for selection bias (random sequence generation), performance bias (blinding of participants and personnel), reporting bias (selective reporting), and other bias. In addition, these studies were unclear for selection bias (allocation concealment) and detection bias (blinding of outcome assessment) and showed high-risk for attrition bias (incomplete outcome data).

4. Discussion

To our knowledge, this is the first systematic review aiming to explore the prebiotic effect of dietary polyphenols investigated in pre-clinical and clinical studies. We found 24 studies of which: 8 showed an increased abundance of *Lactobacillus* spp. and/or *Bifidobacterium* spp. (Cheng, Chen, Liu, et al., 2019; Cho et al., 2016; Dey et al., 2019; Li, Wu, et al., 2019; Ma et al., 2019; Qiao et al., 2014; Vendrame et al., 2011; Xie et al., 2019); 10 showed an increased abundance of *Akkermansia* spp. (Anhê et al., 2015, 2017; Dey et al., 2019; Etxeberria et al., 2017; Liu et al., 2018; Lu et al., 2019; Ma et al., 2019; Masumoto et al., 2016; Xia et al., 2019; Xie et al., 2019); 3 showed an increased abundance of *Faecalibacterium* spp. (González-Sarriás et al., 2018; López et al., 2018; Luo et al., 2019); and 3 showed an increased abundance of *Roseburia* spp. (Lu et al., 2019; Ma et al., 2019; Van Hul et al., 2018). Seven studies reported an increase in the production of SCFA, including butyrate (Guo, Han, et al., 2018; Henning et al., 2018; Li, Wu, et al., 2019; Liu et al., 2018, 2019; Luo et al., 2019; Wang et al., 2018).

The current definition of prebiotic recognizes that prebiotic targets extend beyond the stimulation of *Bifidobacterium* and *Lactobacillus*, and include other microorganisms such as *Roseburia*, *Eubacterium* and *Faecalibacterium* spp., but are not limited to these (Gibson et al., 2017). Thereby, the inclusion criteria applied in this systematic review is in accordance with the current concept of prebiotics. The health benefits associated with prebiotics are immune modulation, increased mineral absorption, improved bowel function, and a positive effect on glucose homeostasis, inflammation, blood lipid profile, satiety and defense against pathogens (Sanders et al., 2019). Although these effects cannot be easily extrapolated to the human gut microbiota, many of them seem to be mediated by SCFA, especially acetate, propionate, and butyrate.

The pathway that explains how polyphenols increase the production of SCFA is not yet fully understood. It is believed that the increase of anaerobic microorganisms, such as *Lactobacillus*, *Lachnospiraceae* and *Ruminococcaceae* can promote the increase of SCFA, especially of butyrate (Li, Wu, et al., 2019; Liu et al., 2018, 2019). Another possible explanation is associated with decaffeinated green tea and black tea polyphenols that have been shown to inhibit α -amylase and α -glucosidase in saliva and small intestine, which may lead to residual carbohydrate in the large intestine providing substrate for the SCFA production (Henning et al., 2018). It is worth mentioning that SCFA are the major mediators among nutrition, gut microbiota, physiology, and pathology (Ríos-Covián et al., 2016). The butyrate has been investigated most extensively, and it is recognized that butyrate-producing bacteria and butyrate per se may be beneficial for human health (Koh, De Vadder, Kovatcheva-Datchary, & Bäckhed, 2016), suggesting it as a biomarker of prebiotic effect.

The studies reviewed here reported that the health benefits associated with the consumption of catechins were reduced concentrations of serum low-density lipoprotein cholesterol (LDL-c) (Ma et al., 2019), glucose and insulin (Wang et al., 2018). Studies also found that catechins could prevent an increase in toll-like receptor 4 (TLR4)/ nuclear factor kappa B (NF κ B)-dependent inflammatory genes (Dey et al., 2019), and increase the hepatic adenosine monophosphate-activated

protein kinase (AMPK) phosphorylation (Henning et al., 2018). Besides, the consumption of green tea extract containing 48% epigallocatechin gallate prevented from a decrease in the expression of intestinal tight junction proteins induced by a high-fat diet (Dey et al., 2019). The treatment with soy isoflavone and HSYA improved intestinal barrier function through increasing the number of goblet cells and the expression of tight junction proteins ZO-1, occludin and Muc-2 (Liu et al., 2018; Luo et al., 2019). Digestive enzymes activities (β -glucosidase, β -galactosidase and β -glucuronidase) in the cecal environment were increased by the supplementation with anthocyanins (Li, Wu, et al., 2019). The health effects of proanthocyanidins consumption were improved glucose tolerance and insulin sensitivity, upregulation of lipid catabolism genes (PPAR α), and downregulation of pro-inflammatory genes (COX2, TNF α) in the liver (Anhê et al., 2015, 2017; Masumoto et al., 2016).

Dietary flavonoids are consumed predominantly as glycosides (conjugated to sugar), which hinders the absorption of these compounds by the small intestine (Kawabata, Yoshioka, & Terao, 2019). Glycosylated flavonoids can serve as the sole source of carbon and energy for some microorganisms in the gut microbiota, which preferentially ferment the sugars linked to flavonoids. Thus, it could explain the prebiotic effect observed for these compounds (Braune & Blaut, 2016). For instance, aqueous extract of jaboticaba, a Brazilian native fruit rich in cyanidin-3-O-glucoside and delphinidin-3-O-glucoside, modulated the abundance of *Lactobacillus* and *Bifidobacterium* in rats with induced colitis (Silva-Maia et al., 2019). Cyanidin-3-O-glucoside ameliorated gut microbial dysbiosis caused by 3-chloro-1,2-propanediol (chemical food contaminant) in rats (Chen et al., 2019). In other study, the Quzhou Fructus Aurantii extract, rich in naringin, a natural flavanone glycoside, increased the genus *Akkermansia* and the expression of tight junction proteins, and reduced metabolic endotoxemia in mice fed high-fat diet (Bai et al., 2019).

Most dietary polyphenols arrive intact in the colon, where they become substrates for the gut microbiota, producing better-absorbing metabolites (Mojzer et al., 2016; Kawabata et al., 2019). The flavonoids, such as epicatechin, catechin, procyanidin and quercetin, when metabolized by gut microbiota, generate hydroxy phenylacetic and hydroxyphenyl propionic acids (Shortt et al., 2018). Equol and O-desmethylanagolensin (ODMA) are active metabolites produced by the action of colonic bacteria on soy isoflavones (Mayo, Vázquez, & Flórez, 2019). The urolithins, in turn, are metabolites produced from ellagitannins and ellagic acid by the human gut microbiota (Tomás-Barberán et al., 2017). These metabolites also have beneficial health effects, such as estrogenic and antioxidant activity (Mayo et al., 2019), anti-inflammatory and antioxidant effects (Lee, Park, Lee, Ahn, & Kim, 2019), and hepatoprotective effect (Zhao et al., 2018). Thus, further studies are warranted to investigate the potential of these metabolites as prebiotic effect markers, such as the SCFA modulation.

Studies with syringaresinol in male C57BL/6 mice (Cho et al., 2016) and pomegranate extract in overweight-obese individuals (González-Sarriás et al., 2018) demonstrated a decrease in serum LBP, indicating that polyphenols could modulate metabolic endotoxemia (Fuks, Nagata, Suganuma, & Ota, 2019). LBP is a glycoprotein mainly synthesized in the hepatocytes with long half-life in the blood and can bind to LPS promoting an LPS-induced immune response via toll-like receptors in macrophages (Jamar, Ribeiro, & Pisani, 2020). LPS, known as endotoxin, is a breakdown product present in the outer membrane of gram-negative bacteria, composed by an O-antigen portion in its outer part and by a lipid-A portion in its inner part. The lipid-A portion exerts most of the immunogenic effects, such as the activation of TLR4, through the formation of the complex containing LBP and the CD14 co-receptor, which signals the NF κ B activation to upregulate pro-inflammatory mediators causing low-grade inflammation. The O-antigen portion activates components of the adaptive immunity, intending to induce the production of antibodies. Therefore, plasma LBP may be an inflammation marker caused by endotoxins (Jamar et al., 2020).

The dysbiosis of the gut microbiota caused by a high-fat diet has been considered as a possible cause for metabolic endotoxemia (Fuks et al., 2019). Thus, it is expected that the consumption of polyphenols may also improve the metabolic endotoxemia associated to dysbiosis, since the consumption of catechins (Liu et al., 2019; Lu et al., 2019), isoflavones (López et al., 2018; Luo et al., 2019), proanthocyanidins (Anhê et al., 2015; Masumoto et al., 2016), and dicaffeoylquinic acids (Xie et al., 2019) decreased plasma or serum LPS in animals fed a high-fat diet. Circulating LPS is released by lysis of a fraction of the bacterial cell wall and flows into the blood by increasing intestinal permeability. The excess intestinal LPS itself, caused by dysbiosis, destroys the narrow junction of intestinal epithelial cells via TLR4 and inhibits mRNA expression of factors related to the restricted junction, such as ZO-1 and occludin, in the intestinal epithelial cell. Therefore, LPS is considered an inflammation marker for dysbiosis (Fuks et al., 2019).

The improvement of the intestinal barrier function through increasing the expressions of ZO-1, occludin and Muc-2 genes, and of the mucin-producing goblet cells number is one of the possible mechanisms associated with the prebiotic effects of catechins (Dey et al., 2019), soy isoflavones (Luo et al., 2019) and HSYA (Liu et al., 2019). Kruppel-like factor 4 (KLF4), a marker of goblet cells, and Muc2 mRNA expression in the proximal colon were also associated with the administration of proanthocyanidins rich-cranberry extract, supporting that these polyphenols could be able to stimulate mucus production (Fig. 2), and therefore create an ecological niche for the *Akkermansia* spp., a mucus-degrading bacterium (Anhê et al., 2015). Xia et al. (2019) reported an increased abundance of *Akkermansia* spp. through an increase in other markers, such as type II and III secretion system proteins, the elongation factor thermo unstable, and a glyceraldehyde-3-phosphate dehydrogenase. This methodology was not observed in other studies. In this review, *Akkermansia* spp. was included as prebiotic target along with the bacteria recognized as probiotics by the Consensus Statement on the Definition and Scope of Prebiotics (Gibson et al., 2017), considering the recent evidences in literature about the relationship among *Akkermansia* spp., gut microbiota and human health (Cani & de Vos, 2017; Jayachandran, Chung, & Xu, 2019; Sanders et al., 2019). *Akkermansia* spp. and *Propionibacterium* spp. are promising candidates among the next generation of microorganisms to be recognized as prebiotic targets (Sanders et al., 2019).

There is also evidence of an additional mechanism by which polyphenols protect the intestinal barrier against oxidative stress. A decrease in the oxidative stress in the liver and ileum was observed in specific pathogen-free mice after a lower-dose consumption of green tea polyphenols (100 mg/kg body weight/day). The improvement of intestinal oxidative stress is supposed to be a potential mechanism for the modulation of tea polyphenols in the gut microbiota (Ma et al., 2019), since excess reactive oxygen species will damage the cell membrane and disrupt the tight junctions leading to an increased intestinal permeability and the development of metabolic disorders (Qiao, Sun, Ding, Le, & Shi, 2013). The tight junctions are multiprotein complexes that maintain barrier function between the enterocytes, creating paracellular barrier properties, which are composed by transmembrane proteins which control the transport across the intercellular space between adjacent cells and cytoplasmic plaque (Costea et al., 2019). The main components of the cytoplasmic plaque are claudin and zonula occludens proteins. In oxidative stress, the interactions of occludin with claudins or proteins of the zonula family are affected directly influencing the formation and function of the tight junctions. Oxidative stress downregulates occludin, reduces its specific membrane localization and regulatory contribution to barrier tightness via multiple signaling pathways (Costea et al., 2019). Nonetheless, most of the evidence is based *in vitro* and animal studies, and then further studies in humans are needed to clarify these mechanisms.

Regarding the methodological quality of the included papers, none of the studies satisfied all criteria established by the SYRCL and Cochrane Collaboration's tool. No preclinical studies provide detailed descriptions of the methods used for allocation concealment, random




Fig. 2. Mechanism of mucin synthesis modulation by proanthocyanidins in the hindgut. KLF4: kruppel-like factor 4, Muc-2: mucin 2, ZO-1: zonula occludens 1. The consumption of polyphenols, represented in the figure by proanthocyanidins, may increase the number of goblet cells, according to the Kruppel-like factor 4 (KLF4) marker, and consequently, the mucin production; so the enhanced mucin provides a favorable environment for the proliferation of *Akkermansia* spp., a mucus-degrading bacterium (Anhê et al., 2015; Xia et al., 2019). In addition, increased expression of ZO-1 and occludin can be observed after the consumption of polyphenols, decreasing the intestinal permeability through the junction of epithelial cells (tight junctions) (Dey et al., 2019; Luo et al., 2019; Liu et al., 2019).

housing and random outcome assessment, or blinding researchers and outcome assessors. The clinical trials did not provide details on allocation concealment and blinding of outcome assessment. The absence of descriptions regarding sample losses in the preclinical and clinical studies also was noted. Therefore, we suggest the use of these tools in experimental planning, as a reference for well-designed studies aiming to reduce the risk of bias and allow more consistent conclusions.

It is noteworthy that the included studies had quite heterogeneous designs, including differences in the intervention period, the methodology for gut microbiome analysis, and the supplementation with polyphenol (food sample and dosage), as well as the polyphenols profile of the samples. Additional studies with isolated polyphenols are needed to eliminate the effects of other compounds in the extract, especially in clinical trials. Moreover, we suggest that further studies aiming to evaluate the prebiotic effect of dietary polyphenols should be carried out according to the internationally accepted prebiotics definition, with complete assessment of the gut microbiota, including the microorganism's genera. The low number of studies on humans is also a limitation in this systematic review.

5. Conclusion

The prebiotic effect of dietary polyphenols, especially catechins, anthocyanins, and proanthocyanidins, has strong evidence based on preclinical studies. Despite the limitations of these studies, it is evident that polyphenols can stimulate the growth of microorganisms recognized as prebiotic targets (*Lactobacillus* spp., *Bifidobacterium* spp., *Akkermansia* spp., *Roseburia* spp., and *Faecalibacterium* spp.), and increase the production of SCFA, including butyrate. Nevertheless, well-designed clinical trials are warranted to prove the prebiotic effect of

polyphenols on humans.

Ethics statement

This research did not include any human subjects and animal experiments.

CRediT authorship contribution statement

Aline Medeiros Alves-Santos: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Writing - original draft, Writing - review & editing. **Clara Sandra Araújo Sugizaki:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Writing - original draft, Writing - review & editing. **Glaucia Carielo Lima:** Conceptualization, Data curation, Supervision, Visualization, Writing - review & editing. **Maria Margareth Veloso Naves:** Conceptualization, Data curation, Supervision, Visualization, Writing - review & editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors thank the CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil) for a research scholarship.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jff.2020.104169>.

References

- Anhê, F. F., Nachbar, R. T., Varin, T. V., Vilela, V., Dudonné, S., Pilon, G., ... Marette, A. (2017). A polyphenol-rich cranberry extract reverses insulin resistance and hepatic steatosis independently of body weight loss. *Molecular Metabolism*, 6(12), 1563–1573. <https://doi.org/10.1016/j.molmet.2017.10.003>.
- Anhê, F. F., Roy, D., Pilon, G., Dudonné, S., Matamoros, S., Varin, T. V., ... Marette, A. (2015). A polyphenol-rich cranberry extract protects from diet-induced obesity, insulin resistance and intestinal inflammation in association with increased *Akkermansia* spp. population in the gut microbiota of mice. *Gut*, 64(6), 872–883. <https://doi.org/10.1136/gutjnl-2014-307142>.
- Anhê, F. F., Varin, T. V., Le Barz, M., Pilon, G., Dudonné, S., Trottier, J., ... Marette, A. (2018). Arctic berry extracts target the gut–liver axis to alleviate metabolic endotoxaemia, insulin resistance and hepatic steatosis in diet-induced obese mice. *Diabetologia*, 61(4), 919–931. <https://doi.org/10.1007/s00125-017-4520-z>.
- Bai, Y., Wang, S., Wang, X., Weng, Y., Fan, X., Sheng, H., ... Zhang, F. (2019). The flavonoid-rich Quzhou Fructus Aurantii extract modulates gut microbiota and prevents obesity in high-fat diet-fed mice. *Nutrition and Diabetes*, 9(1), 1–11. <https://doi.org/10.1038/s41387-019-0097-6>.
- Bian, Y., Wei, J., Zhao, C., & Li, G. (2020). Natural polyphenols targeting senescence: A novel prevention and therapy strategy for cancer. *International Journal of Molecular Sciences*, 21(2), 1–17. <https://doi.org/10.3390/ijms21020684>.
- Braune, A., & Blaut, M. (2016). Bacterial species involved in the conversion of dietary flavonoids in the human gut. *Gut Microbes*, 7(3), 216–234. <https://doi.org/10.1080/19490976.2016.1158395>.
- Cani, P. D., & de Vos, W. M. (2017). Next-generation beneficial microbes: The case of *Akkermansia muciniphila*. *Frontiers in Microbiology*, 8, 1–8. <https://doi.org/10.3389/fmicb.2017.01765>.
- Casanova-Martí, A., Serrano, J., Portune, K. J., Sanz, Y., Blay, M. T., Terra, X., ... Pinent, M. (2018). Grape seed proanthocyanidins influence gut microbiota and enteroendocrine secretions in female rats. *Food and Function*, 9(3), 1672–1682. <https://doi.org/10.1039/c7fo02028g>.
- Chai, S. C., Davis, K., Zhang, Z., Zha, L., & Kirschner, K. F. (2019). Effects of tart cherry juice on biomarkers of inflammation and oxidative stress in older adults. *Nutrients*, 11(2), 1–10. <https://doi.org/10.3390/nu11020228>.
- Chen, G., Wang, G., Zhu, C., Jiang, X., Sun, J., Tian, L., & Bai, W. (2019). Effects of cyanidin-3-O-glucoside on 3-chloro-1,2-propanediol induced intestinal microbiota dysbiosis in rats. *Food and Chemical Toxicology*, 133, 110767. <https://doi.org/10.1016/j.fct.2019.110767>.
- Cheng, L., Chen, Y., Zhang, X., Zheng, X., Cao, J., Wu, Z., ... Cheng, K. (2019). A metagenomic analysis of the modulatory effect of *Cyclocarya paliurus* flavonoids on the intestinal microbiome in a high-fat diet-induced obesity mouse model. *Journal of the Science of Food and Agriculture*, 99(8), 3967–3975. <https://doi.org/10.1002/jsfa.9622>.
- Cheng, N., Chen, S., Liu, X., Zhao, H., & Cao, W. (2019). Impact of Schisandra chinensis bee pollen on nonalcoholic fatty liver disease and gut microbiota in highfat diet induced obese mice. *Nutrients*, 11(2), 1–16. <https://doi.org/10.3390/nu11020346>.
- Chiu, H. F., Chen, Y. J., Lu, Y. Y., Han, Y. C., Shen, Y. C., Venkatakrishnan, K., & Wang, C. K. (2017). Regulatory efficacy of fermented plant extract on the intestinal microflora and lipid profile in mildly hypercholesterolemic individuals. *Journal of Food and Drug Analysis*, 25(4), 819–827. <https://doi.org/10.1016/j.jfda.2016.10.008>.
- Cho, S. Y., Kim, J., Lee, J. H., Sim, J. H., Cho, D. H., Bae, I. H., ... Kim, H. R. (2016). Modulation of gut microbiota and delayed immunosenescence as a result of syringaresinol consumption in middle-aged mice. *Scientific Reports*, 6, 1–14. <https://doi.org/10.1038/srep39026>.
- Cires, M. J., Navarrete, P., Pastene, E., Carrasco-Pozo, C., Valenzuela, R., Medina, D. A., ... Gotteland, M. (2019). Effect of a proanthocyanidin-rich polyphenol extract from avocado on the production of amino acid-derived bacterial metabolites and the microbiota composition in rats fed a high-protein diet. *Food and Function*, 10(7), 4022–4035. <https://doi.org/10.1039/c9fo00700h>.
- Collins, B., Hoffman, J., Martinez, K., Grace, M., Lila, M. A., Cockrell, C., ... McIntosh, M. (2016). A polyphenol-rich fraction obtained from table grapes decreases adiposity, insulin resistance and markers of inflammation and impacts gut microbiota in high-fat-fed mice. *Journal of Nutritional Biochemistry*, 31, 150–165. <https://doi.org/10.1016/j.jnutbio.2015.12.021>.
- Costea, L., Mészáros, A., Bauer, H., Bauer, H. C., Traweger, A., Wilhelm, I., ... Krizbai, I. A. (2019). The blood–brain barrier and its intercellular junctions in age-related brain disorders. *International Journal of Molecular Sciences*, 20(21), 1–28. <https://doi.org/10.3390/ijms20215472>.
- Cowan, T. E., Palmnäs, M. S. A., Yang, J., Bomhof, M. R., Ardell, K. L., Reimer, R. A., ... Shearer, J. (2014). Chronic coffee consumption in the diet-induced obese rat: Impact on gut microbiota and serum metabolomics. *Journal of Nutritional Biochemistry*, 25(4), 489–495. <https://doi.org/10.1016/j.jnutbio.2013.12.009>.
- Cueva, C., Silva, M., Pinillos, I., Bartolomé, B., & Moreno-Arribas, M. V. (2020). Interplay between dietary polyphenols and oral and gut microbiota in the development of colorectal cancer. *Nutrients*, 12(3), 1–19. <https://doi.org/10.3390/nu12030625>.
- Dey, P., Sasaki, G. Y., Wei, P., Li, J., Wang, L., Zhu, J., ... Bruno, R. S. (2019). Green tea extract prevents obesity in male mice by alleviating gut dysbiosis in association with improved intestinal barrier function that limits endotoxin translocation and adipose inflammation. *Journal of Nutritional Biochemistry*, 67, 78–89. <https://doi.org/10.1016/j.jnutbio.2019.01.017>.
- Etxeberria, U., Hijona, E., Aguirre, L., Milagro, F. I., Bujanda, L., Rimando, A. M., ... Portillo, M. P. (2017). Pterostilbene-induced changes in gut microbiota composition in relation to obesity. *Molecular Nutrition and Food Research*, 61(1), 1–12. <https://doi.org/10.1002/mnfr.201500906>.
- Fang, C., Kim, H., Yanagisawa, L., Bennett, W., Sirven, M. A., Alaniz, R. C., ... Mertens-Talcott, S. U. (2019). Gallotannins and *Lactobacillus plantarum* WCFS1 mitigate high-fat diet-induced inflammation and induce biomarkers for thermogenesis in adipose tissue in gnotobiotic mice. *Molecular Nutrition and Food Research*, 63(9), 1–12. <https://doi.org/10.1002/mnfr.201800937>.
- Fuke, N., Nagata, N., Suganuma, H., & Ota, T. (2019). Regulation of gut microbiota and metabolic endotoxemia with dietary factors. *Nutrients*, 11(10), 1–38. <https://doi.org/10.3390/nu11102277>.
- Gao, X., Xie, Q., Kong, P., Liu, L., Sun, S., Xiong, B., ... Xiang, H. (2018). Polyphenol- and caffeine-rich postfermented Pu-erh tea improves diet-induced metabolic syndrome by remodeling intestinal homeostasis in mice. *Infection and Immunity*, 86(1), 1–16. <https://doi.org/10.1128/IAI.00601-17>.
- García-Mazcorro, J. F., Pedreschi, R., Chew, B., Dowd, S. E., Kawan, J. R., & Noratto, G. (2018). Dietary supplementation with raspberry extracts modifies the fecal microbiota in obese diabetic db/db mice. *Journal of Microbiology and Biotechnology*, 28(8), 1247–1259. <https://doi.org/10.4014/jmb.1803.03020>.
- Gibson, G. R., Hutkins, R., Sanders, M. E., Prescott, S. L., Reimer, R. A., Salminen, S. J., ... Reid, G. (2017). Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nature Reviews Gastroenterology and Hepatology*, 14(8), 491–502. <https://doi.org/10.1038/nrgastro.2017.75>.
- Ginés, I., Gil-Cardoso, K., Terra, X., Blay, M., Pérez-Vendrell, A. M., Pinent, M., & Ardévol, A. (2019). Grape seed proanthocyanidins target the enteroendocrine system in cafeteria-diet-fed rats. *Molecular Nutrition and Food Research*, 63(11), 1–7. <https://doi.org/10.1002/mnfr.201800912>.
- González-Sarriás, A., García-Villalba, R., Romo-Vaquero, M., Alasalvar, C., Örem, A., Zafrilla, P., ... Espín, J. C. (2017). Clustering according to urolithin metabolotype explains the interindividual variability in the improvement of cardiovascular risk biomarkers in overweight-obese individuals consuming pomegranate: A randomized clinical trial. *Molecular Nutrition and Food Research*, 61(5), 1–14. <https://doi.org/10.1002/mnfr.201600830>.
- González-Sarriás, A., Romo-Vaquero, M., García-Villalba, R., Cortés-Martín, A., Selma, M. V., & Espín, J. C. (2018). The endotoxemia marker lipopolysaccharide-binding protein is reduced in overweight-obese subjects consuming pomegranate extract by modulating the gut microbiota: A randomized clinical trial. *Molecular Nutrition and Food Research*, 62(11), 1–10. <https://doi.org/10.1002/mnfr.201800160>.
- Griffin, L. E., Witrick, K. A., Klotz, C., Dorenkott, M. R., Goodrich, K. M., Fundaro, G., ... Neilson, A. P. (2017). Alterations to metabolically active bacteria in the mucosa of the small intestine predict anti-obesity and anti-diabetic activities of grape seed extract in mice. *Food and Function*, 8(10), 3510–3522. <https://doi.org/10.1039/c7fo01236e>.
- Grzelak-Błaszczyk, K., Milala, J., Kosmala, M., Kołodziejczyk, K., Sójka, M., Czarnecki, A., ... Jurgoński, A. (2018). Onion quercetin monoglycosides alter microbial activity and increase antioxidant capacity. *Journal of Nutritional Biochemistry*, 56, 81–88. <https://doi.org/10.1016/j.jnutbio.2018.02.002>.
- Guo, J., Han, X., Zhan, J., You, Y., & Huang, W. (2018). Vanillin alleviates high fat diet-induced obesity and improves the gut microbiota composition. *Frontiers in Microbiology*, 9, 1–12. <https://doi.org/10.3389/fmicb.2018.02733>.
- Guo, X., Tang, R., Yang, S., Lu, Y., Luo, J., & Liu, Z. (2018). Rutin and its combination with inulin attenuate gut dysbiosis, the inflammatory status and endoplasmic reticulum stress in Paneth cells of obese mice induced by high-fat diet. *Frontiers in Microbiology*, 9, 1–11. <https://doi.org/10.3389/fmicb.2018.02651>.
- Henning, S. M., Yang, J., Hsu, M., Lee, R. P., Grojean, E. M., Ly, A., ... Li, Z. (2018). Decaffeinated green and black tea polyphenols decrease weight gain and alter microbiome populations and function in diet-induced obese mice. *European Journal of Nutrition*, 57(8), 2759–2769. <https://doi.org/10.1007/s00394-017-1542-8>.
- Higgins, J. P. T., Altman, D. G., Gotzsche, P. C., Jüni, P., Moher, D., Oxman, A. D., ... Sterne, J. A. C. (2011). The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*, 343(7829), 1–9. <https://doi.org/10.1136/bmj.d5928>.
- Hooijmans, C. R., Rovers, M. M., De Vries, R. B. M., Leenaars, M., Ritskes-Hoitinga, M., & Langendam, M. W. (2014). SYRCL's risk of bias tool for animal studies. *BMC Medical Research Methodology*, 14(1), 1–9. <https://doi.org/10.1186/1471-2288-14-43>.
- Jamar, G., Ribeiro, D. A., & Pisani, L. P. (2020). High-fat or high-sugar diets as trigger inflammation in the microbiota-gut-brain axis. *Critical Reviews in Food Science and Nutrition*, 1–19. <https://doi.org/10.1080/10408398.2020.1747046>.
- Jayachandran, M., Chung, S. S. M., & Xu, B. (2019). A critical review of the relationship between dietary components, the gut microbe *Akkermansia muciniphila*, and human health. *Critical Reviews in Food Science and Nutrition*, 1–12. <https://doi.org/10.1080/10408398.2019.1632789>.
- Kawabata, K., Yoshioka, Y., & Terao, J. (2019). Role of intestinal microbiota in the bioavailability and physiological functions of dietary polyphenols. *Molecules*, 24(2), 1–25. <https://doi.org/10.3390/molecules24020370>.
- Koh, A., De Vadder, F., Kovatcheva-Datchary, P., & Bäckhed, F. (2016). From dietary fiber to host physiology: Short-chain fatty acids as key bacterial metabolites. *Cell*, 165(6),

- 1332–1345. <https://doi.org/10.1016/j.cell.2016.05.041>.
- Lee, G., Park, J. S., Lee, E. J., Ahn, J. H., & Kim, H. S. (2019). Anti-inflammatory and antioxidant mechanisms of urolithin B in activated microglia. *Phytomedicine*, 55, 50–57. <https://doi.org/10.1016/j.phymed.2018.06.032>.
- Li, J., Wu, T., Li, N., Wang, X., Chen, G., & Lyu, X. (2019). Bilberry anthocyanin extract promotes intestinal barrier function and inhibits digestive enzyme activity by regulating the gut microbiota in aging rats. *Food and Function*, 10(1), 333–343. <https://doi.org/10.1039/c8fo01962b>.
- Li, Q., Liu, F., Liu, J., Liao, S., & Zou, Y. (2019). Mulberry leaf polyphenols and fiber induce synergistic antiobesity and display a modulation effect on gut microbiota and metabolites. *Nutrients*, 11(5), 1–19. <https://doi.org/10.3390/nu11051017>.
- Liu, J., Hao, W., He, Z., Kwek, E., Zhao, Y., Zhu, H., ... Chen, Z. Y. (2019). Beneficial effects of tea water extracts on the body weight and gut microbiota in C57BL/6J mice fed with a high-fat diet. *Food and Function*, 10(5), 2847–2860. <https://doi.org/10.1039/c8fo02051e>.
- Liu, J., Yue, S., Yang, Z., Feng, W., Meng, X., Wang, A., ... Yan, D. (2018). Oral hydroxysafflor yellow A reduces obesity in mice by modulating the gut microbiota and serum metabolism. *Pharmacological Research*, 134, 40–50. <https://doi.org/10.1016/j.phrs.2018.05.012>.
- López, P., Sánchez, M., Perez-Cruz, C., Velázquez-Villegas, L. A., Syeda, T., Aguilar-López, M., ... Tovar, A. R. (2018). Long-term genistein consumption modifies gut microbiota, improving glucose metabolism, metabolic endotoxemia, and cognitive function in mice fed a high-fat diet. *Molecular Nutrition and Food Research*, 62(16), 1–12. <https://doi.org/10.1002/mnfr.201800313>.
- Lu, X., Liu, J., Zhang, N., Fu, Y., Zhang, Z., Li, Y., ... Cao, Y. (2019). Ripened Pu-erh tea extract protects mice from obesity by modulating gut microbiota composition. *Journal of Agricultural and Food Chemistry*, 67(25), 6978–6994. <https://doi.org/10.1021/acs.jafc.8b04909>.
- Luo, Q., Cheng, D., Huang, C., Li, Y., Lao, C., Xia, Y., ... Chen, Z. (2019). Improvement of colonic immune function with soy isoflavones in high-fat diet-induced obese rats. *Molecules*, 24(6), 1–17. <https://doi.org/10.3390/molecules24061139>.
- Ma, H., Zhang, B., Hu, Y., Wang, J., Liu, J., Qin, R., ... Wang, S. (2019). Correlation analysis of intestinal redox state with the gut microbiota reveals the positive intervention of tea polyphenols on hyperlipidemia in high fat diet fed mice. *Journal of Agricultural and Food Chemistry*, 67(26), 7325–7335. <https://doi.org/10.1021/acs.jafc.9b02211>.
- Masumoto, S., Terao, A., Yamamoto, Y., Mukai, T., Miura, T., & Shoji, T. (2016). Non-absorbable apple procyanidins prevent obesity associated with gut microbial and metabolomic changes. *Scientific Reports*, 6, 1–10. <https://doi.org/10.1038/srep31208>.
- Mayo, B., Vázquez, L., & Flórez, A. B. (2019). Equol: A bacterial metabolite from the daidzein isoflavone and its presumed beneficial health effects. *Nutrients*, 11(9), 1–20. <https://doi.org/10.3390/nu11092231>.
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., ... PRISMA-P Group (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews*, 4(1), 1–9. <https://doi.org/10.1186/2046-4053-4-1>.
- Mojzer, E. B., Knez Hrncić, M., Škerget, M., Knez, Ž., & Bren, U. (2016). Polyphenols: Extraction methods, antioxidative action, bioavailability and anticarcinogenic effects. *Molecules*, 21(7), 1–38. <https://doi.org/10.3390/molecules21070901>.
- Most, J., Penders, J., Lucchesi, M., Goossens, G. H., & Blaak, E. E. (2017). Gut microbiota composition in relation to the metabolic response to 12-week combined polyphenol supplementation in overweight men and women. *European Journal of Clinical Nutrition*, 71(9), 1040–1045. <https://doi.org/10.1038/ejcn.2017.89>.
- Noad, R. L., Rooney, C., McCall, D., Young, I. S., McCance, D., McKinley, M. C., ... McKeown, P. P. (2016). Beneficial effect of a polyphenol-rich diet on cardiovascular risk: A randomised control trial. *Heart*, 102(17), 1371–1379. <https://doi.org/10.1136/heartjnl-2015-309218>.
- Paquette, M., Larqué, A. S. M., Weisnagel, S. J., Desjardins, Y., Marois, J., Pilon, G., ... Jacques, H. (2017). Strawberry and cranberry polyphenols improve insulin sensitivity in insulin-resistant, non-diabetic adults: A parallel, double-blind, controlled and randomised clinical trial. *British Journal of Nutrition*, 117(4), 519–531. <https://doi.org/10.1017/S0007114517000393>.
- Porrás, D., Nistal, E., Martínez-Flórez, S., Pisonero-Vaquero, S., Olcoz, J. L., Jover, R., ... Sánchez-Campos, S. (2017). Protective effect of quercetin on high-fat diet-induced non-alcoholic fatty liver disease in mice is mediated by modulating intestinal microbiota imbalance and related gut-liver axis activation. *Free Radical Biology and Medicine*, 102, 188–202. <https://doi.org/10.1016/j.freeradbiomed.2016.11.037>.
- Qiao, Y., Sun, J., Ding, Y., Le, G., & Shi, Y. (2013). Alterations of the gut microbiota in high-fat diet mice is strongly linked to oxidative stress. *Applied Microbiology and Biotechnology*, 97(4), 1689–1697. <https://doi.org/10.1007/s00253-012-4323-6>.
- Qiao, Y., Sun, J., Xia, S., Tang, X., Shi, Y., & Le, G. (2014). Effects of resveratrol on gut microbiota and fat storage in a mouse model with high-fat-induced obesity. *Food and Function*, 5(6), 1241–1249. <https://doi.org/10.1039/c3fo06030a>.
- Remely, M., Ferk, F., Sterneder, S., Setayesh, T., Roth, S., Kepcija, T., ... Haslberger, A. G. (2017). EGCG prevents high fat diet-induced changes in gut microbiota, decreases of DNA strand breaks, and changes in expression and DNA methylation of Dnmt1 and MLH1 in C57BL/6J male mice. *Oxidative Medicine and Cellular Longevity*, 2017, 1–17. <https://doi.org/10.1155/2017/3079148>.
- Ríos-Covián, D., Ruas-Madiedo, P., Margolles, A., Gueimonde, M., de los Reyes-Gavilán, C. G., & Salazar, N. (2016). Intestinal short chain fatty acids and their link with diet and human health. *Frontiers in Microbiology*, 7, 1–9. 10.3389/fmicb.2016.00185.
- Romo-Vaquero, M., Selma, M. V., Larrosa, M., Obiol, M., García-Villalba, R., González-Barrio, R., ... García-Conesa, M. T. (2014). A rosemary extract rich in carnosic acid selectively modulates caecum microbiota and inhibits β -glucosidase activity, altering fiber and short chain fatty acids fecal excretion in lean and obese female rats. *PLoS ONE*, 9(4), e94687. <https://doi.org/10.1371/journal.pone.0094687>.
- Sanders, M. E., Merenstein, D. J., Reid, G., Gibson, G. R., & Rastall, R. A. (2019). Probiotics and prebiotics in intestinal health and disease: From biology to the clinic. *Nature Reviews Gastroenterology and Hepatology*, 16(10), 605–616. <https://doi.org/10.1038/s41575-019-0173-3>.
- Serrelli, G., & Deiana, M. (2019). In vivo formed metabolites of polyphenols and their biological efficacy. *Food and Function*, 10(11), 6999–7021. <https://doi.org/10.1039/c9fo01733j>.
- Shen, C. Y., Wan, L., Wang, T. X., & Jiang, J. G. (2019). *Citrus aurantium* L. var. *amara* Engl. inhibited lipid accumulation in 3T3-L1 cells and *Caenorhabditis elegans* and prevented obesity in high-fat diet-fed mice. *Pharmacological Research*, 147, 104347. <https://doi.org/10.1016/j.phrs.2019.104347>.
- Shortt, C., Hasselwander, O., Meynier, A., Nauta, A., Fernández, E. N., Putz, P., ... Antoine, J. M. (2018). Systematic review of the effects of the intestinal microbiota on selected nutrients and non-nutrients. *European Journal of Nutrition*, 57(1), 25–49. <https://doi.org/10.1007/s00394-017-1546-4>.
- Silva-Maia, J. K., Batista, A. G., Cazarin, C. B. B., Soares, E. S., Bogusz Junior, S., Leal, R. F., ... Maróstica Junior, M. R. (2019). Aqueous extract of Brazilian berry (*Myrciaria jaboticaba*) peel improves inflammatory parameters and modulates *Lactobacillus* and *Bifidobacterium* in rats with induced colitis. *Nutrients*, 11, 1–13. <https://doi.org/10.3390/nu11112776>.
- Singh, A. K., Cabral, C., Kumar, R., Ganguly, R., & Pandey, A. K. (2019). Beneficial effects of dietary polyphenols on gut microbiota and strategies to improve delivery efficiency. *Nutrients*, 11, 1–21. <https://doi.org/10.3390/nu11092216>.
- Sung, M. M., Kim, T. T., Denou, E., Soltys, C. L. M., Hamza, S. M., Byrne, N. J., ... Dyck, J. R. B. (2017). Improved glucose homeostasis in obese mice treated with resveratrol is associated with alterations in the gut microbiome. *Diabetes*, 66(2), 418–425. <https://doi.org/10.2337/db16-0680>.
- Tan, S., Caparros-Martin, J. A., Matthews, V. B., Koch, H., O'Gara, F., Croft, K. D., & Ward, N. C. (2018). Isoquercetin and inulin synergistically modulate the gut microbiome to prevent development of the metabolic syndrome in mice fed a high fat diet. *Scientific Reports*, 8(1), 1–13. <https://doi.org/10.1038/s41598-018-28521-8>.
- Tomás-Barberán, F. A., González-Sarrías, A., García-Villalba, R., Núñez-Sánchez, M. A., Selma, M. V., García-Conesa, M. T., & Espín, J. C. (2017). Urolithins, the rescue of “old” metabolites to understand a “new” concept: Metabotypes as a nexus among phenolic metabolism, microbiota dysbiosis, and host health status. *Molecular Nutrition and Food Research*, 61(1), 1–35. <https://doi.org/10.1002/mnfr.201500901>.
- Unno, T., & Osakabe, N. (2018). Green tea extract and black tea extract differentially influence cecal levels of short-chain fatty acids in rats. *Food Science and Nutrition*, 6(4), 728–735. <https://doi.org/10.1002/fsn3.607>.
- Van Hul, M., Geurts, L., Plovier, H., Druart, C., Everard, A., Ståhlman, M., ... Cani, P. D. (2018). Reduced obesity, diabetes, and steatosis upon cinnamon and grape pomace are associated with changes in gut microbiota and markers of gut barrier. *American Journal of Physiology - Endocrinology and Metabolism*, 314(4), E334–E352. <https://doi.org/10.1152/ajpendo.00107.2017>.
- Vendrame, S., Guglielmetti, S., Riso, P., Arioli, S., Klimis-Zacas, D., & Porrini, M. (2011). Six-week consumption of a wild blueberry powder drink increases Bifidobacteria in the human gut. *Journal of Agricultural and Food Chemistry*, 59(24), 12815–12820. <https://doi.org/10.1021/jf2028686>.
- Wang, L., Zeng, B., Liu, Z., Liao, Z., Zhong, Q., Gu, L., ... Fang, X. (2018). Green tea polyphenols modulate colonic microbiota diversity and lipid metabolism in high-fat diet treated HFA mice. *Journal of Food Science*, 83(3), 864–873. <https://doi.org/10.1111/1750-3841.14058>.
- Wang, W., Pan, Y., Wang, L., Zhou, H., Song, G., Wang, Y., ... Li, A. (2019). Optimal dietary ferulic acid for suppressing the obesity-related disorders in leptin-deficient obese C57BL/6J-ob/ob mice. *Journal of Agricultural and Food Chemistry*, 67(15), 4250–4258. <https://doi.org/10.1021/acs.jafc.8b06760>.
- Xia, Y., Tan, D., Akbar, R., Kong, J., Seviour, R., & Kong, Y. (2019). Aqueous raw and ripe Pu-erh tea extracts alleviate obesity and alter cecal microbiota composition and function in diet-induced obese rats. *Applied Microbiology and Biotechnology*, 103(4), 1823–1835. <https://doi.org/10.1007/s00253-018-09581-2>.
- Xie, M., Chen, G., Wan, P., Dai, Z., Zeng, X., & Sun, Y. (2019). Effects of dicaffeoylquinic acids from *Ilex kudingcha* on lipid metabolism and intestinal microbiota in high-fat-diet-fed mice. *Journal of Agricultural and Food Chemistry*, 67(1), 171–183. <https://doi.org/10.1021/acs.jafc.8b05444>.
- Yu, J., Guo, H., Xie, J., Luo, J., Li, Y., Liu, L., ... Peng, X. (2019). The alternate consumption of quercetin and alliin in the traditional Asian diet reshaped microbiota and altered gene expression of colonic epithelial cells in rats. *Journal of Food Science*, 84(3), 678–686. <https://doi.org/10.1111/1750-3841.14473>.
- Yuan, L., Li, X., He, S., Gao, C., Wang, C., & Shao, Y. (2018). Effects of natural flavonoid isoorientin on growth performance and gut microbiota of mice. *Journal of Agricultural and Food Chemistry*, 66(37), 9777–9784. <https://doi.org/10.1021/acs.jafc.8b03568>.
- Zhang, C., Wu, W., Li, X., Xin, X., & Liu, D. (2019). Daily supplementation with fresh *Angelica keiskei* Juice alleviates high-fat diet-induced obesity in mice by modulating gut microbiota composition. *Molecular Nutrition and Food Research*, 63(14), 1–9. <https://doi.org/10.1002/mnfr.201900248>.
- Zhang, L., Dong, M., Xu, G., Tian, Y., Tang, H., & Wang, Y. (2018). Metabolomics reveals that dietary ferulic acid and quercetin modulate metabolic homeostasis in rats.

- Journal of Agricultural and Food Chemistry*, 66(7), 1723–1731. <https://doi.org/10.1021/acs.jafc.8b00054>.
- Zhang, X., Zhang, M., Ho, C. T., Guo, X., Wu, Z., Weng, P., ... Cao, J. (2018). Metagenomics analysis of gut microbiota modulatory effect of green tea polyphenols by high fat diet-induced obesity mice model. *Journal of Functional Foods*, 46, 268–277. <https://doi.org/10.1016/j.jff.2018.05.003>.
- Zhao, H., Jiang, Z., Chang, X., Xue, H., Yahefu, W., & Zhang, X. (2018). 4-hydroxyphenylacetic acid prevents acute APAP-induced liver injury by increasing phase II and antioxidant enzymes in mice. *Frontiers in Pharmacology*, 9, 1–10. <https://doi.org/10.3389/fphar.2018.00653>.
- Zhou, J., Tang, L., Shen, C. L., & Wang, J. S. (2018). Green tea polyphenols modify gut-microbiota dependent metabolisms of energy, bile constituents and micronutrients in female Sprague-Dawley rats. *Journal of Nutritional Biochemistry*, 61, 68–81. <https://doi.org/10.1016/j.jnutbio.2018.07.018>.
- Zhou, Y., Zhang, N., Arikawa, A. Y., & Chen, C. (2019). Inhibitory effects of green tea polyphenols on microbial metabolism of aromatic amino acids in humans revealed by metabolomic analysis. *Metabolites*, 9(5), 1–17. <https://doi.org/10.3390/metabo9050096>.