



DRHMC : Delayed Rejection Hamiltonian Monte-Carlo

Chirag Modi

Center for Computational Astrophysics (CCA)
Center for Computational Mathematics (CCM)
Flatiron Institute

StanCon 2023

w/ Alex Barnett, Bob Carpenter
arXiv:2110.00610



Hierarchical models

Q : Researcher is interested in learning the mean SAT score (μ).

Data : Collect students' scores for SAT exams from 5 schools.

Hierarchical models


Q : Researcher is interested in learning the mean SAT score (μ).

Data : Collect students' scores for SAT exams from 5 schools.

Hierarchical model with partial pooling

$$\mu, \sigma \sim \pi(\mu, \sigma); \quad \mu_j \sim \mathbb{N}(\mu, \sigma);$$

$$Y_{ij} \sim \mathbb{N}(\mu_j, \sigma_j)$$



Hierarchical models

Q : Researcher is interested in learning the mean SAT score (μ).

Data : Collect students' scores for SAT exams from 5 schools.



Hierarchical model with partial pooling

$$\mu, \sigma \sim \pi(\mu, \sigma); \quad \mu_j \sim \mathbb{N}(\mu, \sigma);$$

$$Y_{ij} \sim \mathbb{N}(\mu_j, \sigma_j)$$

Degeneracies of hierarchical model

- small σ results in μ_j concentrating around μ
- large σ results in μ_j varying wider range of values



Hierarchical models

Q : Researcher is interested in learning the mean SAT score (μ).

Data : Collect students' scores for SAT exams from 5 schools.

Hierarchical model with partial pooling

$$\mu, \sigma \sim \pi(\mu, \sigma); \quad \mu_j \sim \mathbb{N}(\mu, \sigma);$$



$$Y_{ij} \sim \mathbb{N}(\mu_j, \sigma_j)$$

Degeneracies of hierarchical model

- small σ results in μ_j concentrating around μ
- large σ results in μ_j varying wider range of values


Strong coupling of the individual parameters (μ_j) to the population parameters (μ, σ)

→ **Funnel degeneracy**



Neal's funnel



$$\beta \sim \mathcal{N}(0, \sigma^2)$$
$$\alpha_i \sim \mathcal{N}(0, e^\beta),$$



Neal's funnel

$$\beta \sim \mathcal{N}(0, \sigma^2)$$
$$\alpha_i \sim \mathcal{N}(0, e^\beta),$$




- Need very different step-sizes in different regions
 - Small steps to probe the neck



Neal's funnel


$$\beta \sim \mathcal{N}(0, \sigma^2)$$
$$\alpha_i \sim \mathcal{N}(0, e^\beta),$$

- Need very different step-sizes in different regions
 - Small steps to probe the neck






Neal's funnel

$$\beta \sim \mathcal{N}(0, \sigma^2)$$
$$\alpha_i \sim \mathcal{N}(0, e^\beta),$$







- Need very different step-sizes in different regions
 - Small steps to probe the neck
 - Large steps to probe the mouth



Neal's funnel

$$\beta \sim \mathcal{N}(0, \sigma^2)$$
$$\alpha_i \sim \mathcal{N}(0, e^\beta),$$

- Need very different step-sizes in different regions
 - Small steps to probe the neck
 - Large steps to probe the mouth
- Constant mass matrix is insufficient
 - **Multiscale!** condition number changes
 - Very badly conditioned away from origin





Motivation for DRHMC

For the distributions that do not have globally optimal configurations for the transition kernel,
can we still benefit from different locally optimized transition kernels.*

(*when continuous adaptation is not feasible)



Delayed Rejection HMC

HMC

$$x \bullet \xrightarrow{F_1} \bullet y$$

Standard HMC

- Transition kernel F_1 : `n' leapfrog steps with step size `ε'

Delayed Rejection HMC


When faced with a rejection, make another proposal with a **different transition kernel** that has a better chance of getting accepted

Standard HMC


- Transition kernel F_1 : 'n' leapfrog steps with step size ϵ

2-stage DRHMC

HMC



(a) 2-stage DRHMC



Delayed Rejection HMC


When faced with a rejection, make another proposal with a **different transition kernel** that has a better chance of getting accepted

Standard HMC


$$\alpha(x, y) = \min \left(\frac{\pi(y)}{\pi(x)}, 1 \right)$$

2-stage DRHMC

HMC



(a) 2-stage DRHMC



Detailed balance:

the probability of transitioning from x to y is the same as the probability of transitioning from y to x

Delayed Rejection HMC


When faced with a rejection, make another proposal with a **different transition kernel** that has a better chance of getting accepted

Standard HMC


$$\alpha(x, y) = \min \left(\frac{\pi(y)}{\pi(x)}, 1 \right)$$

2-stage DRHMC

HMC



(a) 2-stage DRHMC



Detailed balance:

the probability of transitioning from x to y is the same as the probability of transitioning from y to x

Delayed Rejection HMC

When faced with a rejection, make another proposal with a **different transition kernel** that has a better chance of getting accepted


Standard HMC

$$\alpha(x, y) = \min \left(\frac{\pi(y)}{\pi(x)}, 1 \right)$$


2-stage DRHMC

$$\alpha_2(x, F_1(x), y) = \min \left(1, \frac{\pi(y) (1 - \alpha_1(y, F_1(y)))}{\pi(x) (1 - \alpha_1(x, F_1(x)))} \right)$$

HMC



(a) 2-stage DRHMC



Detailed balance:

the probability of transitioning from x to y is the same as the probability of transitioning from y to x

Delayed Rejection HMC

When faced with a rejection, make another proposal with a **different transition kernel** that has a better chance of getting accepted


Standard HMC

$$\alpha(x, y) = \min \left(\frac{\pi(y)}{\pi(x)}, 1 \right)$$


2-stage DRHMC

$$\alpha_2(x, F_1(x), y) = \min \left(1, \frac{\pi(y) (1 - \alpha_1(y, F_1(y)))}{\pi(x) (1 - \alpha_1(x, F_1(x)))} \right)$$

HMC



(a) 2-stage DRHMC



Detailed balance:

the probability of transitioning from x to y is the same as the probability of transitioning from y to x

Delayed Rejection HMC

When faced with a rejection, make another proposal with a **different transition kernel** that has a better chance of getting accepted

Standard HMC

$$\alpha(x, y) = \min \left(\frac{\pi(y)}{\pi(x)}, 1 \right)$$


2-stage DRHMC

$$\alpha_2(x, F_1(x), y) = \min \left(1, \frac{\pi(y) [1 - \alpha_1(y, F_1(y))]}{\pi(x) [1 - \alpha_1(x, F_1(x))]} \right)$$


3-stage DRHMC

$$\tilde{\alpha}_3(x) = \min \left[\frac{\pi(y) [1 - \tilde{\alpha}_1(y)] [1 - \tilde{\alpha}_2(y)]}{\pi(x) [1 - \tilde{\alpha}_1(x)] [1 - \tilde{\alpha}_2(x)]}, 1 \right]$$


HMC



(a) 2-stage DRHMC



(b) 3-stage DRHMC




DRHMC for Neal's funnel

Delayed proposals with reduced step size:

- Starting step size: ε_0
- Step size decreases with factor 'a'
 $\varepsilon_0, \varepsilon_0/a, \varepsilon_0/a^2, \dots, \varepsilon_0/a^k, \dots$

Two hyperparameters:

- a : factor of reduction
- k : number of delayed rejections



DRHMC for Neal's funnel

Delayed proposals with reduced step size:



- Starting step size: ϵ_0
- Step size decreases with factor 'a'
 $\epsilon_0, \epsilon_0/a, \epsilon_0/a^2, \dots, \epsilon_0/a^k, \dots$

Two hyperparameters:

- a: factor of reduction
- k: number of delayed rejections

DRHMC requires ~5-10x less gradient evaluations vs standard HMC

- Similar gains for other hard problems



Lower is better

DRHMC for Neal's funnel

Delayed proposals with reduced step size:



- Starting step size: ϵ_0
- Step size decreases with factor 'a'
 $\epsilon_0, \epsilon_0/a, \epsilon_0/a^2, \dots, \epsilon_0/a^k, \dots$

Two hyperparameters:


- a: factor of reduction
- k: number of delayed rejections

DRHMC requires ~5-10x less gradient evaluations vs standard HMC

- Similar gains for other hard problems



Delayed proposals made as we move into the neck of the funnel.




Lower is better

Cost of DRHMC

Number of proposals (density evaluations) for k^{th} order grows as 2^{k-1}

(b) 3-stage DRHMC




Cost of DRHMC

Number of proposals (density evaluations) for k^{th} order grows as 2^{k-1}

But, the dominant cost in HMC is number of gradient evaluations, leapfrog steps!

(b) 3-stage DRHMC



Cost of DRHMC

Number of proposals (density evaluations) for k^{th} order grows as 2^{k-1}

But, the dominant cost in HMC is number of gradient evaluations, leapfrog steps!


DRHMC

- Starting step size: ϵ
- Step size decreases with factor 'a'
- $\epsilon, \epsilon/a, \epsilon/a^2, \dots, \epsilon/a^k$

HMC

- Largest **stable** step size: ϵ_0
 $\epsilon_0 \sim \epsilon/a^{k-1}$

(b) 3-stage DRHMC



Cost of DRHMC

Number of proposals (density evaluations) for k^{th} order grows as 2^{k-1}

But, the dominant cost in HMC is number of gradient evaluations, leapfrog steps!

DRHMC

- Starting step size: ε
- Step size decreases with factor ' a '
- $\varepsilon, \varepsilon/a, \varepsilon/a^2, \dots, \varepsilon/a^k$


HMC

- Largest **stable** step size: ε_0
 $\varepsilon_0 \sim \varepsilon/a^{k-1}$

Integration time for every proposal: $T = n\varepsilon$

- Total number of leapfrog steps for HMC $\sim T/\varepsilon_0 \sim na^{k-2}$

(b) 3-stage DRHMC



Cost of DRHMC

Number of proposals (density evaluations) for k^{th} order grows as 2^{k-1}

But, the dominant cost in HMC is number of gradient evaluations, leapfrog steps!

DRHMC

- Starting step size: ϵ
- Step size decreases with factor 'a'
- $\epsilon, \epsilon/a, \epsilon/a^2, \dots, \epsilon/a^k$

HMC


- Largest **stable** step size: ϵ_0
 $\epsilon_0 \sim \epsilon/a^{k-1}$

Integration time for every proposal: $T = n\epsilon$

- Total number of leapfrog steps for HMC $\sim T/\epsilon_0 \sim na^{k-2}$
- Total number of leapfrog steps for DRHMC*
 (*in the worst case)

$$: n \underset{(\epsilon)}{2^{k-1}} + na \underset{(\epsilon/a)}{2^{k-2}} + \dots + na^{k-1} \underset{(\epsilon/a^k)}{1} \quad (GP)$$

(b) 3-stage DRHMC



Cost of DRHMC

Number of proposals (density evaluations) for k^{th} order grows as 2^{k-1}

But, the dominant cost in HMC is number of gradient evaluations, leapfrog steps!

DRHMC

- Starting step size: ϵ
- Step size decreases with factor ' a '
- $\epsilon, \epsilon/a, \epsilon/a^2, \dots, \epsilon/a^k$

HMC


- Largest **stable** step size: ϵ_0
 $\epsilon_0 \sim \epsilon/a^{k-1}$

Integration time for every proposal: $T = n\epsilon$

- Total number of leapfrog steps for HMC $\sim T/\epsilon_0 \sim na^{k-2}$
- Total number of leapfrog steps for DRHMC*
 (*in the worst case)

$: n \times 2^{k-1} + na \times 2^{k-2} + \dots + na^{k-1} \times 1$ (GP)	
$= nka^{k-1}$	if $a = 2 \rightarrow O(ak)$ more expensive than HMC
$= O(a^{k-1}n)$	if $a > 2 \rightarrow O(a)$ more expensive than HMC, Independent of k !

(b) 3-stage DRHMC





Are delayed proposals always beneficial?

Say you reject a proposal when $\alpha = 0.9$
Should you make a delayed proposal?



Are delayed proposals always beneficial?

Say you reject a proposal when $\alpha = 0.9$

Should you make a delayed proposal?

Not all rejections are bad



Are delayed proposals always beneficial?

Say you reject a proposal when $\alpha = 0.9$

Should you make a delayed proposal?

Not all rejections are bad

Probabilistic DRHMC: Make the next proposal with probability

$$p_{j+1}(x) = 1 - \tilde{\alpha}_j(x)$$

$$\tilde{\alpha}_2(x) = \min \left[\frac{\pi(y)[1 - \tilde{\alpha}_1(y)]p_2(y, F_1(y))}{\pi(x)[1 - \tilde{\alpha}_1(x)]p_2(x, F_1(x))}, 1 \right]$$

Are delayed proposals always beneficial?

Say you reject a proposal when $\alpha = 0.9$
 Should you make a delayed proposal?
 Not all rejections are bad


Probabilistic DRHMC: Make the next proposal with probability

$$p_{j+1}(x) = 1 - \tilde{\alpha}_j(x)$$


$$\tilde{\alpha}_2(x) = \min \left[\frac{\pi(y)[1 - \tilde{\alpha}_1(y)]p_2(y, F_1(y))}{\pi(x)[1 - \tilde{\alpha}_1(x)]p_2(x, F_1(x))}, 1 \right]$$

- Reduces the cost of DRHMC by only making delayed proposals only when needed
- Increases robustness to fitting step-size!

Stochastic
Volatility



(a) Delayed Rejection



(b) Probabilistic Delayed Rejection



Variants of DRHMC

Does not necessarily need to reduce step size

- Different integrators (higher order leapfrog, implicit midpoint)
- Different kinetic energy
- Different mass matrix



Variants of DRHMC

Does not necessarily need to reduce step size

- Different integrators (higher order leapfrog, implicit midpoint)
- Different kinetic energy
- Different mass matrix

Extensions for DRHMC

WIP with Gilad Turok and Bob Carpenter

- Auto-tuning DRHMC hyper-parameters
- Continuous adaptation: Combine DR + Generalized HMC (partial momentum refresh)



Takeaways

- Delayed rejection HMC for pathological distributions (multiscale distributions like funnel)
 - benefit from multiple, locally optimized transition kernels
- Unlike DR for Metropolis Hastings, cost of a well-tuned DRHMC is a constant factor more than a stable HMC
 - if adapting step size
- Probabilistic DRHMC makes proposals probabilistically, and reduces the cost of DRHMC
- More stable to tuning parameters of HMC (for e.g. step size)
- Any valid transition kernels can be combined into delayed proposals



Takeaways

- Delayed rejection HMC for pathological distributions (multiscale distributions like funnel)
 - benefit from multiple, locally optimized transition kernels
- Unlike DR for Metropolis Hastings, cost of a well-tuned DRHMC is a constant factor more than a stable HMC
 - if adapting step size
- Probabilistic DRHMC makes proposals probabilistically, and reduces the cost of DRHMC
- More stable to tuning parameters of HMC (for e.g. step size)
- Any valid transition kernels can be combined into delayed proposals


Thank you




Extra slides

Other experiments


No ideal global proposal!



Mixture of 2 Gaussians with different scales ($\sigma = 0.1, 1$)
~2x gains



Eight school model- hierarchical, mildly multi-scale
~3x gains



Gull's lighthouse: poor data, ill-defined prior, Cauchy posterior
~5x gains

DRHMC for Neal's funnel


Delayed proposals with reduced step size:

- Starting step size: ϵ_0
- Step size decreases with factor 'a'
 $\epsilon_0, \epsilon_0/a, \epsilon_0/a^2, \dots, \epsilon_0/a^k, \dots$

Two hyperparameters:

- a: factor of reduction
- k: number of delayed rejections

Similar gains for other hard problems



Comparing cost of DRHMC vs standard HMC
 ~5-10x gains



Delayed Rejections (DR)

Delayed Rejection : when faced with a rejection, delay it.
Try to make more proposals which might get accepted

Well studied in the context of random-walk Metropolis sampling
Mira 1998, Mira & Tierney 99, Greene & Mira 2001

$$K(x, dy) = \alpha(x, y)Q(x, dy) + r(x)\delta_x(dy)$$

$$\alpha(x, y) = \min\left(\frac{\pi(y)q(y, x)}{\pi(x)q(x, y)}, 1\right)$$

$$K(x, dy) = Q_1(x, dy)\alpha_1(x, y) + \int_{s \in S} Q_1(x, ds)[1 - \alpha_1(x, s)] \underbrace{[Q_2(x, s, dy)\alpha_2(x, s, y) + r_2(x, s)\delta_x(dy)]},$$

$$\alpha_2(x, s, y) = \min\left(\frac{\pi(y)q_2(y, s, x)q_1(y, s)[1 - \alpha_1(y, s)]}{\pi(x)q_2(x, s, y)q_1(x, s)[1 - \alpha_1(x, s)]}, 1\right)$$



Other DR methods in the literature

Extra Chance HMC : when turned down, keep moving on
Sohl-Dickstein et al., 2014; Campos and Sanz-Serna, 2015