



Plagiarism Checker X - Report

Originality Assessment

8%



Overall Similarity

Date: Jan 27, 2024

Matches: 100 / 1307 words

Sources: 8

Remarks: Low similarity detected, check with your supervisor if changes are required.

Verify Report:

Scan this QR Code



Matrix Representation of Graph Theory in Hydrocarbons

Amit Kumar Gupta^{1,3*}, Kumari Pinky^{1,3} and Shalini Gupta^{2,3**}

¹Department of Mathematics, K. C. M. T, Bareilly-243122, Uttar Pradesh, India

²Department of Chemistry, Bareilly College, Bareilly-243005, Uttar Pradesh, India

³M. J. P. Rohilkhand University, Bareilly-243006, Uttar Pradesh, India.

Corresponding Author E-mail: * guptaamit48@rediffmail.com; **

shalu123gupta21@gmail.com

Abstract: Graph theory is one of the best-known research head subjects having many applications. Graph theory serves as a mathematical model to represent any system which has a binary relation. chemical graph theory is branch of graph theory that is concerned with analysis of all consequences of connectivity in a chemical graph. An organic molecule can be represented by a graph, which can be converted to several matrices by using various graph characteristic, connecting of atoms through bonds lead to adjacency, incidence, cut-set, circuit and distance matrices. The object of this paper is to adopt graph theory matrix approach in hydrocarbons.

Index Terms: Matrix (Adjacency, cycle, cut-set, edge, incidence and path) and Undirected graph.

1. Introduction

The graph theory is the study of graph, which is a collection of vertices or nodes and edges that connect pair of vertices and is a branch of graph theory that applies geometry to derive two-dimensional representation of graphs. Graph theory matrix approach is used in modeling and solving decision making problem with multiple and interrelated attributes.

1.1 Graph theoretical terms and chemical terms

Chemical system may be depicted by chemical graphs using a simple conversion rule,

Site \leftrightarrow Vertex

connection \leftrightarrow Edge

1.2 Mapping of graph theoretical and chemical terms

Graph theoretical terms Chemical terms

- ☐ Chemical/ Molecular graph \rightarrow Structural formula
- ☐ Vertex \rightarrow Atom
- ☐ Weighted vertex \rightarrow Atoms of a specified element
- ☐ Edge \rightarrow Chemical bond
- ☐ Weighted Edge \rightarrow Chemical bond between specified element
- ☐ Chain \rightarrow Linear Alkanes/Alkenes/Alkynes
- ☐ Cycle \rightarrow Cycloalkanes

2. Matrix representation of structural graphs

The purpose of matrix representation of structural graph is the reflecting molecular topology and correlating structure. Graphs are converted into a mathematical expression which may be matrix and a polynomial. The first representation for characterizing a graph and discussed below.

2.1 Adjacency Matrix

Adjacency matrix of a graph with n -vertices and no parallel edges, self-loops are allowed is an $n \times n$ order matrix $A=(a_{ij})$

where,

$a_{ij} =$

is weighted adjacency matrix of Aliphatic Hydrocarbons. The valency of carbon is 4 and valency of Hydrogen is 1.

2.2 Incidence Matrix

Let $G=(V, E)$ be graph with n -vertices (v_1, v_2, \dots, v_n) and m -edges (e_1, e_2, \dots, e_m) and no Self-Loop. The matrix $A=(a_{ij})$ of order $n \times m$, where,

$a_{ij} =$

for a given graph $G = (V, E)$, an incidence matrix depends on the ordering of the vertices of G and that edge of G . For different ordering of vertices and edges we get different incident matrix of the same graph G .

2.3 Path Matrix

A path matrix is defined for a specific pair of vertices in a graph say (u, v) and is denoted by $P(u, v)$.

Let 'm' be the number of different paths between the vertices 'u' and 'v', 'n' be the number of edges path between the vertices (u, v) .

then path matrix will be,

$$P(u, v) = (P_{ij}) =$$

The path matrix of any two vertices of alkanes is row matrix. At last, one of the path matrix of any two carbons in alkanes is two rows. At last, one of the path matrix of any two carbons alkynes are three rows.

2.4 Cycle Matrix

Let G be graph with 'm' edges and 'n' different circuits then a cycle matrix $C = (C_{ij})$ of order $n \times m$ is defined as

$$C_{ij} =$$

The cycle matrix is denoted by $C(G)$. The cycle matrix of aliphatic hydrocarbons as

Cycle matrix of Alkanes does not exist.

Cycle matrix of Alkenes is 1.

Cycle matrix of Alkynes is 2.

2.5 Edge Cut-Set Matrix: -

Let the graph G have 'm' edge and 'n' be the number of different Cut-set in G . The Edge Cut-set matrix $E(G)$ is given by

$$E(G) = (E_{ij})_{n \times m} \text{ where } E_{ij} =$$

The number of Edge cut-sets of alkanes are 4, 7, 10, ..., $3n+1$, alkenes are 5, 8, 11, ..., $3n+2$ and alkynes are 3, 6, 9, ..., $3n$.

3. Hydrocarbons

Hydrocarbons in an organic compound made of carbons and hydrogens. It is possible for single (–), double (=) and triple (≡) bonds to form carbene atoms and hydrogens. 3

Aliphatic hydrocarbons are compound of hydrogen and carbon that do not contain benzene ring. There are three types alkanes, alkene and alkynes. The saturated hydrocarbons known

as alkanes consist 8 only carbon and hydrogen connected by single bond. The chemical

formula is C_nH_{2n+2} where n represent the number of carbons and $2n+2$ represents

number of hydrogens. The unsaturated hydrocarbons are 3 alkenes and alkynes. 1

Alkenes are hydrocarbons that contain $C=C$, chemical formula for alkene is

C_nH_{2n} . Alkynes are hydrocarbons that contains at least one triple bond between two

carbon atoms, the number of hydrogen atoms is still less in alkynes as compared to

alkanes or alkenes. The chemical formula of alkynes is C_nH_{2n-2} .

4. Different types of graphs matrix representation in aliphatic hydrocarbons

4.1. Adjacency Matrices

Table-4.1

Hydrocarbons

Structure

Structure into a graph

Adjacency Matrix

Methane (CH_4)

(Alkane)

Ethene (C_2H_4)

(Alkene)

Ethyne (C_2H_2)

(Alkyne)

4.2. Incidence Matrix

Table-4.2

Hydrocarbons

Structure

Structure into a graph

Adjacency Matrix

Methane (CH_4)

(Alkane)

Ethene (C₂H₄)

(Alkene)

Ethyne (C₂H₂)

(Alkyne)

4.3. Path matrix

Table-4.3

Hydrocarbons

Structure

Structure into a graph

Adjacency Matrix

Methane (CH_4)

(Alkane)

6 Path matrix of Methane with vertices C,H1 whose order is 1×4 is denoted by

Ethene (C_2H_4)

(Alkene)

Path matrix of ethane with vertices C1 and C2 whose order is 2×6 is denoted by

Ethyne (C_2H_2)

(Alkyne)

Path matrix of ethane with vertices C1 and C2 whose order is 3×6 is denoted by

4.4. 4 Edge Cut Set matrix

Table-4.4

Hydrocarbons

Structure

Structure into a graph

Adjacency Matrix

Methane (CH_4)

(Alkane)

Edge Cut-sets of methane are $C1=\{e1\}$, $C2=\{e2\}$, $C3=\{e3\}$ and $C4=\{e4\}$

So, the Edge Cut Set matrix of methane is of order 4×4 is

Ethene (C_2H_4)

(Alkene)

Edge Cut-sets of Ethene are $C1=\{e1\}$, $C2=\{e2\}$, $C3=\{e3\}$, $C4=\{e4\}$ and $C5=\{e5,e6\}$

So, the Edge Cut Set matrix of methane is of order 5×6 is

Ethyne (C_2H_2)

(Alkyne)

Edge Cut-sets of Ethyne are $C_1=\{e_1\}$, $C_2=\{e_2\}$, $C_3=\{e_3, e_4, e_5, e_6\}$

So, the ⁴ Edge Cut Set matrix of methane is of order 3×5 is

4.5. Cycle matrix

Table-4.5

Hydrocarbons

Structure

Structure into a graph

Adjacency Matrix

Methane (CH_4)

(Alkane)

Alkanes does not exit cycle matrix.

Ethene (C₂H₄)

(Alkene)

Ethene has one cycle $C_1=\{e_5,e_6\}$. In general alkenes ³ has at least one cycle matrix. The cycle matrix of ethene is of order 1×4 as

Ethyne (C₂H₂)

(Alkyne)

Cycle of Ethyne are $C_1=\{e_1,e_2\}$, $C_2=\{e_4,e_5\}$ and $C_3=\{c_3,c_5\}$. The cycle matrix of the Ethyne is of order 3×5 as

5. Conclusion

Group theory is used in the field of organic chemistry to model chemical compound in the

form of graph where vertices represent the sequences in hydrocarbons and an edge will be drawn between two vertices. In this paper the major role of graph theoretic approach to find the matrix representation in hydrocarbons.

References

- [1] O.P. Gandhi, and V.P. Agrawal; FMEA—A diagraph and matrix approach, Reliability Engineering & System Safety; Volume 35, Issue 2, 147-158 (1992). [https://doi.org/10.1016/0951-8320\(92\)90034-I](https://doi.org/10.1016/0951-8320(92)90034-I)
- [2] R.V. Rao, Graph theory and matrix approach for the performance evaluation of technical institutions, Indian J of Tech Educ, 23 (2) 27-33 (2000).
- [3] M. Darvish, M. Yasaei and A. Saeedi, Application of **7 graph theory and matrix methods to contractor ranking**, Int. J Proj. Manag, 27, 610-619 (2009).
- [4] V. Paramasivam, V. Senthil, and N. Rajam Ramasamy, **2 Decision making in equipment selection: an integrated approach with digraph and matrix approach, AHP and ANP**, The international journal of manufacturing technology, 54 (2010) 1233-1244.
- [5] J.A. **5 Bondy and U.S.R Murty, Graph theory with** application; Machmillan Press Ltd., 1976.
- [6] Nar Singh Deo- **Graph theory with applications** to engineering and computer science, (PHI Publication).
- [7] Murgan. M- Application of algebraic concept of hydrocarbons in graph theory.
- [8] Kenneth. H. Rosen- Discrete mathematics and its application, McGraw Hill.

Sources

1	http://chilbrtxt.org/our/BrdgVll_y_o_unty_nd_Tchncl_oll_g/Fundntl_of_h_try/11:_rgnc_htry/1104:_lkn_nd_lkyn INTERNET 2%
2	http://r_rchgtn_t/profl/V-P_r_v/publcton/2_126_2_1_Dc_on_kng_n_qupnt_lcton_ntgrtd_pproch_th_dgrph_nd_trx_pproch_P_nd_NP/lnk/fc2_2_bf14088_f6_1/Dc_on_kng_n-qupnt_lcton_ntgrtd_pproch_th_dgrph_nd_trx_pproch_P_nd_NP.pdf INTERNET 1%
	http://chilbrtxt.org/Bookh_lv/Introductory_htry/htry_for_hngng_T_ll_nd_Mc_rry)/0:_rgnc_htry/02:_lph_t_c_ydroc_rbon INTERNET 1%
4	http://cnfrco/Grph-Thory/cut-dg-cut-t-cut-vrtx-n-grph-connectvty INTERNET 1%
	http://rouontrlc/~h hn/IFT_4/GTW.pdf INTERNET 1%
6	http://tckovrfloco/qu_ton/1_8602_2/co_putton-of-pth-trx-fro-th-djcncy-trx INTERNET 1%
	http://ntccholrorg/ppr/pplcton-of-th-grph-thory-nd-trx-thod-Drvh-Y/08b2460111db62f6cbf40cb280b0 INTERNET 1%
8	http://opntxorg/book/orgnc-chtry/pg/-2-lkn-nd-lkn-or#:~:txt=lknroftndcrbd_turtddhydroc_rbon:hydroc_rbonn_2n+2_hrn_nntgr INTERNET 1%

EXCLUDE CUSTOM MATCHES OFF

EXCLUDE QUOTES OFF

EXCLUDE BIBLIOGRAPHY OFF