- 22. Novzurov, E.N., Ismailov, N.M., and Mamaedov, S.Sh. (1983). Phenolic compounds from leaves of *Hippophae rhamnoides*. *Rastit. Resur.* 39, 354.
- 23. Aslanov, S.M. (1982). Chemical constituents of fruit of Hippophae growing in Apsherone. Rastit. Resur. 18, 73.
- 24. Jmirko T.G., Goncharova, N.P., Gigienova, E.I., and Glushenkova, A.I. (1984). Neutral lipids in the fruit oil of *Hippophae rhamnoides*. *Khim. Prir. Soedin* 300.
- 25. Novzurov, E.N. (1981). Carotenoids and sterols from Hippophae rhamnoides. Khim. Prir. Soedin 98.
- 26. Jmirko T.G., Rashkes, Ya.V., and Glushenkova, A.I. (1986). Oxyacids in the fruit oil of *Hippophae rhamnoides*. *Khim. Prir. Soedin* 161.
- 27. Baldandorj, D., Khandsuren, S., and Jargal, Ya. (1985). Fruit of *Hippophae rhamnoides* L. Mongolian National Standard 916–85.
- 28. Ulziikhutag, A. (1969). Influence of Hippophae oil on atherosclerosis. (p. 18). A thesis submitted for the degree of Doctor of Philosophy in Medicine. Moscow: Academy of Medical Science, Institute of Nutrition.
- 29. Jamyansan, Ya. (1973). Bioactive compounds from the fruit of *Hippophae* growing in Mongolia and its use. (p. 32). A thesis submitted for the degree of Doctor of Philosophy in Chemistry. Moscow: Institute of Vitamin Research and Investigation.