

## References:

1. Olziikhutag, N. (Ed). (1983). Latin-Mongolian-Russian Dictionary of Vascular Plants of Mongolia (p. 258). Ulaanbaatar: Press of Mongolian Academy of Sciences.
2. Gubanov, I.A. (1996). Conspectus on Mongolian Flora (vascular plants) (p. 107). Moscow: Valang Press.
3. Malishev, L.I., and Peshkova, G.A. (1979). Flora of Central Siberia (Vol. 2, p. 840). Novosibirsk: Science Printing.
4. Sanchir, Ch., Batkhuu, J., Boldsaikhan, B., and Komatsu, K. (2003). Illustrated Guide of Mongolian Useful Plants. (Vol. 1, p. 64). Ulaanbaatar: Admon Printing.
5. Ligaa, U., Davaasuren, B., and Ninjil, N. (2005). Medicinal Plants of Mongolia Used in Western and Eastern Medicine. (p. 238). Ulaanbaatar: JCK Printing.
6. Yuthok Yonten Gonpo., Four Medical Tantras, VIII-IXth century.
7. Chandra, A., Misra, L.N., and Thakur, R.S. (1987). Germacranolides and an alkyl glucoside from *Tanacetum vulgare* L. *Phytochemistry* 26, 1463.
8. Dembitski, A.D., Krotova, G.I., Urina, R.A., Suleeva, R. (1984). Essential oil from *Tanacetum vulgare* L. *Khim. Priir. Soedin.* 716.
9. Vaverkova, S., Mikulasova, M., Haban, M., and Sloboda, P. (2006). A study of qualitative properties of the essential oil of *Tanacetum vulgare* L. *Ceska Slov. Farm.* 55, 181.
10. Grabarczyk, H., Drozd, B., and Mozdzanowska, A. (1973). Sesquiterpene lactones: Lactones in aerial parts of *Tanacetum vulgare* L. *Pol. J. Pharmacol. Pharm.* 25, 95; *Chem. Abstr.* (1973), 78, 156612.
11. Bloszyk, E., and Drozd, B. (1978). Sesquiterpene lactones in species of the genus *Chrysanthemum*. *Acta Soc. Bot. Pol.* 47, 3.
12. Unusov, A.I., Sidyakin, Sh.P., Nigmatullaev, A.M. (1979). Sesquiterpene lactones of *Tanacetum vulgare* L. *Khim. Priir. Soedin.* 101.
13. Nano, G.M., Appendino, G., Bicchi, C., and Frattini, C. (1980). Wild Piedmontese plants: On chemotypes of *Tanacetum vulgare* L., containing sesquiterpene lactones with germacrane skeleton. *Fitoterapia* 51, 135.
14. Appendino, G., Gariboldi, P., and Nana, G.M. (1983). Tanacetols A and B, non-volatile sesquiterpene alcohols from *Tanacetum vulgare* L. *Phytochemistry* 22, 509.
15. Calleri, M., Chiari, G., and Viterbo, D. (1983). The structure of tanacetols A, C<sub>17</sub>H<sub>26</sub>O<sub>4</sub>, and B, C<sub>19</sub>H<sub>30</sub>O<sub>5</sub>, two new sesquiterpene alcohols from *Tanacetum vulgare* L. *Acta Crystallogr. C* 39, 758.
16. Chandler, R.F., Hooper, S.N., Hooper, D.L., Jamieson, W.D., and Lewis, E. (1982). Herbal remedies of the Maritime Indians: Sterol and triterpenes of *Tanacetum vulgare* L. *Lipids* 17, 102.
17. Appendino, G., Valle, M.G., and Nana, G.M. (1982). On a new chemotype of *Tanacetum vulgare* L. *Fitoterapia* 53, 115.
18. Sokolov, P.D. *et al.* (1989). Plants Review of USSR: Family Asteraceae. (p. 189). Leningrad: Science Printing.