

Habitat: Rocky areas along river banks [2–4].

Part used: Shoot

Traditional Uses: The taste is sour and sweet, and the potency is blunt and cool. It is used for the following: treating fever and poisoning. It is an ingredient of the following traditional prescriptions: Braivu-3, Gagol-18, Gontog-7, Khach gurgum -25, Degd-13, and Dedbo-10 [4–8].

Chemical constituents: Flavonoids: quercetin, rhamnetin, tamarisketin, and 3-O-beta-D-quercetin glucopyranoside (isoquercitrin) [9].

References:

1. Olziikhutag, N. (Ed). (1983). Latin-Mongolian-Russian Dictionary of Vascular Plants of Mongolia (p. 146). Ulaanbaatar: Press of Mongolian Academy of Sciences.
2. Gubanov, I.A. (1996). Conspectus on Mongolian Flora (vascular plants) (p. 76). Moscow: Valang Press.
3. Malishev, L.I., and Peshkova, G.A. (1979). Flora of Central Siberia (Vol. 2, p. 657). Novosibirsk: Science Printing.
4. Ligaa, U., Davaasuren, B., and Ninjil, N. (2005). Medicinal Plants of Mongolia Used in Western and Eastern Medicine. (p. 77). Ulaanbaatar: JCK Printing.
5. Yuthok Yonten Gonpo., Four Medical Tantras, VIII-IXth century.
6. Danzanpuntsag., Crystal rosary. XVIIIth century.
7. Boldsaikhan, B. (2004). Encyclopedia of Mongolian Medicinal Plants (p. 111). Ulaanbaatar: Mongolian University of Science and Technology.
8. Khurelchuluun, B., Suran, D., and Zina, C. (2007). Illustrated Guide of Raw Materials Used in Traditional Medicine. (p. 190). Ulaanbaatar: Erkhes Printing.
9. Semenova, L.S. (1993). Flavonoid composition of shoots of *Myricaria longifolia* (Willd.) Ehrenb. *Rastit. Resur.* 29, 40.