

obovate or ovate-lanceolate, longer than secondary umbels, acute. Mericaps oblong, with narrowly winged ribs, two longitudinal furrows on commissure.

Distribution: Khent., Mong-Dag.

Habitat: Larch and birch forests, their fringes in mountain forest-steppe belt [2,3].

Part used: Herb

Traditional Uses: The taste is bitter and the potency is light, severe, and sharp. It is used for the following: as an anti-parasitic agent. It is also said to be beneficial for stomach diseases, and for colds. It is an ingredient of the following traditional prescription: Zadi-15 [4,5].

Chemical constituents: Essential oil: myrcene, α -pinene, β -pinene, limonene, *n*-cymol, phellandrene, sabinene, camphene, carvone, linalool, bornyl acetate, pinocamphene, perolidol and others [6], flavonoids: quercetin, isorhamnetin, rutin [7].

Bioactivity: Antitumor [8].

References:

1. Olziikhutag, N. (Ed). (1983). Latin-Mongolian-Russian Dictionary of Vascular Plants of Mongolia (p. 204). Ulaanbaatar: Press of Mongolian Academy of Sciences.
2. Gubanov, I.A. (1996). Conspectus on Mongolian Flora (vascular plants) (p. 79). Moscow: Valang Press.
3. Malishev, L.I., and Peshkova, G.A. (1979). Flora of Central Siberia (Vol. 2, p. 676). Novosibirsk: Science Printing.
4. Yuthok Yonten Gonpo., Four Medical Tantras, VIII-IXth century.
5. Danzanpuntsag., Crystal rosary. XVIIIth century.
6. Tserendulam, L. (2002). Chemical and chemical-technological investigation of *Bupleurum* species. (p. 53). A thesis submitted for the degree of Doctor of Philosophy in Pharmacy. Ulaanbaatar: Medical University of Mongolia.
7. Sokolov, P.D. *et al.*, Plants Review of USSR: Family Rutaceae-Elaeagnaceae. (p. 86). Leningrad: Science Printing.
8. Cetlin, A.L., Niconov, G.K., Shvarev, I.F., and Pimenov, M.G. (1965). On the antitumor activity of natural coumarins. *Rastit. Resur.* 1, 507.