

References:

1. Olziikhutag, N. (Ed). (1983). Latin-Mongolian-Russian Dictionary of Vascular Plants of Mongolia (p. 205). Ulaanbaatar: Press of Mongolian Academy of Sciences.
2. Gubanov, I.A. (1996). Conspectus on Mongolian Flora (vascular plants) (p. 79). Moscow: Valang Press.
3. Malishev, L.I., and Peshkova, G.A. (1979). Flora of Central Siberia (Vol. 2, p. 678). Novosibirsk: Science Printing.
4. Sanchir, Ch., Batkhuu, J., Boldsaikhan, B., and Komatsu, K. (2003). Illustrated Guide of Mongolian Useful Plants. (Vol. 1, p. 25). Ulaanbaatar: Admon Printing.
5. Ligaa, U., Davaasuren, B., and Ninjil, N. (2005). Medicinal Plants of Mongolia Used in Western and Eastern Medicine. (p. 134). Ulaanbaatar: JCK Printing.
6. Ganbayar, Ya. (2001). Guide to Drug Prescriptions of Mongolia. (p. 123). Ulaanbaatar: Bit Service Printing.
7. Khurelchuluun, B., Suran, D., and Zina, C. (2007). Illustrated Guide of Raw Materials Used in Traditional Medicine. (p. 32). Ulaanbaatar: Erkhes Printing.
8. Max Wichtl. (1994). Herbal Drugs and Phytopharmaceuticals. (p. 123). Stuttgart: Medpharm, Stuttgart.
9. Sokolov, P.D. *et al.* (1988). Plants Review of USSR: Family Rutaceae-Elaeagnaceae. (p. 92). Leningrad: Science Printing.
10. Embong, M.B., Hadzigeve D., and Molnar, S. (1977). Essential oils from species grown in Alberta: Caraway oil (*Carum carvi*). *Can. J. Plant. Sci.* 57, 543.
11. Ikeda, R.M., Stanley, W.L., Vannier, S.H., and Spitler, E.H. (1962). The monoterpene hydrocarbon composition of some essential oils. *J. Food. Sci.* 27, 455.
12. Karrer, W. (1958). Konstitution und Vorkommen der organischen Pflanzenstoffe. (pp. 1207). Basel: Birkhäuser Verlag.
13. Plouvier, V. (1957). Sur la recherche du pinitol chez quelques Caryophyllacees, Magnoliacees et plantes de familles voisines. *Compt. Rend. Acad. Sci.* 244, 382.
14. Rothbächer, H., and Suteu, F. (1975). Über Hydroxylverbindungen des Kümmelöls. *Planta Med.* 28, 112.
15. Salveson, A., and Baerheim Svendsen, A. (1978). Oxygen-containing monoterpenes: Gas chromatographic separation and identification of cumin oil constituents. *Sci. Pharm.* 46, 93; *Chem. Abstr.* (1978). 89, 160134.
16. Schantz, M., and Ek, B.S. (1971). Über die Bildung von ätherischen Öl in Kümmel-*Carum carvi*. *Sci. Pharm.* 39, 82.
17. Schantz, M., and Huhtikengas, A. (1971). Über die Bildung von Limonen und Carvon in Kümmel-*Carum carvi*. *Phytochemistry* 10, 1787.
18. Kartnig, Th. (1969). Über einige Lipoid-Inhaltsstoffe aus den Früchten von *Pimpinella anisum* L. und *Carum carvi* L. Fette, Seifen, Anstrichmittel, 71, 276.
19. Dirks, U., and Herman, K. (1984). 4-(β -D-Glucopyranosyloxy)-benzoic acid, a characteristic phenolic constituent of the Apiaceae. *Phytochemistry* 23, 1811.
20. Hondelmann, W. (1985). Das Vorkommen einer ungewöhnlichen Fettsäure, der Petroselinsäure, in der Familie der Doldengewächse als Ausgangspunkt für die Entwicklung neuer Ölfruchte. Land-bauforsch. Völkendrode, 35, 185.
21. Kleiman, R., and Spencer, G.F. (1982). Search for new industrial oils: Umbelliferae seed oils rich in petroselinic acid. *J. Amer. Oil. Chem. Soc.* 59, 29.
22. Stepananco, N.A., Gusakova, S.D., and Umarov, A.U. (1980). Lipids from seed of *Carum carvi* and *Foeniculum vulgare*. *Khim. Prir. Soedin.* 827.
23. Volodya, Ts., Tserenbaljid, D., and Lamjav, Ts. (2008). Medicinal Plants of Mongolia. (p. 87). Ulaanbaatar.
24. Mashkovsi, M.D (1994). Medicinal Preparations. (p. 431). Moscow: Medicine Printing.