

**Distribution:** Khovs., Khent., Khang., Mong-Dag., Khyang., Khovd (Zuslan), Mong. Alt. (Taishir uul), Gobi-Alt. (Ikh Bogd, Baga Bogd).

**Habitat:** Larch forests, riverbank meadows, forest fringes [2–5].

**Parts used:** Herb and flowers

**Traditional Uses:** The taste is sweet and bitter, and the potency is cool. It is used for the following: wound healing, cardiovascular diseases, blood disorders, lymph diseases, and eye diseases. It is an ingredient of the following traditional prescription: Dugseltan [5–7].

**Chemical constituents:** Herb contains alkaloids, flavonoids; flowers contain alkaloids, vitamin C, flavonoids, coumarins [8].

**Bioactivity:** Antibacterial [8].

#### References:

1. Olziikhutag, N. (Ed). (1983). Latin-Mongolian-Russian Dictionary of Vascular Plants of Mongolia (p. 84). Ulaanbaatar: Press of Mongolian Academy of Sciences.
2. Gubanov, I.A. (1996). Conspectus on Mongolian Flora (vascular plants) (p. 51). Moscow: Valang Press.
3. Malishev, L.I., and Peshkova, G.A. (1979). Flora of Central Siberia (Vol. 2, p. 338). Novosibirsk: Science Printing.
4. Sanchir, Ch., Batkhuu, J., Boldsaikhan, B., and Komatsu, K. (2003). Illustrated Guide of Mongolian Useful Plants. (Vol. 1, p. 206). Ulaanbaatar: Admon Printing.
5. Ligaa, U., Davaasuren, B., and Ninjil, N. (2005). Medicinal Plants of Mongolia Used in Western and Eastern Medicine. (p. 178). Ulaanbaatar: JCK Printing.
6. Yuthok Yonten Gonpo., Four Medical Tantras, VIII-IXth century.
7. Danzanpuntsag., Crystal rosary. XVIIIth century, p. 255.
8. Fedorov, A.A. *et al.* (1985). Plants Review of USSR: Family Magnoliaceae-Limoniaceae. (p. 97). Leningrad: Science Printing.