

Distribution: Khent., Khang., Mong-Dag., Khovd, Mong. Alt., Dund. Khalkh, Ikh n., Dor. Mong, Gobi-Alt., Zyyngar.

Habitat: Rocky and stony slopes, among boulders, dry light forests, nomad camps, inhabited areas [1–5].

Parts used: Herb

Traditional Uses: The taste is bitter and the potency is hot and oily. It is used for the following: wound healing, treating lymph disease, edema, diabetes, scurvy, and lupus erythematosus; also beneficial in nervous diseases [5–7].

Chemical constituents: Organic acids, ascorbic acid, carotene, phenol carboxylic acids, flavonoids [8].

Bioactivities: Haemostatic, anti-inflammatory, diuretic, and antipyretic [8].

References:

1. Olziikhutag, N. (Ed). (1983). Latin-Mongolian-Russian Dictionary of Vascular Plants of Mongolia (p. 120). Ulaanbaatar: Press of Mongolian Academy of Sciences.
2. Gubanov, I.A. (1996). Conspectus on Mongolian Flora (vascular plants) (p. 37). Moscow: Valang Press.
3. Malishev, L.I., and Peshkova, G.A. (1979). Flora of Central Siberia (Vol. 2, p. 273). Novosibirsk: Science Printing.
4. Sanchir, Ch., Batkhuu, J., Boldsaikhan, B., and Komatsu, K. (2003). Illustrated Guide of Mongolian Useful Plants. (Vol. 1, p. 236). Ulaanbaatar: Admon Printing.
5. Ligaa, U., Davaasuren, B., and Ninjil, N. (2005). Medicinal Plants of Mongolia Used in Western and Eastern Medicine. (p. 391). Ulaanbaatar: JCK Printing.
6. Yuthok Yonten Gonpo., Four Medical Tantras, VIII-IXth century.
7. Boldsaikhan, B. (2004). Encyclopedia of Mongolian Medicinal Plants (p. 109). Ulaanbaatar: Mongolian University of Science and Technology.
8. Fedorov, A.A. *et al.* (1985). Plants Review of USSR: Family Magnoliaceae-Limoniaceae. (p. 138). Leningrad: Science Printing.