

**Distribution:** Khent., Khang. (east), Mong-Dag., Khyang., Dund. Khalkh, Dor. Mong., Dor. Gobi.

**Habitat:** Rocky and scree slopes, mountain foothills, ravines, valleys and steppes, filifolium and feather-grass steppes [2–4].

**Part used:** Herb

**Traditional Uses:** The taste is bitter and the potency is cool. It is used for the following: treating blood disorders. It is an ingredient of the following traditional prescriptions: Regonchitan, Utogtanmar, and Olsomei-25 [4–7].

**Chemical constituents:** 0.3% essential oil, phenol carboxylic acids: caffeic, sinapic, and ferulic acids, coumarin: ellagic acid, catechin: (–)-epicatechin, (–)-epigallocatechin, 4.9% tannins, flavonoids: quercetin, quercimetrin, astragalin, and kaempferol [8].

**Bioactivities:** Antihypertensive, spasmolytic, angioprotective, anti-inflammatory, and antibacterial [8].

#### References:

1. Olziikhutag, N. (Ed). (1983). Latin-Mongolian-Russian Dictionary of Vascular Plants of Mongolia (p. 163). UlaanbaaUlaanbaatar: Press of Mongolian Academy of Sciences.
2. Gubanov, I.A. (1996). Conspectus on Mongolian Flora (vascular plants) (p. 62). Moscow: Valang Press.
3. Malishev, L.I., and Peshkova, G.A. (1979). Flora of Central Siberia (Vol. 2, p. 572). Novosibirsk: Science Printing.
4. Ligaa, U., Davaasuren, B., and Ninjil, N. (2005). Medicinal Plants of Mongolia Used in Western and Eastern Medicine. (p. 129). Ulaanbaatar: JCK Printing.
5. Yuthok Yonten Gonpo., Four Medical Tantras, VIII-IXth century.
6. Danzanpuntsag., Crystal rosary. XVIIIth century.
7. Boldsaikhan, B. (2004). Encyclopedia of Mongolian Medicinal Plants (p. 118). Ulaanbaatar: Mongolian University of Science and Technology.
8. Sokolov, P.D. *et al.* (1987). Plants Review of USSR: Family Hydrangeaceae-Haloragaceae. (p. 59). Leningrad: Science Printing.