References:

- 1. Olziikhutag, N. (Ed). (1983). Latin-Mongolian-Russian Dictionary of Vascular Plants of Mongolia (p. 245). Ulaanbaatar: Press of Mongolian Academy of Sciences.
- 2. Gubanov, I.A. (1996). Conspectus on Mongolian Flora (vascular plants) (p. 94). Moscow: Valang Press.
- Malishev, L.I., and Peshkova, G.A. (1979). Flora of Central Siberia (Vol. 2, p. 799). Novosibirsk: Science Printing.
- 4. Sanchir, Ch., Batkhuu, J., Boldsaikhan, B., and Komatsu, K. (2003). Illustrated Guide of Mongolian Useful Plants. (Vol. 1, p. 239). Ulaanbaatar: Admon Printing.
- 5. Ligaa, U., Davaasuren, B., and Ninjil, N. (2005). Medicinal Plants of Mongolia Used in Western and Eastern Medicine. (p. 78). Ulaanbaatar: JCK Printing.
- 6. Yuthok Yonten Gonpo., Four Medical Tantras, VIII-IXth century.
- 7. Danzanpuntsag., Crystal rosary. XVIIIth century.
- 8. Boldsaikhan, B. (2004). Encyclopedia of Mongolian Medicinal Plants (p. 70). Ulaanbaatar: Mongolian University of Science and Technology.
- 9. Khurelchuluun, B., Suran, D., and Zina, C. (2007). Illustrated Guide of Raw Materials Used in Traditional Medicine. (p. 310). Ulaanbaatar: Erkhes Printing.
- 10. Enkhjargal, D., Bayasgalan, B., and Purevsuren, S. (2004). Pharmacognosy. (p. 124). Ulaanbaatar: Erkhes Printing.
- 11. Letchamo, W., Ward, W., Heard, B., and Heard, D. (2004). Essential oil of *Valeriana officinalis* L. cultivars and their antimicrobial activity as influenced by harvesting time under commercial organic cultivation. *J. Agric. Food Chem.* 52, 3915.
- 12. Hikino, H., Hikino, Y., Kato, H., Takeshita, Y., and Takemoto, T. (1969). Constituents of kesso (*Valeriana*) root. *Yakugaku Zasshi* 89, 118.
- 13. Sokolov, P.D. *et al.* (1990). Plants Review of USSR: Family Caprifoliaceae-Plantaginaceae. (p. 23). Leningrad: Science Printing.
- 14. Krepinsky, J., Herout, V., and Sorm, F. (1958). O rostlinnych lattkach: Islace neutral nich latek z korene kozliku lekarrskeho (*Valeriana officinalis* L.). *Chem. Lysty* 52, 1784.
- 15. Thies, P.W., and Funke, S. (1966). Über die Wirkstoffe der Baldrians. Tetrahedron Lett. 11, 1155.
- 16. Van Meer, J.H., Labadie, R.P. (1981). Straight-phase and reversed-phase high-performance liquid chromatographic separations of valepotriate isomers and homologues. *J. Chromatogr.* 205, 206.
- 17. Johnson, R.D. and Waller, G.R. (1971). Isolation of actinidine from Valeriana officinalis L. Phytochemistry 10, 3334.
- 18. Borkowski, F. (1960). Die Isolierung von Alkaloiden aus den Wurzelstöcken des Baldrians (*Valeriana officinalis* L.). *Pharmazie* 15, 568.
- 19. Franck, B., Peterson, U., and Hüper, F. (1970). Valerianine, a tertiary monoterpene alkaloid from *Valeriana*. *Angew. Chem*. 9, 891.
- 20. Casinovi, C.G. (1968). Composition of some Valeriana species. Fitoterapia 39, 45.
- 21. Dagvatseren, B., and Barkasdorj, Ts. (1990). Root and rhizome of Valeriana. Mongolian National Standard 924–90.