

# Chapter 5: Bayesian Inference for Poisson Distribution

## 강의 목표

- ▶ 푸아송분포를 중심으로 베이지안 추론의 이해.
- ▶ Parameter Estimation (모수 추정).
  - ▶ Point Estimation (점추정).
  - ▶ Confidence Interval (구간추정).
- ▶ Prediction (예측).

# Poisson Distribution

- ▶ Probability mass function for Poisson with rate  $\theta$ :

## Poisson Distribution

- ▶ Probability mass function for Poisson with rate  $\theta$ :

$$f(X = x \mid \theta) = \frac{\theta^x e^{-\theta}}{x!}.$$

- ▶ Suppose  $X_1, \dots, X_n$  have  $Poi(\theta)$ . Then, the joint p.d.f is

## Poisson Distribution

- Probability mass function for Poisson with rate  $\theta$ :

$$f(X = x \mid \theta) = \frac{\theta^x e^{-\theta}}{x!}.$$

- Suppose  $X_1, \dots, X_n$  have  $Poi(\theta)$ . Then, the joint p.d.f is

$$f(X_1, X_2, \dots, X_n \mid \theta) = \prod_{i=1}^n f(x_i \mid \theta) \propto \theta^{\sum x_i} e^{-n\theta}.$$

# How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto 1, \quad \theta > 0$$

## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto 1, \quad \theta > 0$$

- ▶ Posterior Dist:

$$P(\theta \mid x_1, \dots, x_n) \propto \theta^{\sum x_i} e^{-n\theta}, \quad \theta > 0$$

## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto 1, \quad \theta > 0$$

- ▶ Posterior Dist:

$$P(\theta \mid x_1, \dots, x_n) \propto \theta^{\sum x_i} e^{-n\theta}, \quad \theta > 0$$

- ▶ Gamma distribution with  $\sum x_i + 1$  and  $n$ .



# How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto \theta^a, \quad \theta > 0.$$

## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto \theta^a, \quad \theta > 0.$$

- ▶ Posterior Dist:

$$P(\theta \mid x_1, \dots, x_n) \propto \theta^{\sum x_i + a} e^{-n\theta}, \quad \theta > 0.$$

## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto \theta^a, \quad \theta > 0.$$

- ▶ Posterior Dist:

$$P(\theta \mid x_1, \dots, x_n) \propto \theta^{\sum x_i + a} e^{-n\theta}, \quad \theta > 0.$$

- ▶ It is Gamma Dist. with  $\alpha = \sum x_i + a + 1$  and  $\beta = n$ .

# How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto e^{-b\theta}, \quad \theta > 0.$$

## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto e^{-b\theta}, \quad \theta > 0.$$

- ▶ Posterior Dist:

$$P(\theta \mid x_1, \dots, x_n) \propto \theta^{\sum x_i} e^{-(n+b)\theta}, \quad \theta > 0.$$

## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto e^{-b\theta}, \quad \theta > 0.$$

- ▶ Posterior Dist:

$$P(\theta \mid x_1, \dots, x_n) \propto \theta^{\sum x_i} e^{-(n+b)\theta}, \quad \theta > 0.$$

- ▶ It is Gamma Dist. with  $\alpha = \sum x_i + 1$  and  $\beta = n + b$ .

## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto \theta^{a+1} e^{-b\theta}, \quad \theta > 0.$$

## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto \theta^{a+1} e^{-b\theta}, \quad \theta > 0.$$

- ▶ Posterior Dist:

$$P(\theta \mid x_1, \dots, x_n) \propto \theta^{\sum x_i + a + 1} e^{-(n+b)\theta}, \quad \theta > 0.$$



## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto \theta^{a+1} e^{-b\theta}, \quad \theta > 0.$$

- ▶ Posterior Dist:

$$P(\theta \mid x_1, \dots, x_n) \propto \theta^{\sum x_i + a + 1} e^{-(n+b)\theta}, \quad \theta > 0.$$

- ▶ It is Gamma Dist. with  $\alpha = \sum x_i + a$  and  $\beta = n + b$ .

## How to Choose Appropriate Prior

- ▶ Prior Dist:

$$P(\theta) \propto \theta^{a+1} e^{-b\theta}, \quad \theta > 0.$$

- ▶ Posterior Dist:

$$P(\theta \mid x_1, \dots, x_n) \propto \theta^{\sum x_i + a + 1} e^{-(n+b)\theta}, \quad \theta > 0.$$

- ▶ It is Gamma Dist. with  $\alpha = \sum x_i + a$  and  $\beta = n + b$ .
- ▶ If the prior distribution is gamma, the posterior is gamma.

# The Gamma / Poisson Bayesian Model

- ▶ Posterior Mean:

# The Gamma / Poisson Bayesian Model

- Posterior Mean:

$$\hat{\lambda} = \frac{\sum x_i + a}{n + b}.$$

# The Gamma / Poisson Bayesian Model

- ▶ Posterior Mean:

$$\hat{\lambda} = \frac{\sum x_i + a}{n + b}.$$

- ▶ It can be decomposed:

# The Gamma / Poisson Bayesian Model

- Posterior Mean:

$$\hat{\lambda} = \frac{\sum x_i + a}{n + b}.$$

- It can be decomposed:

$$\hat{\lambda} = \left( \frac{n}{n + b} \right) \left( \frac{\sum x_i}{n} \right) + \left( \frac{b}{n + b} \right) \left( \frac{a}{b} \right).$$

# The Gamma / Poisson Bayesian Model

- Posterior Mean:

$$\hat{\lambda} = \frac{\sum x_i + a}{n + b}.$$

- It can be decomposed:

$$\hat{\lambda} = \left( \frac{n}{n + b} \right) \left( \frac{\sum x_i}{n} \right) + \left( \frac{b}{n + b} \right) \left( \frac{a}{b} \right).$$

- The data get weighted more heavily as  $n \rightarrow \infty$ .

# Bayesian Learning

- ▶ We can use the Bayesian approach to update our information about the parameter(s) of interest **sequentially** as new data become available.



# Bayesian Learning

- ▶ We can use the Bayesian approach to update our information about the parameter(s) of interest **sequentially** as new data become available.
- ▶ Suppose we formulate a prior for our parameter  $\theta$  and observe a random sample  $x_1$ .

# Bayesian Learning

- ▶ We can use the Bayesian approach to update our information about the parameter(s) of interest **sequentially** as new data become available.
- ▶ Suppose we formulate a prior for our parameter  $\theta$  and observe a random sample  $x_1$ .
- ▶ The posterior is

$$\pi(\theta \mid x_1) \propto p(\theta)L(\theta \mid x_1).$$

# Bayesian Learning

- ▶ We can use the Bayesian approach to update our information about the parameter(s) of interest **sequentially** as new data become available.
- ▶ Suppose we formulate a prior for our parameter  $\theta$  and observe a random sample  $x_1$ .
- ▶ The posterior is

$$\pi(\theta \mid x_1) \propto p(\theta)L(\theta \mid x_1).$$

- ▶ Suppose that we observe a new (independent) sample  $x_2$ .

# Bayesian Learning

- ▶ We can use our previous posterior as the **new prior** and derive a **new posterior**:

$$\pi(\theta \mid x_1, x_2) \propto p(\theta)L(\theta \mid x_1, x_2)$$

# Bayesian Learning

- ▶ We can use our previous posterior as the **new prior** and derive a **new posterior**:

$$\begin{aligned}\pi(\theta \mid x_1, x_2) &\propto p(\theta)L(\theta \mid x_1, x_2) \\ &= p(\theta)L(\theta \mid x_1)L(\theta \mid x_2)\end{aligned}$$

# Bayesian Learning

- ▶ We can use our previous posterior as the **new prior** and derive a **new posterior**:

$$\begin{aligned}\pi(\theta \mid x_1, x_2) &\propto p(\theta)L(\theta \mid x_1, x_2) \\ &= p(\theta)L(\theta \mid x_1)L(\theta \mid x_2) \\ &= p(\theta \mid x_1)L(\theta \mid x_2).\end{aligned}$$

# Bayesian Learning

- ▶ We can use our previous posterior as the **new prior** and derive a **new posterior**:

$$\begin{aligned}\pi(\theta \mid x_1, x_2) &\propto p(\theta)L(\theta \mid x_1, x_2) \\ &= p(\theta)L(\theta \mid x_1)L(\theta \mid x_2) \\ &= p(\theta \mid x_1)L(\theta \mid x_2).\end{aligned}$$

- ▶ Note this is the same posterior we would have obtained had  $x_1$  and  $x_2$  arrived at the same time.

# Bayesian Learning

- ▶ We can use our previous posterior as the **new prior** and derive a **new posterior**:

$$\begin{aligned}\pi(\theta \mid x_1, x_2) &\propto p(\theta)L(\theta \mid x_1, x_2) \\ &= p(\theta)L(\theta \mid x_1)L(\theta \mid x_2) \\ &= p(\theta \mid x_1)L(\theta \mid x_2).\end{aligned}$$

- ▶ Note this is the same posterior we would have obtained had  $x_1$  and  $x_2$  arrived at the same time.
- ▶ This "**sequential updating**" process can continue indefinitely in the Bayesian setup.



## Traffic Example

두 도시 에서 차량 통행량 등 주위의 교통환경이 비슷한 교차로를 하나씩 선택하여 매주 발생한 교통사고 건수를 1년 동안 조사하였다. 첫 번째 도시 에서는 직진 후 좌회전 신호를 사용하고 두 번째 도시 에서는 좌회전 후 직진 신호를 사용한다. 교통사고 건수는 독립적으로 포아송 분포를따 른다고 가정한다. 교통통제 등의 이유로 조사를 할 수 없었던 기간을 제외하고 다음 표와 같은 조사 결과를 얻었다.

## Traffic Example

교통사고 건수	0	1	2	3	4	5	6	7
City 1의 사고 건수	7	14	13	8	4	2	2	0
City 2의 사고 건수	4	13	15	6	2	2	3	1

$$n_1 = 50, \quad \sum x_{1i} = 102, \quad \bar{x}_1 = 2.04$$

$$n_2 = 46, \quad \sum x_{2i} = 104, \quad \bar{x}_2 = 2.26$$

## Traffic Example

- ▶ 두 도시의 실제 평균 교통사고 건수  $\theta_1$ 과  $\theta_2$ 에 대하여  $\text{Gamma}(2, 1)$ 의 사전분포를 가정하자.
- ▶ 그렇다면 다음과 같은 posterior distribution을 찾을 수 있다.

$$\pi(\theta \mid n_1 = 50, \sum x_{1i} = 102) \sim \text{Gamma}(2 + 102, 1 + 50),$$


$$\pi(\theta \mid n_2 = 46, \sum x_{2i} = 104) \sim \text{Gamma}(2 + 104, 1 + 46).$$

## Traffic Example

```
x1 = rep(c(0, 1, 2, 3, 4, 5, 6), c(7, 14, 13, 8, 4, 2, 2))
x2 = rep(c(0, 1, 2, 3, 4, 5, 6, 7), c(4, 13, 15, 6, 2, 2, 3, 1))
a =2; b = 1
n1 = length(x1); s1 = sum(x1)
n2 = length(x2); s2 = sum(x2)
postmean.theta1 = (a+s1)/(b+n1)
postmean.theta2 = (a+s2)/(b+n2)
### plot the posterior
par(mfrow=c(1, 1))
theta <- seq(0, 6, length=100)
plot(theta, dgamma(theta, a+s1, b+n1), type="l", xlab="theta", ylab="p(theta|x)")
lines(theta, dgamma(theta, a+s2, b+n2), lty=2, col = "red")
lines(theta, dgamma(theta, a, b), lty=3, col = "blue")
legend ( 2.5, 1.5, legend=c (paste ("City 1"), paste("City 2"),
paste("Gamma(2, 1) prior")), cex = 1.3, lty=c(1, 2, 3), col=c(1, 2, 4),
      bty="n")
```

## Traffic Example

- ▶ City 1의 사고 발생 건수가 City 2에 비해 작다.
- ▶ 사후 분포의 분산이 사전 분포의 분산보다 작다.
- ▶ Likelihood의 영향으로 사후 분포들이 구간 (1.5, 3)이외에는 매우 비슷한다.



## Prediction Distribution

- Poisson - Gamma Prediction Dist:

$$f(x_{n+1} \mid x_1, \dots, x_n) = \int f(x_{n+1} \mid \theta, x_1, x_2, \dots, x_n) \pi(\theta \mid x_1, x_2, \dots, x_n)$$

## Prediction Distribution

- Poisson - Gamma Prediction Dist:

$$\begin{aligned} f(x_{n+1} \mid x_1, \dots, x_n) &= \int f(x_{n+1} \mid \theta, x_1, x_2, \dots, x_n) \pi(\theta \mid x_1, x_2, \dots, x_n) \\ &= \int f(x_{n+1} \mid \theta) \pi(\theta \mid x_1, x_2, \dots, x_n) d\theta \end{aligned}$$

## Prediction Distribution

► Poisson - Gamma Prediction Dist:

$$\begin{aligned}f(x_{n+1} \mid x_1, \dots, x_n) &= \int f(x_{n+1} \mid \theta, x_1, x_2, \dots, x_n) \pi(\theta \mid x_1, x_2, \dots, x_n) \\&= \int f(x_{n+1} \mid \theta) \pi(\theta \mid x_1, x_2, \dots, x_n) d\theta \\&= \int \frac{\theta^{x_{n+1}} e^{-\theta}}{x_{n+1}!} \times \frac{(b+n)^{a+\sum x_i}}{\Gamma(a+\sum x_i)} \theta^{a+\sum x_i-1} e^{-(b+n)\theta} d\theta\end{aligned}$$



## Prediction Distribution

► Poisson - Gamma Prediction Dist:

$$\begin{aligned}f(x_{n+1} \mid x_1, \dots, x_n) &= \int f(x_{n+1} \mid \theta, x_1, x_2, \dots, x_n) \pi(\theta \mid x_1, x_2, \dots, x_n) \\&= \int f(x_{n+1} \mid \theta) \pi(\theta \mid x_1, x_2, \dots, x_n) d\theta \\&= \int \frac{\theta^{x_{n+1}} e^{-\theta}}{x_{n+1}!} \times \frac{(b+n)^{a+\sum x_i}}{\Gamma(a+\sum x_i)} \theta^{a+\sum x_i-1} e^{-(b+n)\theta} d\theta \\&= \frac{(b+n)^{a+\sum x_i}}{x_{n+1}! \Gamma(a+\sum x_i)} \int \theta^{a+\sum x_i+x_{n+1}-1} e^{-(b+n+1)\theta} d\theta\end{aligned}$$

## Prediction Distribution

► Poisson - Gamma Prediction Dist:

$$\begin{aligned} f(x_{n+1} \mid x_1, \dots, x_n) &= \int f(x_{n+1} \mid \theta, x_1, x_2, \dots, x_n) \pi(\theta \mid x_1, x_2, \dots, x_n) \\ &= \int f(x_{n+1} \mid \theta) \pi(\theta \mid x_1, x_2, \dots, x_n) d\theta \\ &= \int \frac{\theta^{x_{n+1}} e^{-\theta}}{x_{n+1}!} \times \frac{(b+n)^{a+\sum x_i}}{\Gamma(a+\sum x_i)} \theta^{a+\sum x_i-1} e^{-(b+n)\theta} d\theta \\ &= \frac{(b+n)^{a+\sum x_i}}{x_{n+1}! \Gamma(a+\sum x_i)} \int \theta^{a+\sum x_i+x_{n+1}-1} e^{-(b+n+1)\theta} d\theta \\ &= \frac{(b+n)^{a+\sum x_i}}{x_{n+1}! \Gamma(a+\sum x_i)} \times \frac{\Gamma(a+\sum x_i+x_{n+1})}{(b+n+1)^{a+\sum x_i+x_{n+1}}} \end{aligned}$$

## Prediction Distribution

► Poisson - Gamma Prediction Dist:

$$\begin{aligned}f(x_{n+1} \mid x_1, \dots, x_n) &= \int f(x_{n+1} \mid \theta, x_1, x_2, \dots, x_n) \pi(\theta \mid x_1, x_2, \dots, x_n) \\&= \int f(x_{n+1} \mid \theta) \pi(\theta \mid x_1, x_2, \dots, x_n) d\theta \\&= \int \frac{\theta^{x_{n+1}} e^{-\theta}}{x_{n+1}!} \times \frac{(b+n)^{a+\sum x_i}}{\Gamma(a+\sum x_i)} \theta^{a+\sum x_i-1} e^{-(b+n)\theta} d\theta \\&= \frac{(b+n)^{a+\sum x_i}}{x_{n+1}! \Gamma(a+\sum x_i)} \int \theta^{a+\sum x_i+x_{n+1}-1} e^{-(b+n+1)\theta} d\theta \\&= \frac{(b+n)^{a+\sum x_i}}{x_{n+1}! \Gamma(a+\sum x_i)} \times \frac{\Gamma(a+\sum x_i+x_{n+1})}{(b+n+1)^{a+\sum x_i+x_{n+1}}} \\&\propto \frac{1}{x_{n+1}!} \times \frac{\Gamma(a+\sum x_i+x_{n+1})}{(b+n+1)^{a+\sum x_i+x_{n+1}}}\end{aligned}$$

## Prediction Distribution

$$f(x_{n+1} \mid x_1, \dots, x_n)$$

## Prediction Distribution

$$\begin{aligned} & f(x_{n+1} \mid x_1, \dots, x_n) \\ = & \binom{a + \sum x_i + x_{n+1} - 1}{a + \sum x_i - 1} \left( \frac{b + n}{b + n + 1} \right)^{a + \sum x_i} \left( \frac{1}{b + n + 1} \right)^{x_{n+1}} \\ = & \binom{a + \sum x_i + x_{n+1} - 1}{x_{n+1}} \left( \frac{1}{b + n + 1} \right)^{x_{n+1}} \left( \frac{b + n}{b + n + 1} \right)^{a + \sum x_i}. \end{aligned}$$

## Prediction Distribution

$$\Pr(X = x) = \binom{x + r - 1}{x} p^x (1 - p)^r \quad \text{for } x = 0, 1, 2, \dots$$

## Prediction Distribution

$$\Pr(X = x) = \binom{x+r-1}{x} p^x (1-p)^r \quad \text{for } x = 0, 1, 2, \dots$$

Hence, the prediction distribution is

## Prediction Distribution

$$\Pr(X = x) = \binom{x+r-1}{x} p^x (1-p)^r \quad \text{for } x = 0, 1, 2, \dots$$

Hence, the prediction distribution is

$$NB \left( a + \sum_{i=1}^n x_i, \frac{1}{b + n + 1} \right).$$



# Prediction Distribution

▶ 예측 기대치:

## Prediction Distribution

▶ 예측 기대치:

$$\mathbb{E}(X_{n+1} \mid x_1, x_2, \dots, x_n) = \frac{a + \sum x_i}{b + n}.$$

## Prediction Distribution

- ▶ 예측 기대치:

$$\mathbb{E}(X_{n+1} \mid x_1, x_2, \dots, x_n) = \frac{a + \sum x_i}{b + n}.$$

- ▶ 예측 기대치는 사후기대치와 동일.

## Prediction Distribution

- ▶ 예측 기대치:

$$\mathbb{E}(X_{n+1} \mid x_1, x_2, \dots, x_n) = \frac{a + \sum x_i}{b + n}.$$

- ▶ 예측 기대치는 사후기대치와 동일.
- ▶ 예측 분산:

## Prediction Distribution

- ▶ 예측 기대치:

$$\mathbb{E}(X_{n+1} \mid x_1, x_2, \dots, x_n) = \frac{a + \sum x_i}{b + n}.$$

- ▶ 예측 기대치는 사후기대치와 동일.
- ▶ 예측 분산:

$$\text{Var}(X_{n+1} \mid x_1, x_2, \dots, x_n) = \frac{a + \sum x_i}{(b + n)^2} (b + n + 1).$$

## Prediction Distribution

- ▶ 예측 기대치:

$$\mathbb{E}(X_{n+1} \mid x_1, x_2, \dots, x_n) = \frac{a + \sum x_i}{b + n}.$$

- ▶ 예측 기대치는 사후기대치와 동일.
- ▶ 예측 분산:

$$\text{Var}(X_{n+1} \mid x_1, x_2, \dots, x_n) = \frac{a + \sum x_i}{(b + n)^2} (b + n + 1).$$

- ▶ 예측분산은 사후분산보다  $(b + n + 1)$  곱한 만큼 크다.

## Prediction Expectation

- Expectation: using the fact  $\mathbb{E}(X_{n+1} \mid \theta) = \theta$ .

$$\mathbb{E}(X_{n+1} \mid x_1, \dots, x_n)$$

## Prediction Expectation

- Expectation: using the fact  $\mathbb{E}(X_{n+1} \mid \theta) = \theta$ .

$$\mathbb{E}(X_{n+1} \mid x_1, \dots, x_n) = \mathbb{E}(\mathbb{E}(X_{n+1} \mid \theta, x_1, \dots, x_n) \mid x_1, \dots, x_n)$$



## Prediction Expectation

- Expectation: using the fact  $\mathbb{E}(X_{n+1} \mid \theta) = \theta$ .

$$\begin{aligned}\mathbb{E}(X_{n+1} \mid x_1, \dots, x_n) &= \mathbb{E}(\mathbb{E}(X_{n+1} \mid \theta, x_1, \dots, x_n) \mid x_1, \dots, x_n) \\ &= \mathbb{E}(\theta \mid x_1, \dots, x_n).\end{aligned}$$

## Prediction Expectation

- Expectation: using the fact  $\mathbb{E}(X_{n+1} \mid \theta) = \theta$ .

$$\begin{aligned}\mathbb{E}(X_{n+1} \mid x_1, \dots, x_n) &= \mathbb{E}(\mathbb{E}(X_{n+1} \mid \theta, x_1, \dots, x_n) \mid x_1, \dots, x_n) \\ &= \mathbb{E}(\theta \mid x_1, \dots, x_n).\end{aligned}$$

- Hence, the expected prediction is the same as the posterior expectation.

## Prediction Variance

- ▶ Variance: using the fact  $\text{Var}(X_{n+1} \mid \theta) = \theta$ .

## Prediction Variance

- ▶ Variance: using the fact  $\text{Var}(X_{n+1} \mid \theta) = \theta$ .

$$\text{Var}(X_{n+1} \mid x_1, \dots, x_n)$$

## Prediction Variance

- Variance: using the fact  $\text{Var}(X_{n+1} \mid \theta) = \theta$ .

$$\begin{aligned}\text{Var}(X_{n+1} \mid x_1, \dots, x_n) &= \text{Var}(\mathbb{E}(X_{n+1} \mid \theta, x_1, \dots, x_n) \mid x_1, \dots, x_n) \\ &\quad + \mathbb{E}(\text{Var}(X_{n+1} \mid \theta, x_1, \dots, x_n) \mid x_1, \dots, x_n)\end{aligned}$$

## Prediction Variance

- Variance: using the fact  $\text{Var}(X_{n+1} \mid \theta) = \theta$ .

$$\begin{aligned}\text{Var}(X_{n+1} \mid x_1, \dots, x_n) &= \text{Var}(\mathbb{E}(X_{n+1} \mid \theta, x_1, \dots, x_n) \mid x_1, \dots, x_n) \\ &\quad + \mathbb{E}(\text{Var}(X_{n+1} \mid \theta, x_1, \dots, x_n) \mid x_1, \dots, x_n) \\ &= \text{Var}(\theta \mid x_1, \dots, x_n) + \mathbb{E}(\theta \mid x_1, \dots, x_n)\end{aligned}$$

## Prediction Variance

- Variance: using the fact  $\text{Var}(X_{n+1} \mid \theta) = \theta$ .

$$\begin{aligned}\text{Var}(X_{n+1} \mid x_1, \dots, x_n) &= \text{Var}(\mathbb{E}(X_{n+1} \mid \theta, x_1, \dots, x_n) \mid x_1, \dots, x_n) \\ &\quad + \mathbb{E}(\text{Var}(X_{n+1} \mid \theta, x_1, \dots, x_n) \mid x_1, \dots, x_n) \\ &= \text{Var}(\theta \mid x_1, \dots, x_n) + \mathbb{E}(\theta \mid x_1, \dots, x_n)\end{aligned}$$



- Hence, the variance of prediction distribution is larger than the the variance of the posterior expectation.

# Predictive Probability

```
> ## Ch 6
> #predictive distribution of  $X_{n+1}$ 
> x1=c(rep(0,7),rep(1,14),rep(2,13),rep(3,8),rep(4,4),rep(5,2),
+      rep(6,2))
> x2=c(rep(0,4),rep(1,13),rep(2,15),rep(3,6),rep(4,2),rep(5,2),
+      rep(6,3),rep(7,1) )
> a =2;b = 1
> n1 = length(x1); s1 = sum(x1)
> n2 = length(x2); s2 = sum(x2)
> x = seq(0,10)
> par(mfrow=c(1, 2))
> plot(x,dnbinom(x,size=a+s1,prob=(b+n1)/(b+n1+1)), xlab="x_{n+1}",
+      ylab="Predictive Prob", type="h", main="City 1")
> plot(x,dnbinom(x,size=a+s2,prob=(b+n2)/(b+n2+1)), xlab="x_{n+1}",
+      ylab="Predictive Prob" ,type="h" , main="City 2" )
```



# Predictive Probability



## Monte Carlo Method for Traffic Example

- ▶ 두 도시의 실제 평균 교통사고 건수  $\theta_1$ 과  $\theta_2$ 에 대하여 다음과 같은 posterior distribution을 찾았다.

$$\pi(\theta \mid n_1 = 50, \sum x_{1i} = 102) \sim \text{Gamma}(2 + 102, 1 + 50),$$

$$\pi(\theta \mid n_2 = 46, \sum x_{2i} = 104) \sim \text{Gamma}(2 + 104, 1 + 46).$$

## Monte Carlo Method for Traffic Example

- ▶ 두 도시의 실제 평균 교통사고 건수  $\theta_1$ 과  $\theta_2$ 에 대하여 다음과 같은 posterior distribution을 찾았다.

$$\pi(\theta \mid n_1 = 50, \sum x_{1i} = 102) \sim \text{Gamma}(2 + 102, 1 + 50),$$

$$\pi(\theta \mid n_2 = 46, \sum x_{2i} = 104) \sim \text{Gamma}(2 + 104, 1 + 46).$$

- ▶ Gamma 분포는 이미 알려져 있지만, 관련 statistics는 여전히 찾기 어렵다.

## Monte Carlo Method for Traffic Example

- ▶ 이 예제에서 주요 목적은  $\theta_1$ 과  $\theta_2$  차이가 얼마인지 찾는 데에 있다. (i.e.,  $\theta_1 - \theta_2$ ).

## Monte Carlo Method for Traffic Example

- ▶ 이 예제에서 주요 목적은  $\theta_1$ 과  $\theta_2$  차이가 얼마인지 찾는 데에 있다. (i.e.,  $\theta_1 - \theta_2$ ).
  - ▶ Posterior expectation for  $\theta_1 - \theta_2$  given data.
  - ▶ Posterior variance for  $\theta_1 - \theta_2$  given data.

## Monte Carlo Method for Traffic Example

- ▶ 이 예제에서 주요 목적은  $\theta_1$ 과  $\theta_2$  차이가 얼마인지 찾는 데에 있다. (i.e.,  $\theta_1 - \theta_2$ ).
  - ▶ Posterior expectation for  $\theta_1 - \theta_2$  given data.
  - ▶ Posterior variance for  $\theta_1 - \theta_2$  given data.
- ▶ Monte Carlo Method를 통해 두 parameter들의 차이에 관련된 통계량을 찾을 수 있다

## Monte Carlo Method for Traffic Example

- For notational convenience, let  $\eta = \theta_1 - \theta_2$ .

```
a =2 ; b = 1
n1 = 50; s1 = 102 ; n2 = 46; s2 = 104 ;
nsim = 30000
theta1.sim = rgamma(nsim,a+s1,b+n1)
theta2.sim = rgamma(nsim,a+s2,b+n2)
eta=theta1.sim- theta2.sim
mean(eta)
[1] -0.2155787
var( eta)
[1] 0.08880491
```


## Monte Carlo Method for Traffic Example

- ▶ 이론  $\mathbb{E}(\eta \mid \text{data}) = -0.2161$ .
- ▶ 이론  $(\eta \mid \text{data}) = 0.0875$ .



# Monte Carlo Method for Traffic Example

```
HPD=HPDsample(eta)
par(mfrow=c(1,1))
plot(density(eta), type="l", xlab= "theta1-theta2",
ylab="posterior density", main="")
abline( v= HPD, lty=2)
text(mean(eta),0.3, "95% HPD interval" )
```



## Monte Carlo Method for Prediction

- ▶ 예측 분포  $X_{n+1} \mid X_1, \dots, X_n$ 은 Posterior 분포보다 더욱 복잡한 형태를 가진 경우가 많다.

## Monte Carlo Method for Prediction

- ▶ 예측 분포  $X_{n+1} \mid X_1, \dots, X_n$ 은 Posterior 분포보다 더욱 복잡한 형태를 가진 경우가 많다.
- ▶ Monte Carlo Method는  $\mathbb{E}(X_{n+1} \mid X_1, \dots, X_n)$ 과  $\text{Var}(X_{n+1} \mid X_1, \dots, X_n)$ 를 추정하는데 큰 도움을 준다.

## Monte Carlo Method for Prediction

- ▶ 예측 분포  $X_{n+1} \mid X_1, \dots, X_n$ 은 Posterior 분포보다 더욱 복잡한 형태를 가진 경우가 많다.
- ▶ Monte Carlo Method는  $\mathbb{E}(X_{n+1} \mid X_1, \dots, X_n)$ 과  $\text{Var}(X_{n+1} \mid X_1, \dots, X_n)$ 를 추정하는데 큰 도움을 준다.
- ▶ Recall that

$$f(X_{n+1} \mid X_1, \dots, X_n) = \mathbb{E}(f(X_{n+1} \mid \theta) \mid X_1, \dots, X_n).$$

## First Method

1. N개의  $\theta_i$ 를 랜덤하게 생성.

$$\theta_i \sim \pi(\theta \mid x_1, \dots, x_n).$$

## First Method

1. N개의  $\theta_i$ 를 랜덤하게 생성.

$$\theta_i \sim \pi(\theta \mid x_1, \dots, x_n).$$

2. 각  $\theta_i$ 에 대하여  $x_{n+1}$ 를 생성함.

$$x_{n+1,i} \sim f(x_{n+1} \mid \theta_i).$$

## First Method

1.  $N$ 개의  $\theta_i$ 를 랜덤하게 생성.

$$\theta_i \sim \pi(\theta \mid x_1, \dots, x_n).$$

2. 각  $\theta_i$ 에 대하여  $X_{n+1}$ 를 생성함.

$$x_{n+1,i} \sim f(x_{n+1} \mid \theta_i).$$

### ► Estimation

- i.  $\widehat{\mathbb{E}}(X_{n+1} \mid X_1, \dots, X_n) = \frac{1}{N} \sum_{i=1}^N x_{n+1,i}$
- ii.  $\widehat{\text{Var}}(X_{n+1} \mid X_1, \dots, X_n) = \frac{1}{N} \sum_{i=1}^N x_{n+1,i}^2 - \left( \frac{1}{N} \sum_{i=1}^N x_{n+1,i} \right)^2$

## Second Method: Rao-Blackwellization

1. N개의  $\theta_i$ 를 랜덤하게 생성.

$$\theta_i \sim \pi(\theta \mid x_1, \dots, x_n).$$



## Second Method: Rao-Blackwellization

1.  $N$ 개의  $\theta_i$ 를 랜덤하게 생성.

$$\theta_i \sim \pi(\theta \mid x_1, \dots, x_n).$$

2.  $\theta_i$ 를 통해 prediction mass function 을 예측.

$$\hat{f}(x_{n+1} \mid x_1, \dots, x_n) = \frac{1}{N} \sum_{i=1}^N f(x_{n+1} \mid \theta).$$

## Second Method: Rao-Blackwellization

1.  $N$ 개의  $\theta_i$ 를 랜덤하게 생성.

$$\theta_i \sim \pi(\theta \mid x_1, \dots, x_n).$$

2.  $\theta_i$ 를 통해 prediction mass function 을 예측.

$$\hat{f}(x_{n+1} \mid x_1, \dots, x_n) = \frac{1}{N} \sum_{i=1}^N f(x_{n+1} \mid \theta).$$

### ► Estimation

- i.  $\hat{\mathbb{E}}(X_{n+1} \mid X_1, \dots, X_n) = \sum_{\text{all } x_{n+1}} x_{n+1} \hat{f}(x_{n+1} \mid x_1, \dots, x_n)$
- ii.  $\text{Var}(X_{n+1} \mid X_1, \dots, X_n) = \sum_{\text{all } x_{n+1}} x_{n+1}^2 \hat{f}(x_{n+1} \mid x_1, \dots, x_n) - \left( \sum_{\text{all } x_{n+1}} x_{n+1} \hat{f}(x_{n+1} \mid x_1, \dots, x_n) \right)^2.$

## Comparisons

- ▶ Rao-Blackwellization는 probability mass function 을 추측. 또 그 확률을 통해 mean과 variance를 예측.
- ▶ 첫번째 방법이 더 복잡하나, 더 정확한 경향이 있다.