

Geometric stacks in Synthetic Algebraic Geometry

Tim Lichtnau, Hugo Moeneclay

Oktober 2024

Abstract

This is meant as a short summary of the progress of the Master Thesis of Tim Lichtnau so far. We work in Synthetic Algebraic Geometry, i.e. Homotopy Type Theory + 3 Axioms. A type gets interpreted as a Zariski-sheaf on the site given by the opposite category of finitely presented algebras over a fixed ring [CCH23]. Affine types get interpreted as the representable sheaves.

Definition 1.1. A Grothendieck topology \mathbb{T} is a subclass of affine schemes, such that

- $1 \in \mathbb{T}$
- \mathbb{T} is \sum -stable, i.e. if $X : \mathbb{T}, B : X \rightarrow \mathbb{T}$, then $\sum_{x:X} Bx$ belongs to \mathbb{T} .

A map is a \mathbb{T} -cover iff its fibered in \mathbb{T} . Relating to the classical definition, we could call a family of maps of affines $\{U_i \rightarrow U\}_{i=1}^n$ covering iff

$$\sum_{i=1}^n U_i \rightarrow U$$

is a \mathbb{T} -cover.

We fix a topology \mathbb{T} , for which we want to define the notion of (geometric) stack.

Definition 1.2. A type X is a (higher) stack iff its $\|\text{Spec } A\|$ -local for any $\text{Spec } A \in \mathbb{T}$. An n -stack is a stack that is an n -type.

This exactly captures the expected classical notion of a stack for a Grothendieck topology [Moe24b], e.g. a 0-type X is a stack, if for any \mathbb{T} -cover $A \rightarrow B$,

$$X^B \rightarrow X^A \rightrightarrows X^{A \times_B A}$$

is an equalizer diagram.

Example 1.3 ([Moe24a]).

- The fppf topology is given by the faithfully flat affine schemes.
- The étale topology is given by formally-étale + faithfully flat affine schemes.
- The smooth topology is given by smooth + faithfully flat affine schemes.

We start by defining the relative setting (this corresponds to fibers of smooth morphism of geometric stacks in [Sim96]). There is a short inductive definition:

Definition 1.4. A stack X is covering, whenever inductively

- $X \in \mathbb{T}$ or
- X is equipped with a map $\mathbb{T} \ni \text{Spec } A \rightarrow X$ fibered in covering stacks.

We call a map $X \rightarrow Y$ fibered in covering stacks a geometric cover (That corresponds to smooth morphisms in [Sim96])

Definition 1.5. A stack X is geometric iff it merely admits a geometric cover $\text{Spec } A \rightarrow X$.

We have the following classical labels associated to geometric stacks depending on the topologies:

	\mathbb{T}	Geometric stacks for \mathbb{T}
gies:	étale	(Higher) Deligne Mumford Stacks
	smooth	(Higher) Artin stacks
	fppf	Something similar to Artin stacks ?

Example 1.6. *Every stack that is a scheme is geometric.*

Theorem 1.7 (Stability Results).

- The class of covering / geometric stacks is Σ -stable.
- The class of covering / geometric stacks is closed under quotients: If $X \rightarrow Y$ is a geometric cover with X covering / geometric stack, then Y is covering / geometric
- Geometric stacks are closed under taking identity types.
- Every geometric stack is a geometric n -stack for some n
- Covering / Geometric stacks have descent: Both types `GeometricStack` and `CoveringStack` are a stack.

It is maybe worth mentioning, that proving descent was surprisingly easy.

Under some very mild condition on the topology ¹ (e.g. satisfied by fppf), which is equivalent to saying that every geometric cover between affines is a \mathbb{T} -cover we have the following explicit description depending on the truncation level n :

Theorem 1.8. *An n -stack X is geometric if and only if*

($n = 0$): there merely exists a map $\text{Spec } A \rightarrow X$ whose fibers F merely admit a \mathbb{T} -cover $\mathbb{T} \ni \text{Spec } B \rightarrow F$. ²

($n \geq 1$): there merely exist a map $\text{Spec } A \rightarrow X$ whose fibers are covering $(n - 1)$ -stacks. Additionally X is covering iff we can choose $\text{Spec } A$ to lie in \mathbb{T} .

For the étale topology we have the following notable results

Theorem 1.9.

- Every Deligne-Mumford stack is a 1-gerbe, i.e. $X \rightarrow \|X\|_1^{\mathbb{T}}$ is a geometric cover, where the latter means the \mathbb{T} -sheafification of the 1-truncation of X .
- A Deligne-Mumford stack X is covering iff $\pi_0^{\mathbb{T}} X := \|X\|_0^{\mathbb{T}}$ and all higher homotopy groups

$$\pi_i^{\mathbb{T}}(X, x) = \|\Omega^i(X, x)\|_0^{\mathbb{T}}, i \geq 1$$

are covering algebraic spaces for the étale topology.

¹We can always enforce this condition without changing the notion of (covering / geometric) stack

²One can reformulate also as taking a quotient of $\text{Spec } A$ by an equivalence relation satisfying a certain property.

References

- [CCH23] Felix Cherubini, Thierry Coquand, and Matthias Hutzler. *A Foundation for Synthetic Algebraic Geometry*. 2023. arXiv: [2307.00073](https://arxiv.org/abs/2307.00073) [math.AG]. URL: <https://arxiv.org/abs/2307.00073>.
- [Moe24a] Hugo Moeneclay. *Synthetic Differential Geometry*. HoTTUF 2024. 2024. URL: https://hott-uf.github.io/2024/abstracts/HoTTUF_2024_paper_11.pdf.
- [Moe24b] Moeneclay, Hugo. “Sheaves”. In: (2024). URL: www.felix-cherubini.de/sheaves.pdf.
- [Sim96] Carlos Simpson. *Algebraic (geometric) n -stacks*. 1996. arXiv: [alg-geom/9609014](https://arxiv.org/abs/alg-geom/9609014) [alg-geom]. URL: <https://arxiv.org/abs/alg-geom/9609014>.