

A
OBi1
DSS−3
Homarine ($C_7H_7NO_3$)
N−methylglutamic Acid ($C_6H_{11}NO_4$)
N−methylglutamine ($C_6H_{12}N_2O_3$)
Glutamic Acid ($C_5H_9NO_4$)
$C_6H_9NO_3$ (−@rt10)
$C_6H_9NO_3$ (−@rt11)
$C_6H_9NO_4$ (−@rt3.5)
$C_6H_9NO_4$ (+@rt12)
$C_7H_9NO_5$ (+@rt12)
Core metabs (max)
Core metabs (med)
Core metabs (min)
homarine
glucose + homarine
enrichment factor (log2)
−3 0 5 10 15
B
OBi1
DSS−3
N−methylglutamic acid
8 9 10 11 12
C
OBi1
DSS−3
N−methylglutamine
7 8 9 10 11 12
Retention Time (min)
glucose
homarine
D
50°N
25°N
0°
5°S
160°W
130°W
100°W
RC104 St2
TN397 St2
TN412 St2
TN412 St7
TN397 St13
E
Homarine
N−methylglutamic Acid
N−methylglutamine
F
TN397 St2
TN397 St13
$^2H_3$ isotopologues
Homarine
N−methylglutamic Acid
N−methylglutamine
$C_6H_9NO_3$ (−@rt10)
$C_6H_9NO_3$ (−@rt11)
$C_6H_9NO_4$ (−@rt3.5)
$C_7H_9NO_5$ (+@rt12)
Core metabs (max)
Core metabs (med)
Core metabs (min)
2hr 24hr 48hr
2hr 24hr 48hr
G
$^{13}C_{5-7}$, ^{15}N isotopologues
C104 St
N412 St
N412 St
Homarine
N−methylglutamic Acid
N−methylglutamine
Glutamic Acid
$C_6H_9NO_3$ (−@rt10)
$C_6H_9NO_3$ (−@rt11)
$C_6H_9NO_4$ (+@rt12)
$C_7H_9NO_5$ (+@rt12)
Core metabs (max)
Core metabs (med)
Core metabs (min)