

FIT2004

Algorithms and Data Structures

Ian Wern Han Lim
lim.wern.han@monash.edu

Referencing materials by
Rafael Dowsley, Nathan Companeze, Aamir Cheema, Arun Konagurthu and Lloyd Allison



Faculty of Information Technology, Monash University

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

This material has been reproduced and communicated to you by or on behalf of Monash University pursuant to Part VB of the Copyright Act 1968 (the Act). The material in this communication may be subject to copyright under the Act. Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act. Do not remove this notice

Ready?

Agenda

- Grade School Integer Multiplication
- Quick Integer Multiplication
 - Karatsuba algorithm (1960, 1962)
 - Schönhage–Strassen algorithm (1971)
 - Popular for matrix multiplication
 - ... and many more including a 2019 one in UNSW

Agenda

- Grade School Integer Multiplication
- Quick Integer Multiplication
 - Karatsuba algorithm (1960, 1962)
 - Schönhage–Strassen algorithm (1971)
 - Popular for matrix multiplication
 - ... and many more including a 2019 one in UNSW
- Divide and Conquer
 - Simple recap on MergeSort
 - Simple recap on QuickSort

Let us begin!

Grade School Multiplication

An algorithm we all know

- Recall how you multiply from grade school

Grade School Multiplication

An algorithm we all know

- Recall how you multiply from grade school
 - It is an algorithm
 - It is something that you do use till today, right?

Grade School Multiplication

An algorithm we all know

- Recall how you multiply from grade school
 - It is an algorithm
 - It is something that you do use till today, right?

- Consider the following 2 inputs
 - $x = 123$
 - $y = 345$
 - Multiple x with y , what is the answer?

Grade School Multiplication

An algorithm we all know

- Recall how you multiply from grade school
 - It is an algorithm
 - It is something that you do use till today, right?
 - What is your algorithm?
 1. Loop through y from right to left.
 2. For each integer of y , multiply with each integer in x from right to left.
 3. If overflow exist, add to the left value

- Consider the following 2 inputs
 - $x = 123$
 - $y = 345$
 - Multiple x with y , what is the answer?

Grade School Multiplication

An algorithm we all know

- Recall how you multiply from grade school

- It is an algorithm
- It is something that you do use till today, right?
- What is your algorithm?

1. Loop through y from right to left.
2. For each integer of y , multiply with each integer in x from right to left.
3. If overflow exist, add to the left value
4. ...

You can write it in a better way, and code it

- Consider the following 2 inputs

- $x = 123$
- $y = 345$
- Multiple x with y , what is the answer?

Grade School Multiplication

An algorithm we all know

- Consider the following 2 inputs
 - $x = 123$
 - $y = 345$
 - Multiple x with y , what is the answer?
- Have your teacher ever...

$$\begin{array}{r} 1 2 3 \\ \times 3 4 5 \\ \hline 6 1 5 \\ 4 9 2 \\ 3 6 9 \\ \hline 4 2 4 3 5 \end{array}$$

Grade School Multiplication

An algorithm we all know

- Consider the following 2 inputs
 - $x = 123$
 - $y = 345$
 - Multiple x with y , what is the answer?
- Have your teacher ever...
 - Explain why it work?

$$\begin{array}{r} 1 2 3 \\ \times 3 4 5 \\ \hline 6 1 5 \\ 4 9 2 \\ 3 6 9 \\ \hline 4 2 4 3 5 \end{array}$$

Grade School Multiplication

An algorithm we all know

- Consider the following 2 inputs
 - $x = 123$
 - $y = 345$
 - Multiple x with y , what is the answer?
- Have your teacher ever...
 - Explain why it work? For every possible number combinations?

$$\begin{array}{r} 1 2 3 \\ \times 3 4 5 \\ \hline 6 1 5 \\ 4 9 2 \\ 3 6 9 \\ \hline 4 2 4 3 5 \end{array}$$

Grade School Multiplication

An algorithm we all know

- Consider the following 2 inputs
 - $x = 123$
 - $y = 345$
 - Multiple x with y , what is the answer?
- Have your teacher ever...
 - Explain why it work? For every possible number combinations?
 - How efficient it is?



$$\begin{array}{r} 1 2 3 \\ \times 3 4 5 \\ \hline 6 1 5 \\ 4 9 2 \\ 3 6 9 \\ \hline 4 2 4 3 5 \end{array}$$

Grade School Multiplication

An algorithm we all know


- Consider the following 2 inputs
 - $x = 123$
 - $y = 345$
 - Multiple x with y , what is the answer?
- Have your teacher ever...
 - Explain why it work? For every possible number combinations?
 - How efficient it is? **Probably not...** that is why....

$$\begin{array}{r} 1 2 3 \\ \times 3 4 5 \\ \hline 6 1 5 \\ 4 9 2 \\ 3 6 9 \\ \hline 4 2 4 3 5 \end{array}$$



Grade School Multiplication

An algorithm we all know



Tootie Fruity 5 months ago

I'm Asian and years ago I showed my Asian parents the box method and they started laughing hysterically calling it "ridiculous". lol

23 1 REPLY

View 3 replies



Skinny Bo8000 1 year ago

This teacher is trying to score some extra working hours LMAO

69 1 REPLY

View 2 replies



Cadetsc 4 months ago

If the problem was 635×232 .

Would had taken her a decade to solve

12 1 REPLY



V Mano 3 months ago

I'm Filipino. I saw how my American 2nd grade niece was trying to grasp the American way of addition and subtraction. She wanted to cry. I taught her the Asian way of doing it. She was able to add and subtract in the millions. Why are the kids being treated as dumb? 29-16. You have to add 1 on both numbers, then subtract 17 from 30. What????

4 1 REPLY

Grade School Multiplication

An algorithm we all know

- Consider the following 2 inputs
 - $x = 123$
 - $y = 345$
 - Multiple x with y , what is the answer?

$$\begin{array}{r} 1 2 3 \\ \times 3 4 5 \\ \hline 6 1 5 \\ 4 9 2 \\ 3 6 9 \\ \hline 4 2 4 3 5 \end{array}$$

- Have your teacher ever...
 - Explain why it work? For every possible number combinations?
 - How efficient it is?
 - This is not efficient...
 - As the numbers become bigger you need more steps!

Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ \hline 42435 \end{array}$$

← How many operations to produce this?

Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ \hline 42435 \end{array}$$

← How many operations to produce this?
Multiply 5×3
Note the overflow
Multiply 5×2
Add the overflow (from prev)
Note the overflow
Multiply 5×1
Add the overflow (from prev)

Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ \hline 42435 \end{array}$$

← Now repeat for this...

Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ \hline 42435 \end{array}$$


← And this this...

Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ \hline 42435 \end{array}$$

Then add them all up



Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ \hline 42435 \end{array}$$

It is a lot of operations!

... and it become worse as the number becomes bigger!

Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ \hline 42435 \end{array}$$

It is a lot of operations!

... and it become worse as the number becomes bigger!

Each digit in x , increases actions per row

Each digit in y , increases a row!

Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ \hline 42435 \end{array}$$

It is a lot of operations!

... and it become worse as the number becomes bigger!

Each digit in x , increases actions per row

Each digit in y , increases a row!

And in the end, you need to add them all up!!!

Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ 42435 \end{array}$$

And don't forget, you need to store all of these somewhere...
What if they are really big? You are wasting memory....

Grade School Multiplication

An algorithm we all know

$$\begin{array}{r} \times \quad 123 \\ 345 \\ \hline 615 \\ 492 \\ 369 \\ \hline 42435 \end{array}$$

This entire thing?

- 9 multiplications
- 3 noting of overflow
- 3 addition for overflow
- The final addition process that deals with a lot of integers and overflow

Grade School Multiplication

An algorithm we all know

- That is why students are taught the box method... which provides the foundation for Karatsuba!

Questions?

Karatsuba

Simple Quick Integer Multiplication

- A method to multiply 2 integers
- Able to do so quickly, even as the integers scales to be bigger integers

Simple Quick Integer Multiplication

- A method to multiply 2 integers
- Able to do so quickly, even as the integers scales to be bigger integers
- By breaking large numbers into smaller ones
 - $x = 1234$
 - $y = 6789$
 - $x * y = ?$

- A method to multiply 2 integers
- Able to do so quickly, even as the integers scales to be bigger integers
- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = ?$

Simple Quick Integer Multiplication

- A method to multiply 2 integers
- Able to do so quickly, even as the integers scales to be bigger integers
- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$

Simple Quick Integer Multiplication

- A method to multiply 2 integers
- Able to do so quickly, even as the integers scales to be bigger integers
- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $+ (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$

Simple Quick Integer Multiplication

- A method to multiply 2 integers
- Able to do so quickly, even as the integers scales to be bigger integers
- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $+ (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$
 $= (12 * 67 * 10^4) + (12 * 89 * 10^2)$
 $+ (34 * 67 * 10^2) + (34 * 89)$

- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $+ (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$
 $= (12 * 67 * 10^4) + (12 * 89 * 10^2)$
 $+ (34 * 67 * 10^2) + (34 * 89)$

- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $+ (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$
 $= (12 * 67 * 10^4) + (12 * 89 * 10^2)$
 $+ (34 * 67 * 10^2) + (34 * 89)$
 - Can you see it?

- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $+ (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$
 $= (12 * 67 * 10^4) + (12 * 89 * 10^2)$
 $+ (34 * 67 * 10^2) + (34 * 89)$
 - Can you see it?

- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $+ (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$
 $= (12 * 67 * 10^4) + (12 * 89 * 10^2)$
 $+ (34 * 67 * 10^2) + (34 * 89)$
- So why do this matter?

- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $+ (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$
 $= (12 * 67 * 10^4) + (12 * 89 * 10^2)$
 $+ (34 * 67 * 10^2) + (34 * 89)$
- So why do this matter?
 - Quicker to multiply small integers
 - Quicker to multiple simple numbers
 - Example: $123 * 100 = 12300$

- By breaking large numbers into smaller ones
 - $x = 1234 = 12 * 10^2 + 34 * 10^0$
 - $y = 6789 = 67 * 10^2 + 89 * 10^0$
 - $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $+ (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$
 $= (12 * 67 * 10^4) + (12 * 89 * 10^2)$
 $+ (34 * 67 * 10^2) + (34 * 89)$
- So why do this matter? If you count the **operations**???
 - Quicker to multiply small integers
 - Quicker to multiple simple numbers
 - Example: $123 * 100 = 12300$

Simple Quick Integer Multiplication

- By breaking large numbers into smaller ones

- $x = 1234 = 12 * 10^2 + 34 * 10^0$

- $y = 6789 = 67 * 10^2 + 89 * 10^0$

- $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $\quad + (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$
 $= (12 * 67 * 10^4) + (12 * 89 * 10^2)$
 $\quad + (34 * 67 * 10^2) + (34 * 89)$
 $= (12 * 67 * 10^4) + (12 * 89 + 34 * 67) * 10^2 + (34 * 89)$



- And we can do even better!
 - Add small numbers, then only multiply!

- With that, we can generalize:
 - Given integer x with n -digits
 - Given integer y with n -digits

- With that, we can generalize:
 - Given integer x with n -digits
 - Given integer y with n -digits
 - x can be broken down into
 - The most significant half x_l
 - The less significant half x_r

12345678
1234 | 5678
↑ ↑
 x_l x_r

- With that, we can generalize:
 - Given integer x with n -digits
 - Given integer y with n -digits
 - x can be broken down into
 - The most significant half x_l
 - The less significant half x_r

123456789
01234 | 56789
↑ ↑
 x_l x_r

- With that, we can generalize:
 - Given integer x with n -digits
 - Given integer y with n -digits
 - x can be broken down into
 - The most significant half x_l
 - The less significant half x_r
 - y can be broken down the same
- put 0 at the front to make all even digit number into odd digit number

- With that, we can generalize:
 - Given integer x with n -digits
 - Given integer y with n -digits
 - x can be broken down into
 - The most significant half x_l
 - The less significant half x_r
 - y can be broken down the same
 - Therefore $x * y = x_l * y_l * 10^n + (x_l * y_r + x_r * y_l) * 10^{n/2} + x_r * y_r$

- With that, we can generalize:
 - Given integer x with n -digits
 - Given integer y with n -digits
 - x can be broken down into
 - The most significant half x_l
 - The less significant half x_r
 - y can be broken down the same
 - Therefore $x * y = \underbrace{x_l * y_l}_{* 10^n} + (\underbrace{x_l * y_r}_{* 10^{n/2}} + \underbrace{x_r * y_l}_{* 10^{n/2}}) + \underbrace{x_r * y_r}_{* 10^{n/2}}$

Questions?

- By breaking large numbers into smaller ones


- $x = 1234 = 12 * 10^2 + 34 * 10^0$

- $y = 6789 = 67 * 10^2 + 89 * 10^0$

- $x * y = (12 * 10^2 + 34 * 10^0) * (67 * 10^2 + 89 * 10^0)$
 $= (12 * 10^2 * 67 * 10^2) + (12 * 10^2 * 89 * 10^0)$
 $+ (34 * 10^0 * 67 * 10^2) + (34 * 10^0 * 89 * 10^0)$
 $= (12 * 67 * 10^4) + (12 * 89 * 10^2)$
 $+ (34 * 67 * 10^2) + (34 * 89)$
 $= (12 * 67 * 10^4) + (12 * 89 + 34 * 67) * 10^2 + (34 * 89)$

- And we can do even better!

- Add small numbers, then only multiply!
 - What if I tell you **we can do even better?**




- Recall we stopped at the following

$$- \text{ Therefore } x * y = \underbrace{x_l * y_l}_{\text{}} * 10^n + (\underbrace{x_l * y_r}_{\text{}} + \underbrace{x_r * y_l}_{\text{}}) * 10^{n/2} + \underbrace{x_r * y_r}_{\text{}}$$

- Recall we stopped at the following

$$- \text{ Therefore } x * y = \underbrace{x_l * y_l}_{\text{red brace}} * 10^n + (\underbrace{x_l * y_r}_{\text{red brace}} + \underbrace{x_r * y_l}_{\text{red brace}}) * 10^{n/2} + \underbrace{x_r * y_r}_{\text{red brace}}$$



- Recall we stopped at the following
 - Therefore $x * y = x_l * y_l * 10^n + \underbrace{(x_l * y_r + x_r * y_l)} * 10^{n/2} + x_r * y_r$
 - Gauss introduce a trick for us

Simple Quick Integer Multiplication

- Recall we stopped at the following
 - Therefore $x * y = x_l * y_l * 10^n + \underbrace{(x_l * y_r + x_r * y_l)} * 10^{n/2} + x_r * y_r$
 - Gauss introduce a trick for us
 - $(x_l + x_r) * (y_l + y_r) = x_l * y_l + x_l * y_r + x_r * y_l + x_r * y_r$

- Recall we stopped at the following
 - Therefore $x * y = x_l * y_l * 10^n + \underbrace{(x_l * y_r + x_r * y_l)} * 10^{n/2} + x_r * y_r$
 - Gauss introduce a trick for us
 - $(x_l + x_r) * (y_l + y_r) = x_l * y_l + x_l * y_r + x_r * y_l + x_r * y_r$
 - Then we rearrange the above...
 - $\underbrace{x_l * y_r + x_r * y_l} = (x_l + x_r) * (y_l + y_r) - x_l * y_l - x_r * y_r$

- Recall we stopped at the following
 - Therefore $x * y = x_l * y_l * 10^n + \underbrace{(x_l * y_r + x_r * y_l)} * 10^{n/2} + x_r * y_r$
 - Gauss introduce a trick for us
 - $(x_l + x_r) * (y_l + y_r) = x_l * y_l + x_l * y_r + x_r * y_l + x_r * y_r$
 - Then we rearrange the above...
 - $\underbrace{x_l * y_r + x_r * y_l} = (x_l + x_r) * (y_l + y_r) - x_l * y_l - x_r * y_r$
 - Why are we doing this?

- Recall we stopped at the following
 - Therefore $x * y = x_l * y_l * 10^n + \underbrace{(x_l * y_r + x_r * y_l)} * 10^{n/2} + x_r * y_r$
 - Gauss introduce a trick for us
 - $(x_l + x_r) * (y_l + y_r) = x_l * y_l + x_l * y_r + x_r * y_l + x_r * y_r$
 - Then we rearrange the above...
 - $\underbrace{x_l * y_r + x_r * y_l} = (x_l + x_r) * (y_l + y_r) - x_l * y_l - x_r * y_r$
 - Why are we doing this?

- Recall we stopped at the following

- Therefore $x * y = x_l * y_l * 10^n + \underbrace{(x_l * y_r + x_r * y_l)} * 10^{n/2} + x_r * y_r$

- Gauss introduce a trick for us

- $(x_l + x_r) * (y_l + y_r) = x_l * y_l + x_l * y_r + x_r * y_l + x_r * y_r$

- Then we rearrange the above...

- $\underbrace{x_l * y_r + x_r * y_l} = (x_l + x_r) * (y_l + y_r) - x_l * y_l - x_r * y_r$

- Why are we doing this?

- Recall we stopped at the following

- Therefore $x * y = x_l * y_l * 10^n + \underbrace{(x_l * y_r + x_r * y_l)}_{\text{cross terms}} * 10^{n/2} + x_r * y_r$

- Gauss introduce a trick for us

- $(x_l + x_r) * (y_l + y_r) = x_l * y_l + x_l * y_r + x_r * y_l + x_r * y_r$

- Then we rearrange the above...

- $x_l * y_r + x_r * y_l = \underbrace{(x_l + x_r) * (y_l + y_r) - x_l * y_l - x_r * y_r}_{\text{cross terms}}$

Sub this back upper red underlined part to have a quicker computation

- Why are we doing this?

- 1 multiplication instead of 2 multiplication

- Note that it is slower to multiply than it is to add/ subtract in general

Questions?

- Given 2 large numbers
- Divide and conquer the large number into 2 halves
 - Smaller numbers are faster to operate on
 - Only need 3 multiplications, on smaller numbers
- Then combine the result

- Given 2 large numbers
- Divide and conquer the large number into 2 halves
 - Smaller numbers are faster to operate on
 - Only need 3 multiplications, on smaller numbers




We can follow Karatsuba again
for the 3 multiplications!

- Then combine the result

Karatsuba

In summary

- To multiply 2 large numbers of n -digits, Karatsuba can do so in $O(N^{1.59})$, which is much more scalable than $O(N^2)$



Questions?

Divide and Conquer

A Recap

- Take a problem
- **Divide** the problem into smaller subproblems
- **Conquer** the smaller subproblems, getting the solution
- Combine the smaller solutions to obtain the bigger solution

Divide and Conquer

A Recap

- Take a problem
- **Divide** the problem into smaller subproblems
- **Conquer** the smaller subproblems, getting the solution
- Combine the smaller solutions to obtain the bigger solution
- And you done that with:
 - Karatsuba earlier
 - MergeSort and QuickSort from earlier your prerequisite(s)

Divide and Conquer

A Recap

- Take a problem
- Divide the problem into smaller subproblems
 - Karatsuba: Split a number into most-significant digits (MSD) and least-significant digits (LSD)
 - MergeSort and QuickSort: Split a list into left-partition and right-partition
- Conquer the smaller subproblems, getting the solution
- Combine the smaller solutions to obtain the bigger solution

Divide and Conquer

A Recap

- Take a problem
- Divide the problem into smaller subproblems
- Conquer the smaller subproblems, getting the solution
 - Karatsuba: Multiply the smaller digits.
 - MergeSort and QuickSort: Sort the partitions
- Combine the smaller solutions to obtain the bigger solution

Divide and Conquer

A Recap

- Take a problem
- Divide the problem into smaller subproblems
- Conquer the smaller subproblems, getting the solution
- Combine the smaller solutions to obtain the bigger solution
 - Karatsuba: Add and subtract the values together.
 - MergeSort and QuickSort: Combine the partitions in sorted order.

Divide and Conquer

A Recap

- Take a problem
 - Divide the problem into smaller subproblems
 - Conquer the smaller subproblems, getting the solution
 - Combine the smaller solutions to obtain the bigger solution
-
- You would notice that many of them are done in recursively as well; we will explore how to analyze **recursive complexity** in a later lecture.

Questions?

Divide and Conquer

Other DnC Algorithms?

- Finding **closest pair** of **points** in a plane in $O(n \log n)$.
- Counting inversions in $O(n \log n)$, see you Studio question.
- Improving matrix multiplication (Strassen's algorithm).
- Fast Fourier Transform: this algorithm published by James Cooley and John Tukey in 1965 is one of the most influential algorithms, with a wide range of applications in engineering, music, science, mathematics, etc.
 - In fact, it can be traced back to unpublished work by Gauss.

Questions?

Thank You