The Simply Typed Lambda Calculus and Variants Thereof

Sandro Stucki

University of Gothenburg | Chalmers, Sweden

Virtual ITC Lecture 1 - 5 Nov 2020

sandro.stucki@gu.se @stuckintheory





Part I STLC

The plain, old simply typed lambda calculus

Part I STLC

The plain, old simply typed lambda calculus

Part II Explicit substitutions

Adding explicit substitutions and their equational theory

Part I STLC

The plain, old simply typed lambda calculus

Part II Explicit substitutions

Adding explicit substitutions and their equational theory

Part III De Bruijn indices

Doing away with variables...

Part I STLC

The plain, old simply typed lambda calculus

Part II Explicit substitutions

Adding explicit substitutions and their equational theory

Part III De Bruijn indices

Doing away with variables...

Part IV Correspondences

Everything is connected.



Syntax

\mathcal{V}	\ni	x, y, z, \dots	(Variable
\mathcal{T}	\ni	$s, t, u := x \mid \lambda x. t \mid s t$	(Term)
\mathcal{A}	\ni	$A, B, C ::= b \mid A \rightarrow B$	(Type)
\mathcal{C}	\ni	$\Gamma, \Delta, E ::= \cdot \mid \Gamma, x : A$	(Context)

Term typing

$$\Gamma \vdash t : A$$

$$x_n: A_n, \dots, x_0: A_0 \vdash x_i: A_i$$
 (VAR)

$$\frac{\Gamma, x \colon A \vdash t \colon B}{\Gamma \vdash \lambda x \colon t \colon A \to B} \tag{ABS}$$

$$\frac{\Gamma \vdash t : A \to B \quad \Gamma \vdash u : A}{\Gamma \vdash t u : B} \tag{APP}$$

Substitutions

Substitutions are finite maps $\sigma \colon \mathcal{V} \to \mathcal{T}$ from variables to terms.

Application of substitutions

$$x[\sigma] := \sigma(x)$$
$$(\lambda x. t)[\sigma] := \lambda x. t[\sigma\{x := x\}]$$
$$(t u)[\sigma] := t[\sigma] u[\sigma]$$

Equational theory

$$\Gamma \vdash t = u : A$$

$$\Gamma \vdash t = t : A \text{ (EREFL)}$$

$$\frac{\Gamma \vdash t_1 = t_2 : A}{\Gamma \vdash t_2 = t_1 : A} \text{ (ESYM)}$$

$$\frac{\Gamma \vdash t_1 = t_2 : A \quad \Gamma \vdash t_2 = t_3 : A}{\Gamma \vdash t_1 = t_3 : A}$$
 (ETRANS)

$$\frac{\Gamma, x \colon A \vdash t \colon B \quad \Gamma \vdash u \colon A}{\Gamma \vdash (\lambda x \colon t) \, u = t[x \coloneqq u] \colon B}$$
(E- β)

$$\frac{\Gamma \vdash t : A \to B \quad x \notin \mathsf{fv}(t)}{\Gamma \vdash t = \lambda x. \, t \, x : A \to B} \, (\mathsf{E} - \eta)$$

$$\frac{\Gamma, x \colon A \vdash t = t' \colon B}{\Gamma \vdash \lambda x \colon t = \lambda x \colon t' \colon A \to B}$$
(EABS)

$$\frac{\Gamma \vdash t = t' : A \to B \quad \Gamma \vdash u = u' : A}{\Gamma \vdash t \, u = t' \, u' : B}$$
(EAPP)

Explicit Substitutions

Substitutions are finite maps $\sigma \colon \mathcal{V} \to \mathcal{T}$ from variables to terms.

But so far, they were meta-theoretic.

Explicit Substitutions

Substitutions are finite maps $\sigma \colon \mathcal{V} \to \mathcal{T}$ from variables to terms.

But so far, they were meta-theoretic.

Let's make them syntactic!

Explicit Substitutions

Substitutions are finite maps $\sigma \colon \mathcal{V} \to \mathcal{T}$ from variables to terms.

But so far, they were meta-theoretic.

Let's make them syntactic!

$$S \quad \ni \quad \rho, \sigma, \tau ::= \langle \rangle \mid \langle \sigma, t \rangle \mid \uparrow \mid \operatorname{id} \mid \rho \circ \sigma \qquad \text{(Substitution)}$$

$$\Gamma \vdash t : A$$

$$\frac{\Gamma \vdash \sigma : \Delta \quad \Delta \vdash t : A}{\Gamma \vdash t[\sigma] : A} \tag{SUB}$$

Substitution typing

$$\Gamma \vdash \sigma : \Delta$$

$$\Gamma \vdash \langle \rangle : \cdot \qquad \qquad \text{(EMPTY)}$$

$$\frac{\Gamma \vdash \sigma : \Delta \quad \Gamma \vdash t : A}{\Gamma \vdash \langle \sigma, x := t \rangle : \Delta, x : A} \qquad \qquad \text{(PAIR)}$$

$$\Gamma, x : A \vdash \uparrow : \Gamma \qquad \qquad \text{(WEAKEN)}$$

$$\Gamma \vdash \text{id} : \Gamma \qquad \qquad \text{(ID)}$$

$$\frac{\Gamma \vdash \sigma : \Delta \quad \Delta \vdash \rho : E}{\Gamma \vdash \rho \circ \sigma : E} \qquad \qquad \text{(COMP)}$$

$$\Gamma \vdash t = u : A$$

$$\frac{\Gamma, x \colon A \vdash t \colon B \quad \Gamma \vdash u \colon A}{\Gamma \vdash (\lambda x \colon t) \ u = t[\langle \mathbf{id}, x \coloneqq u \rangle] \colon B}$$
 (E-\beta)

Update: term equality

$$|\Gamma \vdash t = u : A|$$

$$\frac{\Gamma, x \colon A \vdash t \colon B \quad \Gamma \vdash u \colon A}{\Gamma \vdash (\lambda x \colon t) \ u = t[\langle \mathbf{id}, x := u \rangle] \colon B}$$
 (E-\beta)

New rules:

$$\frac{\Gamma \vdash \sigma : \Delta \quad \Gamma \vdash t : A}{\Gamma \vdash x[\langle \sigma, x := t \rangle] = t : A} \frac{\Gamma \vdash \sigma : \Delta \quad \Delta \vdash t : A \to B \quad \Delta \vdash u : A}{\Gamma \vdash (t \, u)[\sigma] = t[\sigma] \, u[\sigma] : B}$$
(EAPPS)

$$\frac{\Gamma \vdash \sigma : \Delta \quad \Delta, x : A \vdash t : B}{\Gamma \vdash (\lambda x. t)[\sigma] = \lambda x. t[\langle \sigma \circ \uparrow, x := x \rangle] : A \to B}$$
 (EABSS)

Typing the RHS of β -contraction

$$(\mathsf{SUB}) \ \frac{\Gamma, x \colon A \vdash t \colon B}{\Gamma \vdash \mathsf{t}[\langle \mathsf{id}, x \coloneqq u \rangle] \colon A} \frac{(\mathsf{ID}) \ \frac{\Gamma \vdash \mathsf{id} \colon \Gamma}{\Gamma \vdash \mathsf{id} \colon \Gamma} \quad \Gamma \vdash u \colon A}{\Gamma \vdash \langle \mathsf{id}, x \coloneqq u \rangle \colon \Delta, x \colon A}$$

Substitution equality (1/2)

$$\Gamma \vdash \sigma = \rho : \Delta$$

$$\frac{\Gamma \vdash \rho : \Delta \quad \Delta \vdash \sigma : E \quad E \vdash \tau : \Phi}{\Gamma \vdash (\tau \circ \sigma) \circ \rho = \tau \circ (\sigma \circ \rho) : \Phi} \tag{SEASSOC}$$

$$\frac{\Gamma \vdash \sigma : \Delta}{\Gamma \vdash \operatorname{id} \circ \sigma = \sigma : \Delta} \qquad \frac{\Gamma \vdash \sigma : \Delta}{\Gamma \vdash \sigma \circ \operatorname{id} = \sigma : \Delta} \qquad \Gamma \vdash \sigma = \sigma : \Delta$$
 (SEREFL)

$$\begin{array}{ll} \Gamma \vdash \sigma_1 = \sigma_2 : \Delta \\ \hline \Gamma \vdash \sigma_2 = \sigma_1 : \Delta \\ \text{(SESYM)} \end{array} \qquad \begin{array}{ll} \Gamma \vdash \sigma_1 = \sigma_2 : \Delta & \Gamma \vdash \sigma_2 = \sigma_3 : \Delta \\ \hline \Gamma \vdash \sigma_1 = \sigma_3 : \Delta \\ \end{array} \qquad \qquad \text{(SETRANS)}$$

$$\frac{\Gamma \vdash \rho : \Delta \quad \Delta \vdash \sigma : E \quad \Delta \vdash t : A}{\Gamma \vdash \langle \sigma, x := t \rangle \circ \rho = \langle \sigma \circ \rho, x := t[\rho] \rangle : E, x : A} \text{ (SEPAIRS)}$$

Substitution equality (2/2)

$$\Gamma \vdash \sigma = \rho : \Delta$$

$$\frac{\Gamma \vdash \sigma : \Delta \quad \Gamma \vdash t : A}{\Gamma \vdash \uparrow \circ \langle \sigma, x := t \rangle = \sigma : \Delta}$$
 (SE- β)

$$\Gamma, x : A \vdash id = \langle \uparrow, x := x \rangle : \Gamma, x : A$$
 (SE- η)

$$\frac{\Gamma \vdash \sigma : \cdot \quad \Gamma \vdash \sigma' : \cdot}{\Gamma \vdash \sigma = \sigma' : \cdot}$$
 (SEEMPTY)

$$\frac{\Gamma \vdash \rho = \rho' : \Delta \quad \Delta \vdash \sigma = \sigma' : E}{\Gamma \vdash \sigma \circ \rho = \sigma' \circ \rho' : E} \tag{SECOMP}$$

$$\frac{\Gamma \vdash \sigma = \sigma' : \Delta \quad \Gamma \vdash t = t' : A}{\Gamma \vdash \langle \sigma, x := t \rangle = \langle \sigma, x := t' \rangle : \Delta, x : A}$$
 (SEPAIR)

Getting rid of variables: de Bruijn indices

De Bruijn indices •000

```
(Variable)
       \ni 0, 1, 2, ...
\mathcal{T} \ni s, t, u := i \mid \lambda t \mid st \mid t[\sigma]
                                                                                                  (Term)
A \ni A, B, C ::= b \mid A \rightarrow B
                                                                                                   (Type)
     \ni \Gamma, \Delta, E ::= \cdot \mid \Gamma, A
                                                                                             (Context)
    \Rightarrow \rho, \sigma, \tau ::= \langle \rangle \mid \langle \sigma, t \rangle \mid \uparrow \mid id \mid \rho \circ \sigma
                                                                                     (Substitution)
```

Typing with de Bruijn indices

$$A_n, \dots, A_0 \vdash i : A_i$$
 (VAR)

$$\frac{\Gamma, A \vdash t : B}{\Gamma \vdash \lambda t : A \to B} \tag{ABS}$$

$$\frac{\Gamma \vdash \sigma : \Delta \quad \Gamma \vdash t : A}{\Gamma \vdash \langle \sigma, t \rangle : \Delta, A}$$
 (PAIR)

$$\frac{\Gamma, A \vdash t : B \quad \Gamma \vdash u : A}{\Gamma \vdash (\lambda t) \ u = t[\langle \mathrm{id}, u \rangle] : B} \tag{E-β}$$

Observation: we can express natural numbers in Peano-style.

```
1 = suc(0)
2 = suc(suc(0))
3 = suc(suc(suc(0)))
\vdots
```

Getting rid of indices: use just 0 + weakening

De Bruijn indices 0000

Observation: we can express natural numbers in Peano-style.

$$1 = 0[\uparrow]$$

$$2 = 0[\uparrow][\uparrow]$$

$$3 = 0[\uparrow][\uparrow][\uparrow]$$

$$\vdots$$

Getting rid of indices: use just 0 + weakening

De Bruijn indices 0000

Observation: we can express natural numbers in Peano-style.

```
1 = 0 \uparrow
2 = 0 \uparrow \uparrow \uparrow \uparrow \uparrow
3 = 0 \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow
```

```
(Variable)
     \ni s, t, u := 0 | \lambda t | s t | t[\sigma]
                                                                                                 (Term)
A \ni A, B, C ::= b \mid A \rightarrow B
                                                                                                  (Type)
     \ni \Gamma, \Delta, E := \cdot \mid \Gamma, A
                                                                                             (Context)
       \Rightarrow \rho, \sigma, \tau ::= \langle \rangle \mid \langle \sigma, t \rangle \mid \uparrow \mid id \mid \rho \circ \sigma
                                                                                     (Substitution)
```

Typing with de Bruijn indices

$$\Gamma, A \vdash 0 : A$$
 (VAR)

$$\frac{\Gamma, A \vdash t : B}{\Gamma \vdash \lambda t : A \to B} \tag{Abs}$$

$$\frac{\Gamma \vdash \sigma : \Delta \quad \Gamma \vdash t : A}{\Gamma \vdash \langle \sigma, t \rangle : \Delta, A}$$
 (PAIR)

$$\frac{\Gamma, A \vdash t : B \quad \Gamma \vdash u : A}{\Gamma \vdash (\lambda t) \ u = t[\langle \mathrm{id}, u \rangle] : B} \tag{E-β}$$

Getting rid of contexts: products

$$\begin{array}{lll} \mathcal{V} & \ni & 0 & \text{(Variable)} \\ \mathcal{T} & \ni & s,t,u ::= 0 \mid \lambda t \mid st \mid t[\sigma] & \text{(Term)} \\ \mathcal{A} & \ni & A,B,C ::= b \mid A \to B & \text{(Type)} \\ \mathcal{C} & \ni & \Gamma,\Delta,E ::= \cdot \mid \Gamma,A & \text{(Context)} \\ \mathcal{S} & \ni & \rho,\sigma,\tau ::= \langle \rangle \mid \langle \sigma,t \rangle \mid \uparrow \mid \text{id} \mid \rho \circ \sigma & \text{(Substitution)} \end{array}$$

$$\begin{array}{lll} \mathcal{V} & \ni & \pi_2 & \text{(Variable)} \\ \mathcal{T} & \ni & s,t,u ::= \pi_2 \mid \lambda t \mid st \mid t[\sigma] & \text{(Term)} \\ \mathcal{A} & \ni & A,B,C ::= \mathbf{b} \mid A \to B & \text{(Type)} \\ \mathcal{C} & \ni & \Gamma,\Delta,E ::= \mathbf{1} \mid \Gamma \times A & \text{(Context)} \\ \mathcal{S} & \ni & \rho,\sigma,\tau ::= \langle \rangle \mid \langle \sigma,t \rangle \mid \pi_1 \mid \operatorname{id} \mid \rho \circ \sigma & \text{(Substitution)} \end{array}$$

Getting rid of contexts: products

Getting rid of contexts: products

000 0000000 __

Term typing

$$\Gamma \vdash t : A$$

Term typing

$$\Gamma \vdash t : A$$

Combinator typing

$$t\colon \Gamma \longrightarrow A$$

$$\begin{array}{c} \pi_2\colon \Gamma\times A\longrightarrow A \quad \text{(VAR)} \quad \frac{\sigma\colon \Gamma\longrightarrow \Delta \quad t\colon \Gamma\longrightarrow A}{\langle \sigma,t\rangle\colon \Gamma\longrightarrow \Delta\times A} \text{ (PAIR)} \\ \frac{t\colon \Gamma\times A\longrightarrow B}{\lambda t\colon \Gamma\longrightarrow A\to B} \quad \text{(ABS)} \\ \frac{t\colon \Gamma\to A\to B \quad u\colon \Gamma\longrightarrow A}{tu\colon \Gamma\longrightarrow B} \quad \frac{\rho\colon \Gamma\longrightarrow \Delta \quad \sigma\colon \Delta\longrightarrow E}{\sigma\circ \rho\colon \Gamma\longrightarrow E} \\ \text{(COMP)} \\ \langle \rangle\colon \Gamma\longrightarrow \mathbf{1} \quad \text{(EMPTY)} \qquad \text{id}\colon \Gamma\longrightarrow \Gamma \qquad \text{(ID)} \end{array}$$

Combinatory IPL

$$\Gamma \Longrightarrow A$$

$$\Gamma \times A \Longrightarrow A$$
 (VAR)

$$\frac{\Gamma \times A \Longrightarrow B}{\Gamma \Longrightarrow A \supset B} \qquad \text{(ABS)}$$

$$\frac{\Gamma \Longrightarrow A \supset B \quad \Gamma \Longrightarrow A}{\Gamma \Longrightarrow B} \text{ (APP)}$$


$$\Gamma \Longrightarrow \top$$
 (EMPTY)

$$\frac{\Gamma \Longrightarrow \Delta \quad \Gamma \Longrightarrow A}{\Gamma \Longrightarrow \Delta \times A} \text{ (PAIR)}$$

$$\Gamma imes A \Longrightarrow \Gamma$$
 (Weaken)

$$\frac{\Gamma \Longrightarrow \Delta \quad \Delta \Longrightarrow E}{\Gamma \Longrightarrow E}$$
 (COMP)

$$\Gamma \Longrightarrow \Gamma$$
 (ID)



Except where otherwise noted, this work is licensed under

http://creativecommons.org/licenses/by/3.0/