

# Takeuti's conjecture

## And Prawitz's proof

Jelle Tjeerd Fokkens

Initial Types Club

April 8, 2021

# Outline

Step 1

Interlude

Step 2

Interlude 2

Step 3

Summary



# Introduction



- ▶ Gerhard Gentzen, 1934 proved the Hauptsatz for first order sequent calculus.

# Introduction



- ▶ Gerhard Gentzen, 1934 proved the Hauptsatz for first order sequent calculus.
- ▶ Gaisi Takeuti stated the conjecture for STT in 1953.

# Introduction



- ▶ Gerhard Gentzen, 1934 proved the Hauptsatz for first order sequent calculus.
- ▶ Gaisi Takeuti stated the conjecture for STT in 1953.
- ▶ Proved by Tait for second-order logic in 1966.

# Introduction



- ▶ Gerhard Gentzen, 1934 proved the Hauptsatz for first order sequent calculus.
- ▶ Gaisi Takeuti stated the conjecture for STT in 1953.
- ▶ Proved by Tait for second-order logic in 1966.
- ▶ Proved independently by Motoo Takahashi and Dag Prawitz in 1967.

# The challenge

- ▶ Russell's paradox: 'The class of all classes that do not contain themselves as elements.'

# The challenge

- ▶ Russell's paradox: 'The class of all classes that do not contain themselves as elements.'
- ▶ Supposed underlying problem: self-reference.



# The challenge

- ▶ Russell's paradox: 'The class of all classes that do not contain themselves as elements.'
- ▶ Supposed underlying problem: self-reference.
- ▶ Impredicativity is supposed to be bad.

$$\frac{\Gamma \vdash \forall X A(X)}{\Gamma \vdash A(X := T)} , \text{ where } T \text{ is any term.}$$

# The challenge

- ▶ Russell's paradox: 'The class of all classes that do not contain themselves as elements.'
- ▶ Supposed underlying problem: self-reference.
- ▶ Impredicativity is supposed to be bad.

$$\frac{\Gamma \vdash \forall X A(X)}{\Gamma \vdash A(X := T)} , \text{ where } T \text{ is any term.}$$

- ▶ Furthermore, extensionality:

$$\forall x (P(x) \leftrightarrow Q(x)) \rightarrow (RP \rightarrow RQ)$$

# Proof outline

We're proving three claims which together imply the Hauptsatz.

1. if  $A$  is not derivable without cut then there is a semi-valuation with  $A$  false

# Proof outline

We're proving three claims which together imply the Hauptsatz.

1. if  $A$  is not derivable without cut then there is a semi-valuation with  $A$  false
2. semi-valuations are extendable to total valuations

# Proof outline

We're proving three claims which together imply the Hauptsatz.

1. if  $A$  is not derivable without cut then there is a semi-valuation with  $A$  false
2. semi-valuations are extendable to total valuations
3. if  $A$  is false in a total valuation, then  $A$  is not derivable

So, if  $A$  is derivable, it is derivable without cut.

# STT

- ▶ Types:
  - ▶ 0 (individuals) and 1 (truth values) are types

# STT

- ▶ Types:
  - ▶ 0 (individuals) and 1 (truth values) are types
  - ▶ if  $\tau_1, \dots, \tau_n$  are types, then  $(\tau_1, \dots, \tau_n)$  is a type (predicate with indicated argument places)

# STT

- ▶ Types:
  - ▶ 0 (individuals) and 1 (truth values) are types
  - ▶ if  $\tau_1, \dots, \tau_n$  are types, then  $(\tau_1, \dots, \tau_n)$  is a type (predicate with indicated argument places)
- ▶ Basic symbols:
  - ▶ free  $(a_i^\tau)$  and bound  $(x_i^\tau)$  variables of any type.



# STT

- ▶ Types:
  - ▶ 0 (individuals) and 1 (truth values) are types
  - ▶ if  $\tau_1, \dots, \tau_n$  are types, then  $(\tau_1, \dots, \tau_n)$  is a type (predicate with indicated argument places)
- ▶ Basic symbols:
  - ▶ free ( $a_i^\tau$ ) and bound ( $x_i^\tau$ ) variables of any type.
  - ▶ constants ( $c_i^\tau$ ) of any type.

- ▶ Types:
  - ▶ 0 (individuals) and 1 (truth values) are types
  - ▶ if  $\tau_1, \dots, \tau_n$  are types, then  $(\tau_1, \dots, \tau_n)$  is a type (predicate with indicated argument places)
- ▶ Basic symbols:
  - ▶ free ( $a_i^\tau$ ) and bound ( $x_i^\tau$ ) variables of any type.
  - ▶ constants ( $c_i^\tau$ ) of any type.
  - ▶ function symbols ( $f_i$ ) denoting functions with arguments and values of type 0.

- ▶ Types:
  - ▶ 0 (individuals) and 1 (truth values) are types
  - ▶ if  $\tau_1, \dots, \tau_n$  are types, then  $(\tau_1, \dots, \tau_n)$  is a type (predicate with indicated argument places)
- ▶ Basic symbols:
  - ▶ free ( $a_i^\tau$ ) and bound ( $x_i^\tau$ ) variables of any type.
  - ▶ constants ( $c_i^\tau$ ) of any type.
  - ▶ function symbols ( $f_i$ ) denoting functions with arguments and values of type 0.
  - ▶  $\neg$  for negation,  $\vee$  for disjunction,  $\exists$  for existence,  $\lambda$  for abstraction and  $\in$  for membership.

# STT, continued

Expressions:

## STT, continued

Expressions:

- ▶ free variables and constants of type  $\tau$  is an expression of type  $\tau$

# STT, continued

Expressions:

- ▶ free variables and constants of type  $\tau$  is an expression of type  $\tau$
- ▶ if  $\phi$  is a function symbol with  $n$  argument places and  $e_1, \dots, e_n$  are expressions of type 0, then  $\phi(e_1, \dots, e_n)$  is an expression of type 0

# STT, continued

Expressions:

- ▶ free variables and constants of type  $\tau$  is an expression of type  $\tau$
- ▶ if  $\phi$  is a function symbol with  $n$  argument places and  $e_1, \dots, e_n$  are expressions of type 0, then  $\phi(e_1, \dots, e_n)$  is an expression of type 0
- ▶ if  $e_1, \dots, e_n$  are expressions of types  $\tau_1, \dots, \tau_n$ , and  $e$  is an expression of type  $(\tau_1, \dots, \tau_n)$ , then  $(e_1, \dots, e_n \in e)$  is an expression of type 1

# STT, continued

Expressions:

- ▶ free variables and constants of type  $\tau$  is an expression of type  $\tau$
- ▶ if  $\phi$  is a function symbol with  $n$  argument places and  $e_1, \dots, e_n$  are expressions of type 0, then  $\phi(e_1, \dots, e_n)$  is an expression of type 0
- ▶ if  $e_1, \dots, e_n$  are expressions of types  $\tau_1, \dots, \tau_n$ , and  $e$  is an expression of type  $(\tau_1, \dots, \tau_n)$ , then  $(e_1, \dots, e_n \in e)$  is an expression of type 1
- ▶ if  $A$  is an expression of type 1, then  $\neg A$  is an expression of type 1



# STT, continued

Expressions:

- ▶ free variables and constants of type  $\tau$  is an expression of type  $\tau$
- ▶ if  $\phi$  is a function symbol with  $n$  argument places and  $e_1, \dots, e_n$  are expressions of type 0, then  $\phi(e_1, \dots, e_n)$  is an expression of type 0
- ▶ if  $e_1, \dots, e_n$  are expressions of types  $\tau_1, \dots, \tau_n$ , and  $e$  is an expression of type  $(\tau_1, \dots, \tau_n)$ , then  $(e_1, \dots, e_n \in e)$  is an expression of type 1
- ▶ if  $A$  is an expression of type 1, then  $\neg A$  is an expression of type 1
- ▶ if  $A$  and  $B$  are expressions of type 1, then  $(A \vee B)$  is an expression of type 1

# STT, continued

Expressions:

- ▶ free variables and constants of type  $\tau$  is an expression of type  $\tau$
- ▶ if  $\phi$  is a function symbol with  $n$  argument places and  $e_1, \dots, e_n$  are expressions of type 0, then  $\phi(e_1, \dots, e_n)$  is an expression of type 0
- ▶ if  $e_1, \dots, e_n$  are expressions of types  $\tau_1, \dots, \tau_n$ , and  $e$  is an expression of type  $(\tau_1, \dots, \tau_n)$ , then  $(e_1, \dots, e_n \in e)$  is an expression of type 1
- ▶ if  $A$  is an expression of type 1, then  $\neg A$  is an expression of type 1
- ▶ if  $A$  and  $B$  are expressions of type 1, then  $(A \vee B)$  is an expression of type 1
- ▶ if  $x^\tau$  is a bound variable which does not occur in an expression  $A(a^\tau)$  of type 1, then  $\exists x^\tau A(x^\tau)$  is an expression of type 1.

# STT, continued

## Expressions:

- ▶ free variables and constants of type  $\tau$  is an expression of type  $\tau$
- ▶ if  $\phi$  is a function symbol with  $n$  argument places and  $e_1, \dots, e_n$  are expressions of type 0, then  $\phi(e_1, \dots, e_n)$  is an expression of type 0
- ▶ if  $e_1, \dots, e_n$  are expressions of types  $\tau_1, \dots, \tau_n$ , and  $e$  is an expression of type  $(\tau_1, \dots, \tau_n)$ , then  $(e_1, \dots, e_n \in e)$  is an expression of type 1
- ▶ if  $A$  is an expression of type 1, then  $\neg A$  is an expression of type 1
- ▶ if  $A$  and  $B$  are expressions of type 1, then  $(A \vee B)$  is an expression of type 1
- ▶ if  $x^\tau$  is a bound variable which does not occur in an expression  $A(a^\tau)$  of type 1, then  $\exists x^\tau A(x^\tau)$  is an expression of type 1.
- ▶ if  $x_1^{\tau_1}, \dots, x_n^{\tau_n}$  are different bound variables which do not occur in an expression  $A(a_1^{\tau_1}, \dots, a_n^{\tau_n})$  of type 1, then  $\lambda x_1^{\tau_1} \dots x_n^{\tau_n} A(x_1^{\tau_1}, \dots, x_n^{\tau_n})$  is an expression of type  $(\tau_1, \dots, \tau_n)$ .

# Step 1

## Definition

A *semi-valuation* is an assignment of at most one of the two values  $t$  and  $f$  to any wff with the following conditions:

- ▶ if  $\neg A$  is  $t$ , then  $A$  is  $f$

# Step 1

## Definition

A *semi-valuation* is an assignment of at most one of the two values  $t$  and  $f$  to any wff with the following conditions:

- ▶ if  $\neg A$  is  $t$ , then  $A$  is  $f$
- ▶ if  $\neg A$  is  $f$ , then  $A$  is  $t$

# Step 1

## Definition

A *semi-valuation* is an assignment of at most one of the two values  $t$  and  $f$  to any wff with the following conditions:

- ▶ if  $\neg A$  is  $t$ , then  $A$  is  $f$
- ▶ if  $\neg A$  is  $f$ , then  $A$  is  $t$
- ▶ if  $A \vee B$  is  $t$ , then at least one of the wffs  $A$  and  $B$  is  $t$ .

# Step 1

## Definition

A *semi-valuation* is an assignment of at most one of the two values  $t$  and  $f$  to any wff with the following conditions:

- ▶ if  $\neg A$  is  $t$ , then  $A$  is  $f$
- ▶ if  $\neg A$  is  $f$ , then  $A$  is  $t$
- ▶ if  $A \vee B$  is  $t$ , then at least one of the wffs  $A$  and  $B$  is  $t$ .
- ▶ if  $A \vee B$  is  $f$ , then both wffs  $A$  and  $B$  are  $f$ .

# Step 1

## Definition

A *semi-valuation* is an assignment of at most one of the two values  $t$  and  $f$  to any wff with the following conditions:

- ▶ if  $\neg A$  is  $t$ , then  $A$  is  $f$
- ▶ if  $\neg A$  is  $f$ , then  $A$  is  $t$
- ▶ if  $A \vee B$  is  $t$ , then at least one of the wffs  $A$  and  $B$  is  $t$ .
- ▶ if  $A \vee B$  is  $f$ , then both wffs  $A$  and  $B$  are  $f$ .
- ▶ if  $\exists x^\tau A(x^\tau)$  is  $t$ , then there exists an expression  $e^\tau$  of type  $\tau$  such that  $A(e^\tau)$  is  $t$ .



# Step 1

## Definition

A *semi-valuation* is an assignment of at most one of the two values  $t$  and  $f$  to any wff with the following conditions:

- ▶ if  $\neg A$  is  $t$ , then  $A$  is  $f$
- ▶ if  $\neg A$  is  $f$ , then  $A$  is  $t$
- ▶ if  $A \vee B$  is  $t$ , then at least one of the wffs  $A$  and  $B$  is  $t$ .
- ▶ if  $A \vee B$  is  $f$ , then both wffs  $A$  and  $B$  are  $f$ .
- ▶ if  $\exists x^\tau A(x^\tau)$  is  $t$ , then there exists an expression  $e^\tau$  of type  $\tau$  such that  $A(e^\tau)$  is  $t$ .
- ▶ if  $\exists x^\tau A(x^\tau)$  is  $f$ , then  $A(e^\tau)$  is  $f$  for every expression  $e^\tau$  of type  $\tau$ .

# Step 1

## Definition

A *semi-valuation* is an assignment of at most one of the two values  $t$  and  $f$  to any wff with the following conditions:

- ▶ if  $\neg A$  is  $t$ , then  $A$  is  $f$
- ▶ if  $\neg A$  is  $f$ , then  $A$  is  $t$
- ▶ if  $A \vee B$  is  $t$ , then at least one of the wffs  $A$  and  $B$  is  $t$ .
- ▶ if  $A \vee B$  is  $f$ , then both wffs  $A$  and  $B$  are  $f$ .
- ▶ if  $\exists x^\tau A(x^\tau)$  is  $t$ , then there exists an expression  $e^\tau$  of type  $\tau$  such that  $A(e^\tau)$  is  $t$ .
- ▶ if  $\exists x^\tau A(x^\tau)$  is  $f$ , then  $A(e^\tau)$  is  $f$  for every expression  $e^\tau$  of type  $\tau$ .
- ▶ if  $(e_1, \dots, e_n \in \lambda x_1 \dots x_n A(x_1, \dots, x_n))$  has a value, then  $A(e_1, \dots, e_n)$  has the same value.

# Step 1.1

## Definition

- ▶ Every variable or constant of type 1 is a *prime formula*

# Step 1.1

## Definition

- ▶ Every variable or constant of type 1 is a *prime formula*
- ▶ A formula  $(e_1, \dots, e_n \in e)$  is a *prime formula* if  $e$  is a free variable or a constant.

# Step 1.1

## Definition

- ▶ Every variable or constant of type 1 is a *prime formula*
- ▶ A formula  $(e_1, \dots, e_n \in e)$  is a *prime formula* if  $e$  is a free variable or a constant.

## Definition

By recursion.

- ▶  $F$  is a *positive part* (pp) of  $F$

# Step 1.1

## Definition

- ▶ Every variable or constant of type 1 is a *prime formula*
- ▶ A formula  $(e_1, \dots, e_n \in e)$  is a *prime formula* if  $e$  is a free variable or a constant.

## Definition

By recursion.

- ▶  $F$  is a *positive part* (pp) of  $F$
- ▶ if  $\neg A$  is a pp of  $F$ , then  $A$  is a *negative part* (np) of  $F$

# Step 1.1

## Definition

- ▶ Every variable or constant of type 1 is a *prime formula*
- ▶ A formula  $(e_1, \dots, e_n \in e)$  is a *prime formula* if  $e$  is a free variable or a constant.

## Definition

By recursion.

- ▶  $F$  is a *positive part* (pp) of  $F$
- ▶ if  $\neg A$  is a pp of  $F$ , then  $A$  is a *negative part* (np) of  $F$
- ▶ if  $\neg A$  is an np of  $F$ , then  $A$  is a pp of  $F$

# Step 1.1

## Definition

- ▶ Every variable or constant of type 1 is a *prime formula*
- ▶ A formula  $(e_1, \dots, e_n \in e)$  is a *prime formula* if  $e$  is a free variable or a constant.

## Definition

By recursion.

- ▶  $F$  is a *positive part* (pp) of  $F$
- ▶ if  $\neg A$  is a pp of  $F$ , then  $A$  is a *negative part* (np) of  $F$
- ▶ if  $\neg A$  is an np of  $F$ , then  $A$  is a pp of  $F$
- ▶ if  $A \vee B$  is a pp of  $F$ , then  $A$  and  $B$  are pps of  $F$



# Step 1.1

## Definition

- ▶ Every variable or constant of type 1 is a *prime formula*
- ▶ A formula  $(e_1, \dots, e_n \in e)$  is a *prime formula* if  $e$  is a free variable or a constant.

## Definition

By recursion.

- ▶  $F$  is a *positive part* (pp) of  $F$
- ▶ if  $\neg A$  is a pp of  $F$ , then  $A$  is a *negative part* (np) of  $F$
- ▶ if  $\neg A$  is an np of  $F$ , then  $A$  is a pp of  $F$
- ▶ if  $A \vee B$  is a pp of  $F$ , then  $A$  and  $B$  are pps of  $F$

Remark: if  $A$  is a true pp or false np of  $F$ , then  $F$  is true.

## Step 1.2

### Definition

- ▶ Every axiom  $F[P_+, P_-]$ , where  $P$  is a prime formula, is *derivable*.

## Step 1.2

### Definition

- ▶ Every axiom  $F[P_+, P_-]$ , where  $P$  is a prime formula, is *derivable*.
- ▶ If the premises of a basic inference are *derivable*, then the conclusion is derivable.

## Step 1.2

### Definition

- ▶ Every axiom  $F[P_+, P_-]$ , where  $P$  is a prime formula, is *derivable*.
- ▶ If the premises of a basic inference are *derivable*, then the conclusion is derivable.
- ▶ A wff is *strictly derivable* if no use is made of S5.

## Step 1.2

### Definition

- ▶ Every axiom  $F[P_+, P_-]$ , where  $P$  is a prime formula, is *derivable*.
- ▶ If the premises of a basic inference are *derivable*, then the conclusion is derivable.
- ▶ A wff is *strictly derivable* if no use is made of S5.

The basic inferences:

$$S1 \frac{F[A_-], F[B_-]}{F[(A \vee B)_-]}$$

$$S2 \frac{F[A(a^\tau)_-]}{F[\exists x^\tau A(x^\tau)_-]}$$

$$S3 \frac{F[\exists x^\tau A(x^\tau)_+] \vee A(e^\tau)}{F[\exists x^\tau A(x^\tau)_+]}$$

$$S4a \frac{F[A(e_1, \dots, e_n)_+]}{F[(e_1, \dots, e_n \in \lambda x_1 \dots x_n A(x_1, \dots, x_n))_+]}$$

$$S4b \frac{F[A(e_1, \dots, e_n)_-]}{F[(e_1, \dots, e_n \in \lambda x_1 \dots x_n A(x_1, \dots, x_n))_-]}$$

$$S5 \frac{F \vee \exists x^1 \neg (x^1 \vee \neg x^1)}{F}$$

## Step 1.3

### Definition

A *deduction string* of a wff  $F$  is a finite or infinite sequence of wffs  $F_1, F_2, F_3, \dots$  that satisfies:

- ▶  $F_1 = F$

## Step 1.3

### Definition

A *deduction string* of a wff  $F$  is a finite or infinite sequence of wffs  $F_1, F_2, F_3, \dots$  that satisfies:

- ▶  $F_1 = F$
- ▶ if  $F_n$  is neither an axiom nor a primitive wff, then  $F_{n+1}$  is a reductum or a weakening of  $F_n$ .

## Step 1.3

### Definition

A *deduction string* of a wff  $F$  is a finite or infinite sequence of wffs  $F_1, F_2, F_3, \dots$  that satisfies:

- ▶  $F_1 = F$
- ▶ if  $F_n$  is neither an axiom nor a primitive wff, then  $F_{n+1}$  is a reductum or a weakening of  $F_n$ .
- ▶ if  $F_n$  is an axiom or a primitive wff, then  $F_n$  is the last wff of the string.



## Step 1.3

### Definition

A *deduction string* of a wff  $F$  is a finite or infinite sequence of wffs  $F_1, F_2, F_3, \dots$  that satisfies:

- ▶  $F_1 = F$
- ▶ if  $F_n$  is neither an axiom nor a primitive wff, then  $F_{n+1}$  is a reductum or a weakening of  $F_n$ .
- ▶ if  $F_n$  is an axiom or a primitive wff, then  $F_n$  is the last wff of the string.

No use is made of rule S5.

## Step 1.4

### Theorem

*If every deduction string of a wff  $F$  contains an axiom, then  $F$  is strictly derivable.*

## Step 1.4

### Theorem

*If every deduction string of a wff  $F$  contains an axiom, then  $F$  is strictly derivable.*

### Proof.

All the deduction strings can be combined together to form a deduction tree of  $F$ . This deduction tree is finite, because there are finitely many deduction strings and the tree has finite branching. Only inferences S1-S4 are being used in the derivation, so no use is made of the cut rule. □

## Step 1.5

### Theorem

*If a wff  $F$  has a deduction string which does not contain an axiom, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

## Step 1.5

### Theorem

*If a wff  $F$  has a deduction string which does not contain an axiom, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

### Proof.

Let  $\phi$  be the deduction string  $(F =) F_1, F_2, F_3, \dots$  not containing an axiom. We define  $V(e) = t$  for  $e$  a negative part in  $\phi$  and  $V(e) = f$  for  $e$  a positive part in  $\phi$ .  $V$  is a semi-valuation, by induction on the rank of a wff  $A$ :

## Step 1.5

### Theorem

*If a wff  $F$  has a deduction string which does not contain an axiom, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

### Proof.

Let  $\phi$  be the deduction string  $(F =) F_1, F_2, F_3, \dots$  not containing an axiom. We define  $V(e) = t$  for  $e$  a negative part in  $\phi$  and  $V(e) = f$  for  $e$  a positive part in  $\phi$ .  $V$  is a semi-valuation, by induction on the rank of a wff  $A$ :

- IB: There is no prime formula  $P$  with  $V(P) = t = f$ .

## Step 1.5

### Theorem

*If a wff  $F$  has a deduction string which does not contain an axiom, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

### Proof.

Let  $\phi$  be the deduction string  $(F =) F_1, F_2, F_3, \dots$  not containing an axiom. We define  $V(e) = t$  for  $e$  a negative part in  $\phi$  and  $V(e) = f$  for  $e$  a positive part in  $\phi$ .  $V$  is a semi-valuation, by induction on the rank of a wff  $A$ :

- ▶ IB: There is no prime formula  $P$  with  $V(P) = t = f$ .
- ▶ IS:  $V$  satisfies all the conditions of a semi-valuation.

## Step 1.5

### Theorem

*If a wff  $F$  has a deduction string which does not contain an axiom, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

### Proof.

Let  $\phi$  be the deduction string  $(F =) F_1, F_2, F_3, \dots$  not containing an axiom. We define  $V(e) = t$  for  $e$  a negative part in  $\phi$  and  $V(e) = f$  for  $e$  a positive part in  $\phi$ .  $V$  is a semi-valuation, by induction on the rank of a wff  $A$ :

- ▶ IB: There is no prime formula  $P$  with  $V(P) = t = f$ .
- ▶ IS:  $V$  satisfies all the conditions of a semi-valuation.

Also, because  $F$  is a positive part of itself:  $V(F) = f$ .





# Interlude

We proved:

# Interlude

We proved:

## Theorem

*If every deduction string of a wff  $F$  contains an axiom, then  $F$  is strictly derivable.*

# Interlude

We proved:

## Theorem

*If every deduction string of a wff  $F$  contains an axiom, then  $F$  is strictly derivable.*

And we proved also:

# Interlude

We proved:

## Theorem

*If every deduction string of a wff  $F$  contains an axiom, then  $F$  is strictly derivable.*

And we proved also:

## Theorem

*If a wff  $F$  has a deduction string which does not contain an axiom, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

# Interlude

We proved:

## Theorem

*If every deduction string of a wff  $F$  contains an axiom, then  $F$  is strictly derivable.*

And we proved also:

## Theorem

*If a wff  $F$  has a deduction string which does not contain an axiom, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

Combining this, we get:

# Interlude

We proved:

## Theorem

*If every deduction string of a wff  $F$  contains an axiom, then  $F$  is strictly derivable.*

And we proved also:

## Theorem

*If a wff  $F$  has a deduction string which does not contain an axiom, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

Combining this, we get:

## Theorem

*If a wff  $F$  is not derivable without cut, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

# Interlude

We proved:

## Theorem

*If every deduction string of a wff  $F$  contains an axiom, then  $F$  is strictly derivable.*

And we proved also:

## Theorem

*If a wff  $F$  has a deduction string which does not contain an axiom, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

Combining this, we get:

## Theorem

*If a wff  $F$  is not derivable without cut, then there exists a semi-valuation  $V$  in which  $F$  is  $f$ .*

And thereby we proved the first step.

## Step 2

Assume a semi-valuation  $V$  fixed.

### Definition

*Basic set* contains:

- ▶ variables of any type



## Step 2

Assume a semi-valuation  $V$  fixed.

### Definition

*Basic set* contains:

- ▶ variables of any type
- ▶ arbitrarily many constants of certain types

## Step 2

Assume a semi-valuation  $V$  fixed.

### Definition

*Basic set* contains:

- ▶ variables of any type
- ▶ arbitrarily many constants of certain types
- ▶ arbitrarily many function symbols with certain argument places.

## Step 2

Assume a semi-valuation  $V$  fixed.

### Definition

*Basic set* contains:

- ▶ variables of any type
- ▶ arbitrarily many constants of certain types
- ▶ arbitrarily many function symbols with certain argument places.

### Definition

*Basic set*  $S$ :

- ▶  $P = \{e \mid V(e) \text{ is defined}\}$

## Step 2

Assume a semi-valuation  $V$  fixed.

### Definition

*Basic set* contains:

- ▶ variables of any type
- ▶ arbitrarily many constants of certain types
- ▶ arbitrarily many function symbols with certain argument places.

### Definition

*Basic set*  $S$ :

- ▶  $P = \{e \mid V(e) \text{ is defined}\}$
- ▶  $Q = \{s \mid s \text{ is a symbol occurring in } P\}$

## Step 2

Assume a semi-valuation  $V$  fixed.

### Definition

*Basic set* contains:

- ▶ variables of any type
- ▶ arbitrarily many constants of certain types
- ▶ arbitrarily many function symbols with certain argument places.

### Definition

*Basic set*  $S$ :

- ▶  $P = \{e \mid V(e) \text{ is defined}\}$
- ▶  $Q = \{s \mid s \text{ is a symbol occurring in } P\}$
- ▶  $R = \{e \mid e \text{ is expression with only symbols from } Q\}$

## Step 2

Assume a semi-valuation  $V$  fixed.

### Definition

*Basic set* contains:

- ▶ variables of any type
- ▶ arbitrarily many constants of certain types
- ▶ arbitrarily many function symbols with certain argument places.

### Definition

*Basic set*  $S$ :

- ▶  $P = \{e \mid V(e) \text{ is defined}\}$
- ▶  $Q = \{s \mid s \text{ is a symbol occurring in } P\}$
- ▶  $R = \{e \mid e \text{ is expression with only symbols from } Q\}$
- ▶  $S = \text{least basic set containing } R$

## Step 2.1

### Definition

*Entities* of finite type are defined by induction on  $\tau$ :

- ▶  $E$  is an entity of type 0 iff  $E$  is an expression of type 0 over  $S$ .

## Step 2.1

### Definition

*Entities* of finite type are defined by induction on  $\tau$ :

- ▶  $E$  is an entity of type 0 iff  $E$  is an expression of type 0 over  $S$ .
- ▶  $E$  is an entity of type 1 iff  $E$  is  $t$  or  $f$ .



## Step 2.1

### Definition

*Entities* of finite type are defined by induction on  $\tau$ :

- ▶  $E$  is an entity of type 0 iff  $E$  is an expression of type 0 over  $S$ .
- ▶  $E$  is an entity of type 1 iff  $E$  is  $t$  or  $f$ .
- ▶  $E$  is an entity of type  $(\tau_1, \dots, \tau_n)$  iff  $E$  is a set of  $n$ -tuples  $((e_1, E_1), \dots, (e_n, E_n))$  where  $e_i$  is an expression of type  $\tau_i$  over  $S$  and  $E_i$  is an entity of type  $\tau_i$ .

## Step 2.2

### Definition

*Possible values of expressions.* By induction over the types:

## Step 2.2

### Definition

*Possible values of expressions.* By induction over the types:

- ▶ if  $e$  is of type 0, then it is a possible value of itself.

## Step 2.2

### Definition

*Possible values of expressions.* By induction over the types:

- ▶ if  $e$  is of type 0, then it is a possible value of itself.
- ▶ if  $e$  is of type 1, then  $E$  is a possible value of  $e$  iff

## Step 2.2

### Definition

*Possible values of expressions.* By induction over the types:

- ▶ if  $e$  is of type 0, then it is a possible value of itself.
- ▶ if  $e$  is of type 1, then  $E$  is a possible value of  $e$  iff
  - ▶  $E$  is  $t$  or  $f$ , and

## Step 2.2

### Definition

*Possible values of expressions.* By induction over the types:

- ▶ if  $e$  is of type 0, then it is a possible value of itself.
- ▶ if  $e$  is of type 1, then  $E$  is a possible value of  $e$  iff
  - ▶  $E$  is  $t$  or  $f$ , and
  - ▶ if  $V$  assigns some value to  $e$ , then  $E = V(e)$ .

## Step 2.2

### Definition

*Possible values of expressions.* By induction over the types:

- ▶ if  $e$  is of type 0, then it is a possible value of itself.
- ▶ if  $e$  is of type 1, then  $E$  is a possible value of  $e$  iff
  - ▶  $E$  is  $t$  or  $f$ , and
  - ▶ if  $V$  assigns some value to  $e$ , then  $E = V(e)$ .
- ▶ if  $e$  is of type  $(\tau_1, \dots, \tau_n)$ , then  $E$  is a possible value of  $e$  iff:

## Step 2.2

### Definition

*Possible values of expressions.* By induction over the types:

- ▶ if  $e$  is of type 0, then it is a possible value of itself.
- ▶ if  $e$  is of type 1, then  $E$  is a possible value of  $e$  iff
  - ▶  $E$  is  $t$  or  $f$ , and
  - ▶ if  $V$  assigns some value to  $e$ , then  $E = V(e)$ .
- ▶ if  $e$  is of type  $(\tau_1, \dots, \tau_n)$ , then  $E$  is a possible value of  $e$  iff:
  - ▶  $E$  is a set of elements are  $n$ -tuples  $((e_1, E_1), \dots, (e_n, E_n))$  where  $e_i$  is an expression of type  $\tau_i$  over  $S$  and  $E_i$  is some possible value of  $e_i$ .



## Step 2.2

### Definition

*Possible values of expressions.* By induction over the types:

- ▶ if  $e$  is of type 0, then it is a possible value of itself.
- ▶ if  $e$  is of type 1, then  $E$  is a possible value of  $e$  iff
  - ▶  $E$  is  $t$  or  $f$ , and
  - ▶ if  $V$  assigns some value to  $e$ , then  $E = V(e)$ .
- ▶ if  $e$  is of type  $(\tau_1, \dots, \tau_n)$ , then  $E$  is a possible value of  $e$  iff:
  - ▶  $E$  is a set of elements are  $n$ -tuples  $((e_1, E_1), \dots, (e_n, E_n))$  where  $e_i$  is an expression of type  $\tau_i$  over  $S$  and  $E_i$  is some possible value of  $e_i$ .
  - ▶ if  $V$  assigns  $t$  to an expression  $(e_1, \dots, e_n) \in e$  over  $S$  and each  $E_i$  is a possible value of  $e_i$ , then  $((e_1, E_1), \dots, (e_n, E_n))$  belongs to  $E$ .

## Step 2.2

### Definition

*Possible values of expressions.* By induction over the types:

- ▶ if  $e$  is of type 0, then it is a possible value of itself.
- ▶ if  $e$  is of type 1, then  $E$  is a possible value of  $e$  iff
  - ▶  $E$  is  $t$  or  $f$ , and
  - ▶ if  $V$  assigns some value to  $e$ , then  $E = V(e)$ .
- ▶ if  $e$  is of type  $(\tau_1, \dots, \tau_n)$ , then  $E$  is a possible value of  $e$  iff:
  - ▶  $E$  is a set of elements are  $n$ -tuples  $((e_1, E_1), \dots, (e_n, E_n))$  where  $e_i$  is an expression of type  $\tau_i$  over  $S$  and  $E_i$  is some possible value of  $e_i$ .
  - ▶ if  $V$  assigns  $t$  to an expression  $(e_1, \dots, e_n) \in e$  over  $S$  and each  $E_i$  is a possible value of  $e_i$ , then  $((e_1, E_1), \dots, (e_n, E_n))$  belongs to  $E$ .
  - ▶ if  $V$  assigns  $f$  to an expression  $(e_1, \dots, e_n) \in e$  over  $S$  and each  $E_i$  is a possible value of  $e_i$ , then  $((e_1, E_1), \dots, (e_n, E_n))$  does not belong to  $E$ .

## Step 2.3

### Definition

*New constants.* To each pair  $(e, E)$  where  $e$  is an expression of type  $\tau$  and  $E$  is a possible value of  $e$ , we assign a constant  $c_{e,E}$  of type  $\tau$ , different from each other and from all the symbols in the basic set  $S$ . The set of these constants is  $S_c$  and we define  $S' = S \cup S_c$ .

## Step 2.3

### Definition

*New constants.* To each pair  $(e, E)$  where  $e$  is an expression of type  $\tau$  and  $E$  is a possible value of  $e$ , we assign a constant  $c_{e,E}$  of type  $\tau$ , different from each other and from all the symbols in the basic set  $S$ . The set of these constants is  $S_c$  and we define  $S' = S \cup S_c$ .

### Definition

If  $e$  is an expression over  $S'$ , then  $e^*$  is the expression obtained from  $e$  by replacing each constant  $c_{e',E}$  with  $e'$ .

## Step 2.4

### Definition

$V'$ , the extension of  $V$  defined over expressions  $e$  over  $S'$ .

## Step 2.4

### Definition

$V'$ , the extension of  $V$  defined over expressions  $e$  over  $S'$ .

- ▶ if  $e$  is a constant  $c_{e',E}$ , then  $V'(e) = E$ .

## Step 2.4

### Definition

$V'$ , the extension of  $V$  defined over expressions  $e$  over  $S'$ .

- ▶ if  $e$  is a constant  $c_{e',E}$ , then  $V'(e) = E$ .
- ▶ if  $e$  is not in  $S_c$  and is of type 0, then  $V'(e) = e^*$ .

## Step 2.4

### Definition

$V'$ , the extension of  $V$  defined over expressions  $e$  over  $S'$ .

- ▶ if  $e$  is a constant  $c_{e',E}$ , then  $V'(e) = E$ .
- ▶ if  $e$  is not in  $S_c$  and is of type 0, then  $V'(e) = e^*$ .
- ▶ if  $e$  is not in  $S_c$  and is of type 1, then  $V'(e) = t$  if  $V(e) = t$  and otherwise  $V'(e) = f$ .



## Step 2.4

### Definition

$V'$ , the extension of  $V$  defined over expressions  $e$  over  $S'$ .

- ▶ if  $e$  is a constant  $c_{e',E}$ , then  $V'(e) = E$ .
- ▶ if  $e$  is not in  $S_c$  and is of type 0, then  $V'(e) = e^*$ .
- ▶ if  $e$  is not in  $S_c$  and is of type 1, then  $V'(e) = t$  if  $V(e) = t$  and otherwise  $V'(e) = f$ .
- ▶ if  $e$  is of type  $(\tau_1, \dots, \tau_n)$ , then  $V'(e)$  is the set of all  $n$ -tuples  $((e_1, E_1), \dots, (e_n, E_n))$  such that  $E_i$  is a possible value of  $e_i$  and  $V'((e_1, \dots, e_n \in e)) = t$ .

## Step 2.4

### Definition

$V'$ , the extension of  $V$  defined over expressions  $e$  over  $S'$ .

- ▶ if  $e$  is a constant  $c_{e',E}$ , then  $V'(e) = E$ .
- ▶ if  $e$  is not in  $S_c$  and is of type 0, then  $V'(e) = e^*$ .
- ▶ if  $e$  is not in  $S_c$  and is of type 1, then  $V'(e) = t$  if  $V(e) = t$  and otherwise  $V'(e) = f$ .
- ▶ if  $e$  is of type  $(\tau_1, \dots, \tau_n)$ , then  $V'(e)$  is the set of all  $n$ -tuples  $((e_1, E_1), \dots, (e_n, E_n))$  such that  $E_i$  is a possible value of  $e_i$  and  $V'((e_1, \dots, e_n \in e)) = t$ .
- ▶ if  $e$  is  $(e_1, \dots, e_n \in e)$ , then  $V'(e) = t$  if  $((e_1^*, V'(e_1)), \dots, (e_n^*, V'(e_n)))$  belongs to  $V'(e)$  and  $V'(e) = f$  otherwise.

## Step 2.4, continued

### Definition

- ▶ if  $e$  is  $\neg A$ , then  $V'(e) = t$  if  $V'(A) = f$  and vice versa.

## Step 2.4, continued

### Definition

- ▶ if  $e$  is  $\neg A$ , then  $V'(e) = t$  if  $V'(A) = f$  and vice versa.
- ▶ if  $e$  is  $A \vee B$ , then  $V'(e) = t$  if  $V'(A) = t$  or  $V'(B) = t$  and  $V'(e) = f$  otherwise.

## Step 2.4, continued

### Definition

- ▶ if  $e$  is  $\neg A$ , then  $V'(e) = t$  if  $V'(A) = f$  and vice versa.
- ▶ if  $e$  is  $A \vee B$ , then  $V'(e) = t$  if  $V'(A) = t$  or  $V'(B) = t$  and  $V'(e) = f$  otherwise.
- ▶ if  $e$  is  $\exists x^\tau A(x^\tau)$ , then  $V'(e) = t$  if  $V'(A(c)) = t$  for some constant  $c$  of type  $\tau$  in  $S_c$  and  $V'(e) = f$  if  $V'(A(c)) = f$  for all such constants.

## Step 2.4, continued

### Definition

- ▶ if  $e$  is  $\neg A$ , then  $V'(e) = t$  if  $V'(A) = f$  and vice versa.
- ▶ if  $e$  is  $A \vee B$ , then  $V'(e) = t$  if  $V'(A) = t$  or  $V'(B) = t$  and  $V'(e) = f$  otherwise.
- ▶ if  $e$  is  $\exists x^\tau A(x^\tau)$ , then  $V'(e) = t$  if  $V'(A(c)) = t$  for some constant  $c$  of type  $\tau$  in  $S_c$  and  $V'(e) = f$  if  $V'(A(c)) = f$  for all such constants.
- ▶ if  $e$  is  $\lambda x_1^{\tau_1} \dots x_n^{\tau_n} A(x_1^{\tau_1} \dots x_n^{\tau_n})$ , then  $V'(e)$  is the set of all  $n$ -tuples  $((e_1, E_1), \dots, (e_n, E_n))$  such that  $e_i$  is an expression of type  $\tau_i$  over  $S$ ,  $E_i$  is a possible value of  $e_i$  and  $V'(A(x_1^{\tau_1} \dots x_n^{\tau_n})) = t$ .

## Step 2.5

### Theorem

*For each expression  $e$  over  $S_1$ ,  $V'(e)$  is a possible value of  $e^*$ .*

## Step 2.5

### Theorem

*For each expression  $e$  over  $S_1$ ,  $V'(e)$  is a possible value of  $e^*$ .*

### Proof.

By induction over the length of  $e$ . There are several cases to consider.



## Step 2.5

### Theorem

*For each expression  $e$  over  $S_1$ ,  $V'(e)$  is a possible value of  $e^*$ .*

### Proof.

By induction over the length of  $e$ . There are several cases to consider.

- ▶  $e$  is  $c_{e',E}$ . Then, by definition,  $E$  is a possible value of  $e' = e^*$  and by definition of  $V'$ ,  $V'(e) = E$ .

## Step 2.5

### Theorem

*For each expression  $e$  over  $S_1$ ,  $V'(e)$  is a possible value of  $e^*$ .*

### Proof.

By induction over the length of  $e$ . There are several cases to consider.

- ▶  $e$  is  $c_{e',E}$ . Then, by definition,  $E$  is a possible value of  $e' = e^*$  and by definition of  $V'$ ,  $V'(e) = E$ .
- ▶  $e$  is of type 0 or a constant or variable in  $S$ . Then by definition  $V'(e)$  is a possible value of  $e^*$ .

## Step 2.5

### Theorem

*For each expression  $e$  over  $S_1$ ,  $V'(e)$  is a possible value of  $e^*$ .*

### Proof.

By induction over the length of  $e$ . There are several cases to consider.

- ▶  $e$  is  $c_{e', E}$ . Then, by definition,  $E$  is a possible value of  $e' = e^*$  and by definition of  $V'$ ,  $V'(e) = E$ .
- ▶  $e$  is of type 0 or a constant or variable in  $S$ . Then by definition  $V'(e)$  is a possible value of  $e^*$ .
- ▶  $e$  is  $(e_1, \dots, e_n \in e')$ . Assume  $V$  assigns some value to  $e^*$ . We have  $e^* = (e_1^*, \dots, e_n^* \in e'^*)$ . By the IH, for all  $i$ ,  $V'(e_i)$  is a possible value  $E_i$  of  $e_i^*$ . In case  $V(e^*) = t$ , since  $V'(e')$  is a possible value of  $e'^*$ , it follows that  $((e_1^*, E_1), \dots, (e_n^*, E_n))$  belongs to  $V'(e')$ , so  $V'(e) = t$ . Similar in case  $V(e^*) = f$ .



## Step 2.5, continued

- ▶  $e$  is  $\neg A$ . Assume  $V$  assigns some value to  $e^*$ . We have  $e^* = \neg A^*$ , so if  $V(e^*)$  is  $t$ , then  $V(A^*) = f$ . By IH  $V'(A) = f$  and hence  $V'(e) = t$ . Similar if  $V(e^*)$  is  $f$ .

## Step 2.5, continued

- ▶  $e$  is  $\neg A$ . Assume  $V$  assigns some value to  $e^*$ . We have  $e^* = \neg A^*$ , so if  $V(e^*)$  is  $t$ , then  $V(A^*) = f$ . By IH  $V'(A) = f$  and hence  $V'(e) = t$ . Similar if  $V(e^*)$  is  $f$ .
- ▶  $e$  is  $A \vee B$ . Assume  $V$  assigns some value to  $e^*$ . We have  $e^* = A^* \vee B^*$ . If  $V(e^*) = t$ , then  $V(A^*) = t$  or  $V(B^*) = t$ . By IH  $V'(A) = t$  or  $V'(B) = t$  and hence  $V'(e) = t$ . Similar if  $V(e^*) = f$ .

## Step 2.5, continued

- ▶  $e$  is  $\neg A$ . Assume  $V$  assigns some value to  $e^*$ . We have  $e^* = \neg A^*$ , so if  $V(e^*)$  is  $t$ , then  $V(A^*) = f$ . By IH  $V'(A) = f$  and hence  $V'(e) = t$ . Similar if  $V(e^*)$  is  $f$ .
- ▶  $e$  is  $A \vee B$ . Assume  $V$  assigns some value to  $e^*$ . We have  $e^* = A^* \vee B^*$ . If  $V(e^*) = t$ , then  $V(A^*) = t$  or  $V(B^*) = t$ . By IH  $V'(A) = t$  or  $V'(B) = t$  and hence  $V'(e) = t$ . Similar if  $V(e^*) = f$ .
- ▶  $e$  is  $\exists x^\tau A(x^\tau)$  and assume  $V$  assigns some value to  $e^*$ . We have  $(\exists x^\tau A(x^\tau))^* = \exists x^\tau (A(x^\tau))^*$ . If  $V(e^*) = t$ , then  $V((A(e^\tau))^*) = t$  for some expression  $e^\tau$  over  $S$ . For each possible value  $E$  of  $e^\tau$ ,  $(A(c_{e^\tau, E}))^* = A^*(e^\tau)$ . By IH  $V'(A(c_{e^\tau, E})) = t$  for some  $c_{e^\tau, E}$  and therefore  $V'(e) = t$ . Similar if  $V(e^*) = f$ .

## Step 2.5, continued

- ▶  $e$  is  $\lambda x_1^{\tau_1} \dots x_n^{\tau_n} A(x_1^{\tau_1} \dots x_n^{\tau_n})$ . By construction,  $V'(e)$  is the set of  $n$ -tuples  $((e_1, E_1), \dots, (e_n, E_n))$ , where  $e_i$  is an expression over  $S$  of type  $\tau_i$  and  $E_i$  is a corresponding possible value. Assume  $V$  assigns  $t$  to  $(e_1, \dots, e_n \in e^*)$  and  $E_1, \dots, E_n$  are possible values for  $e_1, \dots, e_n$ . We note that  $e^* = \lambda x_1^{\tau_1} \dots x_n^{\tau_n} A^*(x_1^{\tau_1} \dots x_n^{\tau_n})$ . Now:  $V(A^*(e_1, \dots, e_n)) = t$ . Since  $A(c_{e_1, E_1}, \dots, c_{e_n, E_n})^* = A^*(e_1, \dots, e_n)$ , it follows from the IH that  $V'(A(c_{e_1, E_1}, \dots, c_{e_n, E_n})) = t$  and hence  $((e_1, E_1), \dots, (e_n, E_n))$  belongs to  $V'(e)$ .

QED.

## Step 2.6

Two useful results.

### Lemma

*For each expression  $e$  over  $S'$ , there is a constant  $c_{e^*,E}$  such that  $V'(e) = V'(c_{e^*,E})$ .*



## Step 2.6

Two useful results.

### Lemma

*For each expression  $e$  over  $S'$ , there is a constant  $c_{e^*,E}$  such that  $V'(e) = V'(c_{e^*,E})$ .*

### Proof.

$V'(e)$  is a possible value of  $e^*$ , so there exists a constant  $c_{e^*,V'(e)}$ .  
Now by definition:  $V'(c_{e^*,V'(e)}) = V'(e)$ . □

## Step 2.6

Two useful results.

### Lemma

*For each expression  $e$  over  $S'$ , there is a constant  $c_{e^*,E}$  such that  $V'(e) = V'(c_{e^*,E})$ .*

### Proof.

$V'(e)$  is a possible value of  $e^*$ , so there exists a constant  $c_{e^*,V'(e)}$ .  
Now by definition:  $V'(c_{e^*,V'(e)}) = V'(e)$ . □

### Lemma

*If  $e_2$  is obtained from  $e_1$  by replacing some occurrences of  $c_{e^*,E}$  by  $e$  and  $V'(e) = V'(c_{e^*,E})$  then  $V'(e_1) = V'(e_2)$ .*

## Step 2.6

Two useful results.

### Lemma

*For each expression  $e$  over  $S'$ , there is a constant  $c_{e^*,E}$  such that  $V'(e) = V'(c_{e^*,E})$ .*

### Proof.

$V'(e)$  is a possible value of  $e^*$ , so there exists a constant  $c_{e^*,V'(e)}$ .  
Now by definition:  $V'(c_{e^*,V'(e)}) = V'(e)$ . □

### Lemma

*If  $e_2$  is obtained from  $e_1$  by replacing some occurrences of  $c_{e^*,E}$  by  $e$  and  $V'(e) = V'(c_{e^*,E})$  then  $V'(e_1) = V'(e_2)$ .*

### Proof.

By induction on the length of  $e_1$ . □

## Step 2.6, continued

### Theorem

$V'$  is a total valuation

## Step 2.6, continued

### Theorem

$V'$  is a total valuation

### Proof.

$V'$  assigns exactly one truth value to all the formulas over  $S'$ . By definition, it satisfies the first five conditions of a semi-valuation. The remaining two conditions need to be checked.

## Step 2.6, continued

### Theorem

$V'$  is a total valuation

### Proof.

$V'$  assigns exactly one truth value to all the formulas over  $S'$ . By definition, it satisfies the first five conditions of a semi-valuation. The remaining two conditions need to be checked.

- Suppose  $V'$  assigns  $f$  to  $\exists x^\tau A(x^\tau)$ . Let  $e$  be an expression of type  $\tau$  over  $S'$ . By definition  $V'$  assigns  $f$  to each formula  $A(c_{e^*,E})$ . By a previous lemma,  $V'(e) = V(c_{e^*,E})$  for some constant  $c_{e^*,E}$ . By another previous lemma,  $V'(A(c_{e^*,E})) = V(A(e))$  for such a constant, so  $V'(A(e)) = f$ .

## Step 2.6, continued

### Theorem

$V'$  is a total valuation

### Proof.

$V'$  assigns exactly one truth value to all the formulas over  $S'$ . By definition, it satisfies the first five conditions of a semi-valuation. The remaining two conditions need to be checked.

- ▶ Suppose  $V'$  assigns  $f$  to  $\exists x^\tau A(x^\tau)$ . Let  $e$  be an expression of type  $\tau$  over  $S'$ . By definition  $V'$  assigns  $f$  to each formula  $A(c_{e^*,E})$ . By a previous lemma,  $V'(e) = V(c_{e^*,E})$  for some constant  $c_{e^*,E}$ . By another previous lemma,  $V'(A(c_{e^*,E})) = V(A(e))$  for such a constant, so  $V'(A(e)) = f$ .
- ▶ Suppose  $V'$  assigns  $t$  to  $e_1, \dots, e_n \in \lambda x_1, \dots, x_n A(x_1, \dots, x_n)$ . By definition of  $V'$ ,  $((e_1^*, V'(e_1)), \dots, (e_n^*, V'(e_n)))$  belongs to  $V'(\lambda x_1, \dots, x_n A(x_1, \dots, x_n))$ . So  $V'$  assigns  $t$  to  $A(c_{e_1^*, V'(e_1)}, \dots, c_{e_n^*, V'(e_n)})$ . Because  $V'(c_{e_i^*, V'(e_i)}) = V'(e_i)$  and a lemma,  $V'(A(e_1, \dots, e_n)) = t$ . Idem for  $V'(e) = f$ .

## Interlude 2

Earlier we saw that, for a wff  $F$  which is not derivable without cut, there exists a semi-valuation  $V$  that makes it  $f$ . We have proved that a semi-valuation  $V$  can be extended to a total valuation  $V'$ . Now we need to prove that any wff  $F$  which is  $f$  in a total valuation, is not derivable. The argument runs as follows:



## Interlude 2

Earlier we saw that, for a wff  $F$  which is not derivable without cut, there exists a semi-valuation  $V$  that makes it  $f$ . We have proved that a semi-valuation  $V$  can be extended to a total valuation  $V'$ . Now we need to prove that any wff  $F$  which is  $f$  in a total valuation, is not derivable. The argument runs as follows:

- ▶ Assume  $V(F) = f$  for some total valuation  $V$ .

## Interlude 2

Earlier we saw that, for a wff  $F$  which is not derivable without cut, there exists a semi-valuation  $V$  that makes it  $f$ . We have proved that a semi-valuation  $V$  can be extended to a total valuation  $V'$ . Now we need to prove that any wff  $F$  which is  $f$  in a total valuation, is not derivable. The argument runs as follows:

- ▶ Assume  $V(F) = f$  for some total valuation  $V$ .
- ▶ then there is a semi-valuation in which  $F \vee E$  is false for a certain  $E$ .

## Interlude 2

Earlier we saw that, for a wff  $F$  which is not derivable without cut, there exists a semi-valuation  $V$  that makes it  $f$ . We have proved that a semi-valuation  $V$  can be extended to a total valuation  $V'$ . Now we need to prove that any wff  $F$  which is  $f$  in a total valuation, is not derivable. The argument runs as follows:

- ▶ Assume  $V(F) = f$  for some total valuation  $V$ .
- ▶ then there is a semi-valuation in which  $F \vee E$  is false for a certain  $E$ .
- ▶ then  $F \vee E$  is not strictly derivable.

## Interlude 2

Earlier we saw that, for a wff  $F$  which is not derivable without cut, there exists a semi-valuation  $V$  that makes it  $f$ . We have proved that a semi-valuation  $V$  can be extended to a total valuation  $V'$ . Now we need to prove that any wff  $F$  which is  $f$  in a total valuation, is not derivable. The argument runs as follows:

- ▶ Assume  $V(F) = f$  for some total valuation  $V$ .
- ▶ then there is a semi-valuation in which  $F \vee E$  is false for a certain  $E$ .
- ▶ then  $F \vee E$  is not strictly derivable.
- ▶ then  $F$  is not derivable.

## Step 3

### Definition

$$E = \exists x^1 \neg(x^1 \vee \neg x^1)$$

## Step 3

### Definition

$$E = \exists x^1 \neg(x^1 \vee \neg x^1)$$

$V'(E) = f$  iff  $V'$  is total. So, if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, namely  $V'$  itself, such that  $V'(F \vee E) = f$ .

## Step 3.1

### Definition

The *reducible parts* of a wff  $F$  are its negative parts of the form  $(A \vee B)$  and  $\exists x A(x)$  and the positive and negative parts of the form  $(e_1, \dots, e_n \in \lambda x_1, \dots, x_n A(x_1, \dots, x_n))$ . i.e. corresponding to conclusions of rules S1, S2, S4a and S4b.

## Step 3.1

### Definition

The *reducible parts* of a wff  $F$  are its negative parts of the form  $(A \vee B)$  and  $\exists x A(x)$  and the positive and negative parts of the form  $(e_1, \dots, e_n \in \lambda x_1, \dots, x_n A(x_1, \dots, x_n))$ . i.e. corresponding to conclusions of rules S1, S2, S4a and S4b.

### Definition

The *rank* of a wff  $F$  is the length of the longest subformula string of  $F$ .



## Step 3.1

### Definition

The *reducible parts* of a wff  $F$  are its negative parts of the form  $(A \vee B)$  and  $\exists xA(x)$  and the positive and negative parts of the form  $(e_1, \dots, e_n \in \lambda x_1, \dots, x_n A(x_1, \dots, x_n))$ . i.e. corresponding to conclusions of rules S1, S2, S4a and S4b.

### Definition

The *rank* of a wff  $F$  is the length of the longest subformula string of  $F$ .

### Definition

The *reducibility rank* of a wff  $F$  is the ordinal number  $\omega r + s$ , where  $r$  is the maximal rank of reducible parts of  $F$  and  $s$  is the number of reducible parts of  $F$ .

## Step 3.1, continued

### Definition

The *derivability order* is defined inductively:

## Step 3.1, continued

### Definition

The *derivability order* is defined inductively:

- ▶ axioms have derivability order 0.

## Step 3.1, continued

### Definition

The *derivability order* is defined inductively:

- ▶ axioms have derivability order 0.
- ▶ if premises of S1 have derivability orders  $n_1$  and  $n_2$ , the the conclusion has derivability order  $\max(n_1, n_2) + 1$

## Step 3.1, continued

### Definition

The *derivability order* is defined inductively:

- ▶ axioms have derivability order 0.
- ▶ if premises of S1 have derivability orders  $n_1$  and  $n_2$ , the the conclusion has derivability order  $\max(n_1, n_2) + 1$
- ▶ if premises of any other basic inference has derivability order  $n$ , then the conclusion has derivability order  $n + 1$ .

## Step 3.1, continued

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

## Step 3.1, continued

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

Let  $F$  be a wff of reducibility rank  $\rho$  which has derivability order  $n$ .  
We use two IHs:

## Step 3.1, continued

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

Let  $F$  be a wff of reducibility rank  $\rho$  which has derivability order  $n$ .  
We use two IHs:

- ▶ for every wff  $F$ , strictly derivable with order  $< n$  there is no semi-valuation in which  $F$  is  $f$ .



## Step 3.1, continued

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

Let  $F$  be a wff of reducibility rank  $\rho$  which has derivability order  $n$ .  
We use two IHs:

- ▶ for every wff  $F$ , strictly derivable with order  $< n$  there is no semi-valuation in which  $F$  is  $f$ .
- ▶ for every wff  $F$  with reducibility rank  $< \rho$  which is strictly derivable with order  $< n$  there is no semi-valuation in which  $F$  is  $f$ .

There are the following cases:

## Step 3.1, continued

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

Let  $F$  be a wff of reducibility rank  $\rho$  which has derivability order  $n$ . We use two IHs:

- ▶ for every wff  $F$ , strictly derivable with order  $< n$  there is no semi-valuation in which  $F$  is  $f$ .
- ▶ for every wff  $F$  with reducibility rank  $< \rho$  which is strictly derivable with order  $< n$  there is no semi-valuation in which  $F$  is  $f$ .

There are the following cases:

- ▶ if  $F$  is an axiom, i.e. of the form  $G(P_+, P_-)$  then it cannot be  $f$  in any semi-valuation.

## Step 3.1, continued

- ▶ if  $F$  is of the form  $G((A \vee B)_-)$ , then  $G(A_-)$  and  $G(B_-)$  are strictly derivable with order  $\leq n$  and reducibility rank  $< \rho$ . So there is no semi-valuation in which they are  $f$ , so there is no semi-valuation in which  $F$  is  $f$ .

## Step 3.1, continued

- ▶ if  $F$  is of the form  $G((A \vee B)_-)$ , then  $G(A_-)$  and  $G(B_-)$  are strictly derivable with order  $\leq n$  and reducibility rank  $< \rho$ . So there is no semi-valuation in which they are  $f$ , so there is no semi-valuation in which  $F$  is  $f$ .
- ▶ if  $F$  is of the form  $G(e_1, \dots, e_n \in \lambda x_1, \dots, x_n A(x_1, \dots, x_n))$ , then  $G(A(e_1, \dots, e_n))$  is strictly derivable with order  $\leq n$  and reducibility rank  $< \rho$ . So, by IH there is no semi-valuation making  $G(A(e_1, \dots, e_n))$  false, so the same thing holds for  $F$ .

## Step 3.1, continued

- ▶ if  $F$  is of the form  $G((A \vee B)_-)$ , then  $G(A_-)$  and  $G(B_-)$  are strictly derivable with order  $\leq n$  and reducibility rank  $< \rho$ . So there is no semi-valuation in which they are  $f$ , so there is no semi-valuation in which  $F$  is  $f$ .
- ▶ if  $F$  is of the form  $G(e_1, \dots, e_n \in \lambda x_1, \dots, x_n A(x_1, \dots, x_n))$ , then  $G(A(e_1, \dots, e_n))$  is strictly derivable with order  $\leq n$  and reducibility rank  $< \rho$ . So, by IH there is no semi-valuation making  $G(A(e_1, \dots, e_n))$  false, so the same thing holds for  $F$ .
- ▶ if  $F$  is of the form  $G(\exists x^\tau A(x^\tau)_-)$ , then ... (complicated)

## Step 3.1, continued

- ▶ if  $F$  is of the form  $G((A \vee B)_-)$ , then  $G(A_-)$  and  $G(B_-)$  are strictly derivable with order  $\leq n$  and reducibility rank  $< \rho$ . So there is no semi-valuation in which they are  $f$ , so there is no semi-valuation in which  $F$  is  $f$ .
- ▶ if  $F$  is of the form  $G(e_1, \dots, e_n \in \lambda x_1, \dots, x_n A(x_1, \dots, x_n))$ , then  $G(A(e_1, \dots, e_n))$  is strictly derivable with order  $\leq n$  and reducibility rank  $< \rho$ . So, by IH there is no semi-valuation making  $G(A(e_1, \dots, e_n))$  false, so the same thing holds for  $F$ .
- ▶ if  $F$  is of the form  $G(\exists x^\tau A(x^\tau)_-)$ , then ... (complicated)
- ▶ if  $F$  is of the form  $G(\exists x^\tau A(x^\tau)_+)$  and follows from rule S3. Then the premise of that rule is strictly derivable with order  $n - 1$ . By IH, there is no semi-valuation with  $F$  false.

## Step 3.2

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

## Step 3.2

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

### Proof.

By induction on order  $n$ .

- ▶ if  $F$  is an axiom, then  $F \vee E$  is also an axiom.



## Step 3.2

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

### Proof.

By induction on order  $n$ .

- ▶ if  $F$  is an axiom, then  $F \vee E$  is also an axiom.
- ▶ if  $F$  is the conclusion of a basic inference S1, S2 or S4, whose premises  $F_i$  are derivable with orders  $< n$ , then  $F_i \vee E$  are strictly derivable by IH, so by applying S1, S2, S4,  $F \vee E$  is strictly derivable.

## Step 3.2

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

### Proof.

By induction on order  $n$ .

- ▶ if  $F$  is an axiom, then  $F \vee E$  is also an axiom.
- ▶ if  $F$  is the conclusion of a basic inference S1, S2 or S4, whose premises  $F_i$  are derivable with orders  $< n$ , then  $F_i \vee E$  are strictly derivable by IH, so by applying S1, S2, S4,  $F \vee E$  is strictly derivable.
- ▶ if  $F$  is the conclusion of a basic inference S3, whose premise  $F \vee G$  is derivable with order  $n - 1$ , then  $F \vee E$  is strictly derivable by a short inference.

## Step 3.2

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

### Proof.

By induction on order  $n$ .

- ▶ if  $F$  is an axiom, then  $F \vee E$  is also an axiom.
- ▶ if  $F$  is the conclusion of a basic inference S1, S2 or S4, whose premises  $F_i$  are derivable with orders  $< n$ , then  $F_i \vee E$  are strictly derivable by IH, so by applying S1, S2, S4,  $F \vee E$  is strictly derivable.
- ▶ if  $F$  is the conclusion of a basic inference S3, whose premise  $F \vee G$  is derivable with order  $n - 1$ , then  $F \vee E$  is strictly derivable by a short inference.
- ▶ if  $F$  is the conclusion of a basic inference S5, whose premise  $F \vee E$  is derivable with order  $n - 1$ , then  $F \vee E$  is strictly derivable by a short inference.

## Step 3.3

So, we've seen:

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ .

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

and



## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

and

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

and

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

So,

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

and

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

So,

- If  $V(F) = f$  for some total valuation  $V$ .

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

and

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

So,

- ▶ If  $V(F) = f$  for some total valuation  $V$ .
- ▶ then there is a semi-valuation in which  $F \vee E$  is false.

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

and

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

So,

- ▶ If  $V(F) = f$  for some total valuation  $V$ .
- ▶ then there is a semi-valuation in which  $F \vee E$  is false.
- ▶ so,  $F \vee E$  is not strictly derivable.

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

and

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

So,

- ▶ If  $V(F) = f$  for some total valuation  $V$ .
- ▶ then there is a semi-valuation in which  $F \vee E$  is false.
- ▶ so,  $F \vee E$  is not strictly derivable.
- ▶ so,  $F$  is not derivable.

## Step 3.3

So, we've seen: if  $V'(F) = f$  for some total valuation  $V'$ , then there exists a semi-valuation, such that  $V'(F \vee E) = f$ . and

### Theorem

*If  $F$  is strictly derivable, then there exists no semi-valuation in which  $F$  is  $f$ .*

and

### Theorem

*If  $F$  is derivable, then  $F \vee E$  is strictly derivable.*

So,

- ▶ If  $V(F) = f$  for some total valuation  $V$ .
- ▶ then there is a semi-valuation in which  $F \vee E$  is false.
- ▶ so,  $F \vee E$  is not strictly derivable.
- ▶ so,  $F$  is not derivable.

And we proved the claim: if  $V(F) = f$  for some total valuation  $V$ , then  $F$  is not derivable.

# Summary

Having just proved the last step, we proved all three:



# Summary

Having just proved the last step, we proved all three:

1. if  $A$  is not derivable without cut then there is a semi-valuation with  $A$  false

# Summary

Having just proved the last step, we proved all three:

1. if  $A$  is not derivable without cut then there is a semi-valuation with  $A$  false
2. semi-valuations are extendable to total valuations

# Summary

Having just proved the last step, we proved all three:

1. if  $A$  is not derivable without cut then there is a semi-valuation with  $A$  false
2. semi-valuations are extendable to total valuations
3. if  $A$  is false in a total valuation, then  $A$  is not derivable

# Summary

Having just proved the last step, we proved all three:

1. if  $A$  is not derivable without cut then there is a semi-valuation with  $A$  false
2. semi-valuations are extendable to total valuations
3. if  $A$  is false in a total valuation, then  $A$  is not derivable

So, if  $A$  is derivable, it is derivable without cut.

# References

- ▶ Gaisi Takeuti, 1953, "On a generalized logic calculus".
- ▶ Dag Prawitz, 1968, "Hauptsatz for higher order logic".
- ▶ Motoo Takahashi, 1967, "Proof of cut-elimination in simple type theory".
- ▶ Kurt Schütte, 1960, "Syntactical and semantical properties of type theory."
- ▶ Thierry Coquand, "Type Theory", The Stanford Encyclopedia of Philosophy (Fall 2018 Edition).