



EPEA516

ANALYTICAL SKILLS II

Dr. Harish Mittu
Associate Professor

Learning Outcomes



After this lecture, you will be able to

- solve problems based on arithmetic and geometric progression.

Problem 1

- Is the sequence given by $T_n = 12n - 9$ form an A. P.?
- $T_n = 12n - 9$
- $T_{n-1} = 12(n-1) - 9$
- $T_{n-1} = 12n - 12 - 9$
- $T_{n-1} = 12n - 21$
- $T_n - T_{n-1} = (12n - 9) - (12n - 21)$
 $= \cancel{12}n - 9 - \cancel{12}n + 21$
 $= 12$

Problem 2

- Find the 15th term of the sequence 12, 7, 2, - 3, - 8,
- First term = $a = 12$
- Common Difference = $d = - 5$
- $n = 15$

$$T_n = a + (n - 1) d$$

$$T_{15} = 12 + (15 - 1) (-5)$$

$$T_{15} = 12 - 70$$

$$T_{15} = - 58$$

Problem 3

- Find the sum of the series $5 + 9 + 13 + \dots$ to 50 terms.
- $a = 5$, $d = 9 - 5 = 4$, and $n = 50$

$$S_n = \frac{n}{2} \{2a + (n - 1) d\}$$

$$S_{50} = \frac{50}{2} \{2 \times 5 + (50 - 1) \times 4\}$$

$$S_{50} = 25\{10 + (49) \times 4\}$$

$$S_{50} = 25\{10 + 196\}$$

$$S_{50} = 25\{206\}$$

$$S_{50} = 5150$$

Problem 4

- Find the sum of 30 terms of an A.P., whose first and last terms are 25 and 216 respectively.
- $a = 25$, $l = 216$, and $n = 30$

$$S_n = \frac{n}{2} \{a + l\}$$

$$S_{30} = \frac{30}{2} \{25 + 216\}$$

$$S_{30} = 15\{241\}$$

$$S_{30} = 3615$$

Problem 5

- Find the sum of 20 terms of an A.P., whose common difference and last term are 7 and 93 respectively.
- $d = 7$, $l = 93$, and $n = 20$

$$S_n = \frac{n}{2} [2l - (n - 1) d]$$

$$S_{20} = \frac{20}{2} [2 \times 93 - (20 - 1) \times 7]$$

$$S_{20} = 10 [186 - 133]$$

$$S_{20} = 10 [53]$$

$$S_{20} = 530$$

Problem 6

- If sum of n terms of an A. P. is given by $S_n = 100 n^2 + 25$, then find its n th term.

- $S_n = 100 n^2 + 25$

- $S_{n-1} = 100 (n - 1)^2 + 25$

$$T_n = S_n - S_{n-1}$$

$$T_n = 100 n^2 + 25 - [100 (n - 1)^2 + 25]$$

$$T_n = 100 n^2 + 25 - [100(n^2 - 2n + 1) + 25]$$

$$T_n = 100 n^2 + 25 - [100n^2 - 200n + 100 + 25]$$


$$T_n = \cancel{100 n^2} + \cancel{25} - \cancel{100 n^2} + 200n - 100 - \cancel{25}$$

$$T_n = 200n - 100$$

Problem 7

- Determine x so that $x + 2$, $4x - 6$ and $3x - 2$ are the three consecutive terms of an A.P.
- Three numbers a , b , c are in A.P. iff $b - a = c - b$ iff $a + c = 2b$

• $x + 2$, $4x - 6$, and $3x - 2$



$$x + \cancel{2} + 3x - \cancel{2} = 2(4x - 6)$$

$$4x = 8x - 12$$

$$12 = 8x - 4x$$

Problem 7

$$12 = 8x - 4x$$

$$\cancel{4x} = \cancel{12} \quad 3$$

$$x = 3$$

Three consecutive terms are $x + 2$, $4x - 6$ and $3x - 2$

i.e., $3 + 2$, $4(3) - 6$ and $3(3) - 2$

i.e., 5, 6 and 7

Problem 8

- Find the 7th term from the end of the A. P. of 50 terms whose first term and common difference are 5 and 8 respectively.
- n^{th} term from the end = $(m - n + 1)^{\text{th}}$ term from beginning

$$T_{m-n+1} = a + (m - n) d$$

- $a = 5, d = 8, m = 50, \text{ and } n = 7$

$$T_{50-7+1} = 5 + (50 - 7) \times 8$$

$$T_{44} = 5 + (43) \times 8$$

$$T_{44} = 5 + 344$$

$$T_{44} = 349$$

Problem 9

- Is the sequence 3, -6, 12, -24, form a G. P.?
- $T_1 = 3, T_2 = -6, T_3 = 12, T_4 = -24, T_5 = 48, \dots$
- $\frac{T_2}{T_1} = \frac{-6}{3} = -2$
- $\frac{T_3}{T_2} = \frac{12}{-6} = -2$
- $\frac{T_4}{T_3} = \frac{-24}{12} = -2$
- $\frac{T_2}{T_1} = \frac{T_3}{T_2} = \frac{T_4}{T_3} = -2 \quad (\text{G.P.})$

Problem 10

- Find the 9th term of the sequence 2, 4, 8, 16, 32
- First term = $a = 2$
- Common ratio = $r = \frac{4}{2} = 2$
- $n = 9$

$$T_n = ar^{n-1}$$

$$T_9 = (2) 2^{9-1}$$

$$T_9 = (2) 2^8 = 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2$$

$$T_9 = 512$$

Problem 11

- Find the sum of 7 terms of the series $\frac{1}{4} + \frac{1}{2} + 1 \dots\dots$

- $a = \frac{1}{4}$, $r = \frac{\frac{1}{2}}{\frac{1}{4}}$ or $\frac{1}{\frac{1}{2}} = 2 (> 1)$, and $n = 7$

$$S_n = \frac{a(r^n - 1)}{(r - 1)} = \frac{\frac{1}{4}(2^7 - 1)}{(2 - 1)}$$

$$S_n = \frac{\frac{1}{4}(128 - 1)}{1}$$

$$S_n = \frac{127}{4}$$

Problem 12

- Find the sum of 12 terms of a G.P., whose first term, common ratio, and last terms are 3, (-2) and 48 respectively.

- $a = 3$, $r = -2$, $l = 48$, and $n = 12$

$$S_n = \frac{lr - a}{r - a}$$

$$S_n = \frac{48 \times (-2) - 3}{-2 - 3}$$

$$S_n = \frac{(-96) - 3}{-5} = \frac{-99}{-5}$$

$$S_n = \frac{99}{5}$$

Problem 13

- Find the sum of 5 terms of a G.P., whose first term, and common ratio are 3 and (-2) respectively.
- $a = 3$, $r = -2$ (< 1), and $n = 5$

$$S_n = \frac{a(1 - r^n)}{(1 - r)}$$

$$S_5 = \frac{3(1 - (-2)^5)}{(1 - (-2))}$$

$$S_5 = \frac{3(1 - (-32))}{(1 - (-2))} = \frac{3(1 + 32)}{(1 + 2)} = \frac{\cancel{3}(33)}{\cancel{3}}$$

$$S_5 = 33$$

Problem 14

- Find the sum of the infinite sequence $9, -3, 1, -\frac{1}{3}, \dots$

- $a = 9, r = \frac{-3}{9} = -\frac{1}{3}$ where $|\frac{-1}{3}| < 1$ i.e., $-\frac{1}{3} < r < \frac{1}{3}$

$$S_{\infty} = \frac{a}{(1 - r)}$$


$$S_{\infty} = \frac{9}{(1 - (-\frac{1}{3}))}$$


$$S_{\infty} = \frac{9}{(1 + \frac{1}{3})} = \frac{9}{(\frac{3 + 1}{3})} = \frac{9 \times 3}{4}$$


$$S_{\infty} = \frac{27}{4}$$

Problem 15

- Determine x so that x , $x + 2$, and $x + 6$ are the three consecutive terms of a G.P.
- Three numbers a , b , c are in G.P. iff $\frac{b}{a} = \frac{c}{b}$ iff $b^2 = ac$

x ,


$x + 2$,


and $x + 6$


$$(x + 2)^2 = x(x + 6)$$

$$\cancel{x^2} + 4x + 4 = \cancel{x^2} + 6x$$

Problem 15

$$\cancel{x^2} + 4x + 4 = \cancel{x^2} + 6x$$

$$4 = 6x - 4x$$

$$\cancel{2x} = \cancel{4} \quad 2$$

$$x = 2$$

Three consecutive terms are x , $x + 2$, and $x + 6$

i.e., 2, $2 + 2$, and $2 + 6$

i.e., 2, 4, and 8

Problem 16

- Find the 5th term from the end of the G. P. of 15 terms whose first term and common ratio are 9 and 3 respectively.

- n^{th} term from the end = $(m - n + 1)^{\text{th}}$ term from beginning

- n^{th} term from the end = $T_{m-n+1} = ar^{m-n}$

{Because $T_n = ar^{n-1}$ }

$$T_{m-n+1} = ar^{m-n}$$

- $a = 9, r = 3, m = 15, \text{ and } n = 5$

Problem 16

$$T_{m-n+1} = ar^{m-n}$$

- $a = 9$, $r = 3$, $m = 15$, and $n = 5$

$$T_{15-5+1} = 9 \times (3)^{15-5}$$

$$T_{11} = 9 \times (3)^{10}$$

$$T_{11} = 9 \times 59049 \text{ or } (3)^2 \times (3)^{10}$$

$$T_{11} = 531441 \text{ or } (3)^{12}$$

Problem 17

- Find the 10th term from the end of the G. P. with last term $\frac{512}{729}$ and common ratio $\frac{2}{3}$.

- n^{th} term from the end in terms of last term 'l' & common ratio 'r'


is $T_n = \frac{l}{r^{n-1}}$

$$l = \frac{512}{729}, r = \frac{2}{3}, \text{ and } n = 10$$

$$T_{10} = \frac{\frac{512}{729}}{\left(\frac{2}{3}\right)^{10-1}}$$

Problem 17

- $l = \frac{512}{729}$, $r = \frac{2}{3}$, and $n = 10$

$$T_{10} = \frac{\frac{512}{729}}{\left(\frac{2}{3}\right)^{10-1}} = \frac{\frac{(2)^9}{(3)^6}}{\frac{(2)^9}{(3)^9}}$$


$$T_{10} = \frac{(2)^{9-9}}{(3)^{6-9}} = \frac{(2)^0}{(3)^{-3}}$$

$$T_{10} = 1 \times (3)^3$$

$$T_{10} = 27$$

Conclusion

- Verification of Sequence
 - A.P. or G.P.
- Compute n term of Progression
 - In A.P.
 - $T_n = a + (n - 1) d$
 - $T_n = S_n - S_{n-1}$
 - In G.P.
 - $T_n = ar^{n-1}$
- Three consecutive terms of an A.P. and G.P.

Conclusion

- n^{th} term from the end of a G.P. having m terms is $(m - n + 1)^{\text{th}}$ term from beginning i.e.,

$$T_{m-n+1} = ar^{m-n}$$

- n^{th} term from the end of an A.P. having m terms is $(m - n + 1)^{\text{th}}$ term from beginning i.e.,

$$T_{m-n+1} = a + (m - n) d$$

- n^{th} term from the end of a G.P. in terms of last term 'l' & common ratio 'r' is

$$T_n = \frac{l}{r^n - 1}$$

Conclusion

- Sum of n terms of A.P.

- $S_n = \frac{n}{2} \{2a + (n - 1) d\}$

- $S_n = \frac{n}{2} \{a + l\}$

- $S_n = \frac{n}{2} [2l - (n - 1) d]$

Conclusion

- Sum of n terms of G.P.

- $S_n = \frac{lr - a}{r - a}$

- $S_n = \frac{a(r^n - 1)}{(r - 1)}$ when $r > 1$

- $S_n = \frac{a(1 - r^n)}{(1 - r)}$ when $r < 1$

- Sum of infinite terms of G.P.

- $S_\infty = \frac{a}{(1 - r)}$ when $|r| < 1$ i.e., $-1 < r < 1$

Summary

- Problems
 - Arithmetic Progression
 - Geometric Progression

That's all for now...