

A hand is shown placing a blue L-shaped block on top of a colorful cube constructed from various other blocks. The background is a solid light blue, and the surface is a light-colored wooden table. Several other blocks are scattered on the table in the foreground.

EMTH403

Mathematical Foundation
for Computer Science

Nitin K. Mishra (Ph.D.)

Associate Professor

Lecture Outcomes



After this lecture, you will be able to

- understand what are duals of Boolean expression.
- understand what is a Complemented Lattice.
- understand what is Boolean Algebra.

Duality

To explain the **relationship between the two identities** in each pair we use the concept of a **dual**.

The dual of a Boolean expression is obtained by **interchanging** Boolean sums and Boolean products and interchanging 0s and 1s.

Duality

Ques:- Find the duals of $x(y + 0)$ and $\bar{x} \cdot 1 + (\bar{y} + z)$.

Sol:- Interchanging \cdot signs and $+$ signs and interchanging 0s and 1s in these expressions produces their duals.

The duals are $x + (y \cdot 1)$ and $(x + 0)(yz)$, respectively.

Boolean algebra

A **Boolean algebra** is a set B with two binary operations \vee and \wedge , elements 0 and 1 , and a unary operation $\bar{}$ such that **these properties hold** for all x , y , and z in B :

$$\left. \begin{array}{l} x \vee 0 = x \\ x \wedge 1 = x \end{array} \right\}$$

Identity laws

$$\left. \begin{array}{l} x \vee \bar{x} = 1 \\ x \wedge \bar{x} = 0 \end{array} \right\}$$

Complement laws

$$\left. \begin{array}{l} (x \vee y) \vee z = x \vee (y \vee z) \\ (x \wedge y) \wedge z = x \wedge (y \wedge z) \end{array} \right\}$$

Associative laws

Boolean algebra

A Boolean algebra is a set B with two binary operations \vee and \wedge , elements 0 and 1 , and a unary operation $\bar{}$ such that these properties hold for all x, y , and z in B :

$$\left. \begin{aligned} x \vee y &= y \vee x \\ x \wedge y &= y \wedge x \end{aligned} \right\}$$

Commutative laws

$$\left. \begin{aligned} x \vee (y \wedge z) &= (x \vee y) \wedge (x \vee z) \\ x \wedge (y \vee z) &= (x \wedge y) \vee (x \wedge z) \end{aligned} \right\}$$

Distributive laws

Boolean algebra

Boolean algebras **may also be defined** using the notion of a lattice, discussed earlier.

Lattice L is a **partially ordered set** in which every pair of elements x, y has a least upper bound, denoted by $\text{lub}(x, y)$ and a greatest lower bound denoted by $\text{glb}(x, y)$.

Boolean algebra

Given two elements x and y of L , we can define two operations \vee and \wedge on pairs of elements of L by

$$x \vee y = \text{lub}(x, y) \text{ and } x \wedge y = \text{glb}(x, y).$$

Boolean algebra

For a lattice L to be a Boolean algebra, it must have two properties.

First, it must be **complemented**.

For a lattice to be complemented it must have a least element 0 and a greatest element 1

and for every element x of the lattice there must exist an element x such that $x \vee x = 1$ and $x \wedge x = 0$.

Boolean algebra

For a lattice L to be a Boolean algebra, it must have two properties.





First, it must be **complemented**.

Second, it must be **distributive**.

This means that for every x , y , and z in L , $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$ and $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$.

Complemented Lattice


A complemented lattice is a **bounded lattice** (with least element 0 and greatest element 1), in which **every element a** has a complement, i.e. an element b satisfying $a \vee b = 1$ and $a \wedge b = 0$. Complements **need not be unique**.



Complemented Lattice


D_{30} is a complemented lattice.

Element	Its Complement
1	30
2	15
3	10
5	6
6	5
10	3
15	2
30	1



Complemented Lattice

D_{20} is not complemented lattice.



D_{20}

Element	Its Complement
1	20
2	10
4	5
5	4
10	2
20	1

$$2 \wedge 10 \neq 0 \quad (2 \wedge 10 = 2)$$

Boolean algebra

For a lattice L to be a Boolean algebra, it must have two properties.

First, it must be **complemented**.


Second, it must be **distributive**.

This means that for every x , y , and z in L , $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$ and $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$.

Boolean algebra

Consider the lattices D_{20} of all positive integer divisors of 20.


D_{20} is not a boolean algebra.



Boolean algebra



Consider the lattices D_{30} of all positive integer divisors of 30.

D_{30} is a Boolean algebra.



Boolean algebra

$S = \{a, b, c\}$ and $T = \{2, 3, 5\}$ consider the Hasse diagram of the two lattices $(P(S), \subseteq)$ and $(P(T), \subseteq)$.



Boolean algebra

Ques:- Show that in a Boolean algebra, the idempotent laws $x \vee x = x$ and $x \wedge x = x$ hold for every element x .

Sol:- By the domination, distributive, and identity laws,
$$x \vee x = (x \vee x) \wedge 1 = (x \vee x) \wedge (x \vee \bar{x}) = x \vee (x \wedge \bar{x}) = x \vee 0 = x.$$

Similarly, $x \wedge x = (x \wedge x) \vee 0 = (x \wedge x) \vee (x \wedge \bar{x}) = x \wedge (x \vee \bar{x}) = x \wedge 1 = x.$

That's all for now...