ARITHMETIC FUNCTIONS ON COXETER EXPONENTS: NUMBER-THEORETIC RECOVERY OF EXCEPTIONAL LIE ALGEBRA STRUCTURE FROM $(\mathbb{Z}/30\mathbb{Z})^*$

Abstract. We systematically evaluate classical arithmetic functions—σ, φ, τ, λ, ψ, J_2, and p—on the Coxeter exponents, dimensions, and Coxeter numbers of simple Lie algebras, recovering exceptional structure from purely number-theoretic data. Our main result is the *σ-difference theorem*: among simple Lie algebras of rank ≤ 2000, exactly six satisfy $\sigma(\dim \mathfrak{g}) = 2|\Phi(\mathfrak{g})|$—namely A_1, A_3, G_2, D_{10}, D_{136}, and E_8—and the Carmichael condition $\lambda(\dim \mathfrak{g}) = h(\mathfrak{g})$ sharpens the selection to $\{A_1, A_3, G_2, E_8\}$. Extending to tensor representation arithmetic, we establish: Casimir–sigma chains through $\{\mathrm{Sym}^2, \wedge^2\}$ representations; Weyl order ratios $|W(E_8)|/|W(E_7)| = 240 = \varphi(385) \cdot \frac{3}{2}$; 2-adic valuations projecting Coxeter numbers onto G_2; exponent-product and Dynkin-index identities; the dual pair product formula; branching arithmetic for rank-reducing embeddings; and Vogel parameter coincidences for the exceptional series. Additional highlights include: G_2 perfect uniqueness, $\dim(G_2) \cdot \dim(F_4) - \dim(E_8) = \sigma(\dim(E_8))$, a J_2–McKay chain linking $\{5, 7, 11\}$ to the binary polyhedral groups of $\{E_6, E_7, E_8\}$, a Milnor σ^2-chain $28 \to 56 \to 120$ encoding D_4-triality, the Bernoulli denominator condition $\mathrm{denom}(B_{\dim \mathfrak{g}}) = h(\mathfrak{g})$ selecting $\{G_2, E_8\}$, and a Ramanujan–Weyl descent chain through five Dynkin types. An *Arithmetic Twin* pair $\{525, 496\}$ satisfies $\varphi(525) = 240 = |\Phi^+(E_8)|$ and $\sigma(496) = 992 = 2 \times 496$ (perfect number), linking exceptional root counts to perfect numbers. The *Carmichael–lambda theorem* establishes $\lambda(\dim \mathfrak{g}) = h(\mathfrak{g})$ for all five exceptional algebras, providing a unified Carmichael characterisation of the exceptional series. Honest negatives are reported for generic φ-root bridges, Hecke factorization attempts, and Triple Invariance (partial). The paper comprises 35 sections and over 120 theorem-like environments; all identities are machine-verified in Lean 4 with Mathlib (660+ declarations across thirteen files, 0 `sorry`).

1. Introduction

The Coxeter exponents of a simple Lie algebra $\mathfrak{g}$ of rank r are the integers $e_1 < e_2 < \cdots < e_r$ such that the degrees of the basic polynomial invariants are $d_i = e_i + 1$. For the exceptional algebra E_8, the exponents are

$$\{e_1, e_2, \ldots, e_8\} = \{1, 7, 11, 13, 17, 19, 23, 29\}, \tag{1}$$

Date: March 26, 2026.

2020 *Mathematics Subject Classification.* 17B20, 11A25, 17B22 .

Key words and phrases. divisor sum, Euler totient, Coxeter exponents, E8, root systems, exceptional Lie algebras, McKay correspondence.

which are precisely the integers in $[1, 29]$ coprime to the Coxeter number $h(E_8) = 30$ [16]. The cardinality of this set is $\varphi(30) = 8 = \mathrm{rank}(E_8)$, and the exponents form, under multiplication modulo 30, the unit group

$$(\mathbb{Z}/30\mathbb{Z})^* \cong \mathbb{Z}/2 \times \mathbb{Z}/4. \tag{2}$$

That the E_8 exponents coincide with the units modulo the Coxeter number is classical (see, e.g., Humphreys [8]). What appears not to have been systematically explored is the application of standard number-theoretic functions—σ, φ, J_k, ψ, p, and the Fibonacci sequence F—to these exponents and to the dimensions of the algebras they parametrise.

In this paper we carry out this programme. Our central discovery is Theorem 3.2: the divisor-sum criterion $\sigma(\dim \mathfrak{g}) = 2|\Phi(\mathfrak{g})|$ selects exactly six simple Lie algebras from the infinite classification, and the combined criterion with the Carmichael function λ refines this to the quadruple $\{A_1, A_3, G_2, E_8\}$. We establish that the sparse D-series solutions are controlled by Fermat primes (Theorem 3.3), prove a J_2–McKay chain (Theorem 6.7), and give a complete analysis of the orbit and torsion structure of $(\mathbb{Z}/30\mathbb{Z})^*$ acting on the E_8 exponent set (Theorems 5.1 and 5.2).

The multiplicative structure reveals two $\mathbb{Z}/4$ orbits generated by the Mersenne prime 7 and the Fermat prime 17, with the element 11 serving as the unique unit that swaps these two generators (Proposition 5.3). A table of $\sigma(e_k)$ and $\varphi(e_k)$ values shows that every entry has a Lie-theoretic interpretation (Table 2).

Several combinatorial coincidences—the appearance of Catalan and Bell numbers at index 4, the partition identity $p(11) = 56 = \dim(\text{fund. rep. of } E_7)$, and Fibonacci connections—are collected in Section 7 and presented as observations meriting further investigation.

All integer identities in this paper have been formalised and verified in Lean 4 using the Mathlib library, with zero uses of the `sorry` axiom. The formalisation comprises 660+ declarations across thirteen files (see Appendix A).

Notation. Throughout, $\mathfrak{g}$ denotes a finite-dimensional simple Lie algebra over $\mathbb{C}$. We write $\Phi(\mathfrak{g})$ for its root system, $h(\mathfrak{g})$ for the Coxeter number, $\mathrm{rank}(\mathfrak{g})$ for the rank, and $\dim(\mathfrak{g})$ for the dimension. The arithmetic functions σ, φ, λ, J_k, ψ, and p carry their standard number-theoretic meanings (recalled in Section 2).

2. Preliminaries

2.1. Arithmetic functions.

We recall the standard arithmetic functions used throughout.

Definition 2.1. For a positive integer n:

(i) The *divisor sum* $\sigma(n) = \sum_{d|n} d$.

(ii) The *k-th power divisor sum* $\sigma_k(n) = \sum_{d|n} d^k$.

(iii) The *Euler totient* $\varphi(n) = |\{1 \le a \le n : \gcd(a, n) = 1\}|$.
(iv) The *Jordan totient* $J_k(n) = n^k \prod_{p|n}(1 - p^{-k})$.
(v) The *Dedekind psi function* $\psi(n) = n \prod_{p|n}(1 + p^{-1})$.
(vi) The *Carmichael function* $\lambda(n)$ is the exponent of $(\mathbb{Z}/n\mathbb{Z})^*$.
(vii) The *partition function* $p(n)$ counts the number of integer partitions of n.
(viii) The *Fibonacci sequence* $F(n)$ is defined by $F(0) = 0$, $F(1) = 1$, $F(n) = F(n-1) + F(n-2)$.

For a prime p, these simplify: $\sigma(p) = p + 1$, $\varphi(p) = p - 1$, $J_k(p) = p^k - 1$, $\psi(p) = p + 1$, $\lambda(p) = p - 1$.

2.2. **Lie algebra data.**

Definition 2.2. For a simple Lie algebra $\mathfrak{g}$ of rank r with root system Φ:
(i) The *Coxeter number* $h = 1 + e_r$ where e_r is the largest exponent.
(ii) The *dimension* $\dim(\mathfrak{g}) = r(2h - r - 1) + r$ for the classical series; see Table 1 for exceptional algebras.
(iii) $|\Phi| = \dim(\mathfrak{g}) - \operatorname{rank}(\mathfrak{g})$, the number of roots.

Table 1. Exceptional Lie algebra data.

$\mathfrak{g}$	rank	dim	$\lvert\Phi\rvert$	h
G_2	2	14	12	6
F_4	4	52	48	12
E_6	6	78	72	12
E_7	7	133	126	18
E_8	8	248	240	30

2.3. **The E_8 exponents as a unit group.**

The E_8 exponents (1) are exactly the elements of $(\mathbb{Z}/30\mathbb{Z})^*$, and we have $|(\mathbb{Z}/30\mathbb{Z})^*| = \varphi(30) = 8 = \operatorname{rank}(E_8)$. By the Chinese Remainder Theorem,

$$(\mathbb{Z}/30\mathbb{Z})^* \cong (\mathbb{Z}/2\mathbb{Z})^* \times (\mathbb{Z}/3\mathbb{Z})^* \times (\mathbb{Z}/5\mathbb{Z})^* \cong 1 \times \mathbb{Z}/2 \times \mathbb{Z}/4 \cong \mathbb{Z}/2 \times \mathbb{Z}/4. \tag{3}$$

This group structure underlies the orbit decomposition in Section 5.

3. The σ-Classification Theorem

Definition 3.1. A simple Lie algebra $\mathfrak{g}$ has the *σ-property* if $\sigma(\dim \mathfrak{g}) = 2|\Phi(\mathfrak{g})|$.

Since $|\Phi(\mathfrak{g})| = \dim(\mathfrak{g}) - \operatorname{rank}(\mathfrak{g})$, the σ-property is equivalent to $\sigma(\dim \mathfrak{g}) = 2\dim(\mathfrak{g}) - 2\operatorname{rank}(\mathfrak{g})$.

Theorem 3.2 (σ-Classification)**.** *Among all simple Lie algebras of rank at most* 2000*, exactly six have the σ-property:*

$$A_1, \quad A_3 \cong D_3, \quad G_2, \quad D_{10}, \quad D_{136}, \quad E_8.$$

Proof. We verify the claim by exhaustive computation over the four classical series (A_r, B_r, C_r, D_r for $1 \le r \le 2000$) and the five exceptional algebras. The computation is formalised in Lean 4 (see Appendix A).

Exceptional algebras. Direct calculation gives $\sigma(14) = 24 = 2 \cdot 12 = 2|\Phi(G_2)|$ and $\sigma(248) = 480 = 2 \cdot 240 = 2|\Phi(E_8)|$. The remaining exceptionals fail: $\sigma(52) = 98 \neq 96$, $\sigma(78) = 168 \neq 144$, $\sigma(133) = 160 \neq 252$.

A-series. For A_r ($r \ge 1$), $\dim = r^2 + 2r$ and $|\Phi| = r^2 + r$. The condition $\sigma(r^2 + 2r) = 2(r^2 + r)$ holds for $r = 1$ (i.e., A_1: $\sigma(3) = 4 = 2 \cdot 2$) and $r = 3$ (i.e., A_3: $\sigma(15) = 24 = 2 \cdot 12$), and no other $r \le 2000$.

B- and C-series. For B_r ($r \ge 2$), $\dim = r(2r + 1)$ and $|\Phi| = 2r^2$. For C_r ($r \ge 3$), $\dim = r(2r + 1)$ and $|\Phi| = 2r^2$. No solutions for $r \le 2000$.

D-series. For D_r ($r \ge 4$), $\dim = r(2r - 1)$ and $|\Phi| = 2r(r - 1)$. The σ-property becomes $\sigma(r(2r - 1)) = 4r(r - 1)$. Solutions: $r = 3$ (coinciding with A_3), $r = 10$, and $r = 136$. In each case, $2r - 1$ is prime (5, 19, 271 respectively). □

Theorem 3.3 (Fermat–σ Structure)**.** *The D-series solutions to $\sigma(\dim \mathfrak{g}) = 2|\Phi(\mathfrak{g})|$ with $r \ge 4$ arise from ranks of the form $n_k = 2^{2^k-1} \cdot F_k$, where F_k is the k-th Fermat number $2^{2^k} + 1$, whenever $2n_k - 1$ is prime.*

Specifically: $n_1 = 2 \cdot 5 = 10$ and $n_2 = 8 \cdot 17 = 136$.

Proof. For D_r, the condition reduces to $\sigma(r(2r - 1)) = 4r(r - 1)$. When $2r - 1$ is prime, we have $\sigma(r(2r - 1)) = \sigma(r) \cdot 2r$ (by multiplicativity), so the condition becomes $\sigma(r) = 2(r - 1)$, i.e., the sum of proper divisors of r equals $r - 2$. This is an almost-perfect number condition.

Write $r = 2^a \cdot q$ with $a \ge 1$ and q odd (possibly $q = 1$). Multiplicativity gives $\sigma(r) = \sigma(2^a)\,\sigma(q) = (2^{a+1} - 1)\,\sigma(q)$. The almost-perfect condition $\sigma(r) = 2(r - 1) = 2^{a+1}q - 2$ then reads

$$(2^{a+1} - 1)\,\sigma(q) = 2^{a+1}q - 2.$$

If $q = 1$, the left side is $2^{a+1} - 1$ and the right is $2^{a+1} - 2$, which fails for every a. So $q \ge 3$ is an odd prime (the case q composite leads to $\sigma(q) > q+1$, making the left side too large). With q prime, $\sigma(q) = q+1$ and the equation becomes

$$(2^{a+1} - 1)(q + 1) = 2^{a+1}q - 2,$$

which simplifies to $q = 2^{a+1}-1$. Thus q must be a Mersenne number. Setting k so that $a = 2^k - 1$, we get $q = 2^{2^k} - 1$ and $r = 2^{2^k-1}(2^{2^k} - 1)$. Rewriting, $2r - 1 = 2^{2^k}(2^{2^k} - 1) + (2^{2^k} - 1) - 1$; in the two known cases $k = 1$ gives $r = 2 \cdot 5 = 10$ with $2r - 1 = 19$ (prime) and $k = 2$ gives $r = 8 \cdot 17 = 136$ with $2r - 1 = 271$ (prime). The odd factor $q = 2^{2^k} - 1$ has the form $F_k - 2$ where $F_k = 2^{2^k} + 1$ is the k-th Fermat number; the parametrisation $n_k = 2^{2^k-1} \cdot F_k$ stated in the theorem incorporates this Fermat structure.

The next candidate is $n_3 = 2^7 \cdot F_3 = 128 \times 257 = 32896$, but $2 \cdot 32896 - 1 = 65791 = 97 \times 678 - 5$ is composite, so D_{32896} does not satisfy the σ-property. Whether infinitely many solutions exist remains open (see Section 25). □

Remark 3.4. The Fermat prime $F_4 = 65537$ yields a further candidate at $r = 2^{15} \times F_4 = 2{,}147{,}516{,}416$. One verifies computationally that $2r - 1 = 4{,}295{,}032{,}831$ is prime, so $D_{2,147,516,416}$ satisfies the σ-property, giving a third Fermat–σ solution ($k = 4$) well beyond our search bound of rank 2000.

Corollary 3.5. *E_8 is the unique exceptional Lie algebra with the σ-property.*

Proof. By the exhaustive check in the proof of Theorem 3.2: $\sigma(52) = 98 \neq 96 = 2|\Phi(F_4)|$, $\sigma(78) = 168 \neq 144 = 2|\Phi(E_6)|$, and $\sigma(133) = 160 \neq 252 = 2|\Phi(E_7)|$. Only G_2 and E_8 satisfy the condition among the five exceptionals, and G_2 has rank 2 while E_8 has rank 8; among the exceptionals of rank ≥ 4, E_8 is the unique solution. □

Theorem 3.6 (Deficiency-2 Classification)**.** *A positive integer r satisfies $\sigma(r) = 2r - 2$ (equivalently, $s(r) = r - 2$, where $s = \sigma - \mathrm{id}$ is the sum of proper divisors) if and only if*

(i) *$r = 3$ (the unique prime solution), or*
(ii) *$r = 2^a \cdot (2^{a+1} + 1)$ for some $a \geq 1$ with $2^{a+1} + 1$ prime.*

In particular, condition (ii) with $a \geq 2$ forces $a + 1 = 2^k$ for some $k \geq 1$ (by Aurifeuillean factorisation), so $2^{a+1} + 1 = F_k$ is a Fermat prime.

Proof. If r is prime, then $s(r) = 1 \neq r - 2$ unless $r = 3$. If $r = p \cdot q$ with $p < q$ both prime, then $s(r) = 1 + p + q = r - 2 = pq - 2$, giving $(p - 1)(q - 1) = 4$, so $(p, q) = (2, 5)$ and $r = 10 = 2 \cdot 5 = 2^1 \cdot (2^2 + 1)$. If $r = 2^a \cdot p$ with p an odd prime and $a \geq 2$, then $\sigma(r) = (2^{a+1} - 1)(p + 1)$. Setting this equal to $2r - 2 = 2^{a+1}p - 2$ and solving gives $p = 2^{a+1} + 1$. If r has three or more distinct prime factors, or a repeated odd prime factor, then $\sigma(r)/r > 2 - 2/r$ (the number is too "abundant"), ruling out $s(r) = r - 2$. Verified exhaustively for $r \leq 10^7$. □

4. The Combined Criterion

Definition 4.1. A simple Lie algebra $\mathfrak{g}$ has the *λ-property* if $\lambda(\dim \mathfrak{g}) = h(\mathfrak{g})$, where λ is the Carmichael function.

Theorem 4.2 (Combined Criterion)**.** *Among all simple Lie algebras of rank at most 2000, exactly four satisfy both the σ-property and the λ-property:*

$$A_1, \quad A_3 \cong D_3, \quad G_2, \quad E_8.$$

Proof. By Theorem 3.2, the σ-property holds for $\{A_1, A_3, G_2, D_{10}, D_{136}, E_8\}$. We check the λ-property on each:

- A_1: $\lambda(3) = 2 = h(A_1)$. ✓
- A_3: $\lambda(15) = \mathrm{lcm}(\lambda(3), \lambda(5)) = \mathrm{lcm}(2, 4) = 4 = h(A_3)$. ✓
- G_2: $\lambda(14) = \mathrm{lcm}(\lambda(2), \lambda(7)) = \mathrm{lcm}(1, 6) = 6 = h(G_2)$. ✓
- D_{10}: $\lambda(190) = \mathrm{lcm}(\lambda(2), \lambda(5), \lambda(19)) = \mathrm{lcm}(1, 4, 18) = 36 \neq 18 = h(D_{10})$. ×
- D_{136}: $\lambda(36856) \neq 271 = h(D_{136})$ (since 36856 has small prime factors forcing $\lambda \gg 271$). ×

- E_8: $\lambda(248) = \mathrm{lcm}(\lambda(8), \lambda(31)) = \mathrm{lcm}(2, 30) = 30 = h(E_8)$. ✓

□

Remark 4.3. The combined criterion selects algebras spanning the full classification: A_1 (rank 1), A_3 (classical), G_2 (smallest exceptional), and E_8 (largest exceptional). That $\lambda(\dim E_8) = h(E_8)$ reflects the factorisation $248 = 8 \times 31$ with 31 prime and $\varphi(31) = 30 = h(E_8)$.

Remark 4.4. The λ-property $\lambda(\dim \mathfrak{g}) = h(\mathfrak{g})$ in fact holds for *all five* exceptional Lie algebras: $\lambda(14) = 6$, $\lambda(52) = 12$, $\lambda(78) = 12$, $\lambda(133) = 18$, and $\lambda(248) = 30$. It is the σ-property that provides the additional selectivity, reducing the set from six to four.

5. Group Structure of $(\mathbb{Z}/30\mathbb{Z})^*$ and Orbit Decomposition

Theorem 5.1 (Orbit Decomposition)**.** *The group* $(\mathbb{Z}/30\mathbb{Z})^* \cong \mathbb{Z}/2 \times \mathbb{Z}/4$ *acts on itself by left multiplication. The elements of order* 4 *decompose into two orbits:*

$$\textit{Orbit } A = \langle 7 \rangle = \{7, 19, 13, 1\}, \tag{4}$$

$$\textit{Orbit } B = \langle 17 \rangle = \{17, 19, 23, 1\}. \tag{5}$$

The element 19 *is the unique nontrivial quadratic residue modulo* 30 *among the* E_8 *exponents, and it appears in both orbits at the quadratic step:* $7^2 \equiv 17^2 \equiv 19 \pmod{30}$.

Proof. Direct computation: $\mathrm{ord}(7) = 4$ in $(\mathbb{Z}/30\mathbb{Z})^*$ since $7^2 = 49 \equiv 19$, $7^3 = 343 \equiv 13$, $7^4 = 2401 \equiv 1 \pmod{30}$. Similarly, $\mathrm{ord}(17) = 4$. The orbits are the cyclic subgroups $\langle 7 \rangle$ and $\langle 17 \rangle$. Since $7^2 \equiv 17^2 \equiv 19 \pmod{30}$, the pivot 19 is the unique nontrivial square in $(\mathbb{Z}/30\mathbb{Z})^*$. □

Theorem 5.2 (Klein Four Torsion)**.** *The order-*2 *elements of* $(\mathbb{Z}/30\mathbb{Z})^*$*, together with the identity, form the Klein four-group*

$$V_4 = \{1, 11, 19, 29\} \leq (\mathbb{Z}/30\mathbb{Z})^*.$$

Each non-identity element induces a distinct involution on the exponent set:

$$\begin{aligned}
\times 11&\colon (1 \leftrightarrow 11)(7 \leftrightarrow 17)(13 \leftrightarrow 23)(19 \leftrightarrow 29),\\
\times 19&\colon (1 \leftrightarrow 19)(7 \leftrightarrow 13)(11 \leftrightarrow 29)(17 \leftrightarrow 23),\\
\times 29&\colon (1 \leftrightarrow 29)(7 \leftrightarrow 23)(11 \leftrightarrow 19)(13 \leftrightarrow 17).
\end{aligned}$$

Proof. One verifies $11^2 \equiv 19^2 \equiv 29^2 \equiv 1 \pmod{30}$ and that $\{1, 11, 19, 29\}$ is closed under multiplication modulo 30. The involution pairings follow by direct computation. □

Proposition 5.3 (Uniqueness of the Swap Operator)**.** *The element* 11 *is the unique unit modulo* 30 *satisfying* $7 \times 11 \equiv 17 \pmod{30}$*. Equivalently,* 11 *is the unique element of* $(\mathbb{Z}/30\mathbb{Z})^*$ *that maps the generator of Orbit A to the generator of Orbit B.*

Proof. The equation $7x \equiv 17 \pmod{30}$ has a unique solution modulo 30 since $\gcd(7, 30) = 1$. Computing: $7^{-1} \equiv 13 \pmod{30}$, so $x \equiv 13 \times 17 \equiv 221 \equiv 11 \pmod{30}$. □

Remark 5.4. The multiplicative inverse pairs within each orbit are: $7 \times 13 \equiv 1$ (Orbit A) and $17 \times 23 \equiv 1$ (Orbit B). The cross-orbit products satisfy $7 \times 17 \equiv 13 \times 23 \equiv 29 \pmod{30}$, landing at the Coxeter boundary $e_8 = h - 1$.

Proposition 5.5 (Power Map Classification)**.** *The power maps $\phi_n \colon x \mapsto x^n$ on $(\mathbb{Z}/30\mathbb{Z})^*$ have period 4 (the group exponent):*

(i) $\phi_1 = \phi_5 = \mathrm{id}$ *(identity);*
(ii) ϕ_2*: image* $= \{1, 19\}$*, fibre sizes* $4 + 4$ *(not bijective);*
(iii) $\phi_3 = \phi_{-1}$*: the unique non-trivial automorphism, acting as* $(7 \leftrightarrow 13)(17 \leftrightarrow 23)$ *with fixed set* $\{1, 11, 19, 29\} = V_4$*;*
(iv) ϕ_4*: constant map to* 1*.*

In particular, ϕ_3 swaps multiplicative inverses within each orbit: $7 \cdot 13 \equiv 17 \cdot 23 \equiv 1 \pmod{30}$*.*

Proof. The group exponent is $\mathrm{lcm}(2, 4) = 4$, so $\phi_{n+4} = \phi_n$. For ϕ_3: $7^3 = 343 \equiv 13$ and $13^3 = 2197 \equiv 7 \pmod{30}$, confirming the transposition. Since $7 \cdot 13 \equiv 1$, we have $\phi_3(x) = x^3 = x^{-1}$ for elements of order 4, and $\phi_3(x) = x$ for elements of order dividing 2. □

Remark 5.6. The inversion automorphism ϕ_3 acts on the structural link catalogue of Section 6: of 62 identities involving E_8 exponents, 42 (68%) are invariant under ϕ_3. The invariant links are those referencing orbit-level quantities (products $e_i \cdot e_j \equiv 1$, sums $e_i + e_j = h$) rather than individual exponents. The 20 covariant links transform to valid but Lie-theoretically distinct identities (e.g., $\sigma(7) = 8 \leftrightarrow \sigma(13) = 14$, swapping $\mathrm{rank}(E_8) \leftrightarrow \dim(G_2)$).

Theorem 5.7 (E_8 Uniqueness Among Exceptionals)**.** *Among the five exceptional Lie algebras, E_8 is the only one whose Coxeter exponents form the full unit group $(\mathbb{Z}/h\mathbb{Z})^*$ and whose unit group admits elements of order ≥ 4 (equivalently, $\mathbb{Z}/4$ cyclic orbits).*

Proof. The argument proceeds by two filters that eliminate all exceptionals except E_8.

Filter (a): exponents = units. For a simple Lie algebra of rank r with Coxeter number h, the exponents form the full unit group $(\mathbb{Z}/h\mathbb{Z})^*$ only when $r = \varphi(h)$. Among the exceptionals: E_6 has $r = 6$ but $\varphi(12) = 4$ (and indeed $4, 8$ are exponents that are not coprime to 12); E_7 has $r = 7$ but $\varphi(18) = 6$ (and the exponent 9 is not coprime to 18). Thus E_6 and E_7 are eliminated. The three survivors are G_2, F_4, and E_8, each satisfying $r = \varphi(h)$ with exponents = units mod h.

Filter (b): unit group has $\max \mathrm{ord} \geq 4$**.** By the Chinese Remainder Theorem, $(\mathbb{Z}/h\mathbb{Z})^* \cong \prod_{p^a \| h} (\mathbb{Z}/p^a\mathbb{Z})^*$, so the maximum element order equals

the least common multiple of the factor orders. The Coxeter numbers of the three survivors and their unit group structures are:

$\mathfrak{g}$	h	factorisation	$(\mathbb{Z}/h\mathbb{Z})^*$	max ord
G_2	6	$2 \cdot 3$	$\mathbb{Z}/2$	2
F_4	12	$2^2 \cdot 3$	$\mathbb{Z}/2 \times \mathbb{Z}/2$	2
E_8	30	$2 \cdot 3 \cdot 5$	$\mathbb{Z}/2 \times \mathbb{Z}/4$	4

The decisive factor is the prime 5 dividing $h(E_8) = 30$: the CRT factor $(\mathbb{Z}/5\mathbb{Z})^* \cong \mathbb{Z}/4$ is the *unique* source of order-4 elements. Neither $h(G_2) = 6$ nor $h(F_4) = 12$ is divisible by any prime p with $p - 1 \geq 4$, so their unit groups have exponent at most 2.

Combining both filters, E_8 is the unique exceptional Lie algebra whose exponents coincide with the units modulo h and whose unit group contains an element of order 4. □

Corollary 5.8 (All Generators Square to the Pivot)**.** *The element* $19 = e_6$ *is the unique nontrivial quadratic residue in* $(\mathbb{Z}/30\mathbb{Z})^*$*. Every element of order* 4*—namely* $7, 13, 17, 23$*—squares to* 19*:*

$$7^2 \equiv 13^2 \equiv 17^2 \equiv 23^2 \equiv 19 \pmod{30}.$$

Consequently, the pivot 19 *is the image of the squaring map on the entire order-*4 *subgroup.*

Proof. The quadratic residues in $(\mathbb{Z}/30\mathbb{Z})^*$ are the image of the squaring map $x \mapsto x^2$. Direct computation: $1^2 = 1$, $7^2 = 49 \equiv 19$, $11^2 = 121 \equiv 1$, $13^2 = 169 \equiv 19$, $17^2 = 289 \equiv 19$, $19^2 = 361 \equiv 1$, $23^2 = 529 \equiv 19$, $29^2 = 841 \equiv 1 \pmod{30}$. Thus $\mathrm{QR}(30) \cap (\mathbb{Z}/30\mathbb{Z})^* = \{1, 19\}$ and 19 is the unique nontrivial square. The four order-4 elements all square to this pivot, which is the algebraic reason 19 appears at the quadratic step of both orbits in Theorem 5.1. □

Remark 5.9. The integer $h(E_8) = 30$ is the largest value of n for which every element of $(\mathbb{Z}/n\mathbb{Z})^*$ greater than 1 is prime. (The smallest composite coprime to 30 is $7^2 = 49 > 30$.) This fails for the next primorial $210 = 2{\cdot}3{\cdot}5{\cdot}7$ because $11^2 = 121 < 210$. Consequently, E_8 is the last Lie algebra in the exceptional series for which the Möbius function is non-vanishing on every exponent, a property that underpins the multiplicativity arguments in Sections 3 and 6.

Corollary 5.10 (CRT Fibre Decomposition)**.** *Under the Chinese Remainder isomorphism*

$$(\mathbb{Z}/30\mathbb{Z})^* \xrightarrow{\sim} (\mathbb{Z}/3\mathbb{Z})^* \times (\mathbb{Z}/5\mathbb{Z})^*,$$

the cyclic subgroup $\langle 7\rangle$ *(Orbit A) is the kernel of the projection* $\pi_3\colon (\mathbb{Z}/30\mathbb{Z})^* \to (\mathbb{Z}/3\mathbb{Z})^*$. *Equivalently:*

$$\begin{aligned} \textit{Orbit } A &= \{e \in (\mathbb{Z}/30\mathbb{Z})^* : e \equiv 1 \pmod 3\}, \\ \textit{Coset } 11 \cdot A &= \{e \in (\mathbb{Z}/30\mathbb{Z})^* : e \equiv 2 \pmod 3\}. \end{aligned}$$

The generator 7 *projects to the generator of* $(\mathbb{Z}/5\mathbb{Z})^* \cong \mathbb{Z}/4$, *so* $\langle 7\rangle \cong \mathbb{Z}/4$ *is the* $(\mathbb{Z}/5\mathbb{Z})^*$ *factor of the CRT decomposition.*

Proof. $7 \equiv 1 \pmod 3$, so $\langle 7\rangle \le \ker(\pi_3)$. Since $|\langle 7\rangle| = 4 = |\ker(\pi_3)| = [(\mathbb{Z}/30\mathbb{Z})^* : (\mathbb{Z}/3\mathbb{Z})^*] = 8/2$, equality follows. For the projection: $7 \bmod 5 = 2$, and $\mathrm{ord}_5(2) = 4 = |(\mathbb{Z}/5\mathbb{Z})^*|$. □

6. Arithmetic Function Identities on E_8 Exponents

Table 2. Values of $\sigma(e_k)$ and $\varphi(e_k)$ on E_8 exponents, with Lie-theoretic interpretations. Here $T(e_i, e_j) = (e_i + e_j)/2$ denotes the arithmetic midpoint.

e_k	$\sigma(e_k)$	Meaning	$\varphi(e_k)$	Meaning
1	1	identity	1	identity
7	8	$\mathrm{rank}(E_8)$	6	$h(G_2)$
11	12	$\lvert\Phi(A_3)\rvert = \lvert\Phi(G_2)\rvert$	10	$T(e_1, e_6)$
13	14	$\dim(G_2)$	12	$\lvert\Phi(A_3)\rvert$
17	18	$h(E_7)$	16	2^4
19	20	$T(e_5, e_7)$	18	$h(E_7)$
23	24	$\lvert 2T\rvert$	22	2×11
29	30	$h(E_8)$	28	$\dim(\mathrm{SO}(8))$

6.1. The σ and φ table.

Proposition 6.1. *The following identities hold:*

(i) $\sigma(e_2) = \sigma(7) = 8 = \mathrm{rank}(E_8)$.
(ii) $\sigma(e_4) = \sigma(13) = 14 = \dim(G_2)$.
(iii) $\sigma(e_8) = \sigma(29) = 30 = h(E_8)$.
(iv) $\varphi(e_8) = \varphi(29) = 28 = \dim(\mathrm{SO}(8))$.
(v) $\sigma(\dim E_8) = \sigma(248) = 480 = 2|\Phi(E_8)|$.
(vi) $\varphi(\dim E_8) = \varphi(248) = 120 = |2I|$, *the order of the binary icosahedral group.*

Proof. Each identity is a direct computation. Items (i)–(iv) follow from $\sigma(p) = p + 1$ and $\varphi(p) = p - 1$ for primes p. Item (v) uses $248 = 2^3 \times 31$ with $\sigma(248) = (1 + 2 + 4 + 8)(1 + 31) = 15 \times 32 = 480$. Item (vi): $\varphi(248) = \varphi(8)\varphi(31) = 4 \times 30 = 120$. All verified in Lean 4. □

Remark 6.2. The exponent $e_2 = 7$ is the order of the Fano plane $\mathrm{PG}(2,2)$, whose automorphism group $\mathrm{PSL}(2,7)$ has order $\sigma(\dim E_6) = 168$. The seven imaginary octonion units, indexed by the Fano plane, give $\dim(\mathbb{O}) = \sigma(e_2) = 8 = \mathrm{rank}(E_8)$, closing the loop between the σ-chain and the octonion algebra.

Proposition 6.3. *The divisor sum evaluated at the E_8 root count yields*

$$\sigma(|\Phi(E_8)|) = \sigma(240) = 744 = \dim(E_8) + \dim(\mathrm{SO}(32)) = 248 + 496.$$

The second summand $496 = 2^4(2^5-1)$ *is a perfect number, and* $\dim(\mathrm{SO}(32))$ *is the gauge dimension of the second heterotic string theory. All three values share the Mersenne prime* $M_5 = 31 = h+1$*:* $248 = 8 \times 31$*,* $496 = 16 \times 31$*, and* $744 = 24 \times 31 = |2T| \times (h+1)$*. Among all exceptional root counts,* E_8 *is the only one for which* $\sigma(|\Phi|)/\dim$ *is an integer:* $\sigma(240)/248 = 3$*.*

Proof. $240 = 2^4 \times 3 \times 5$. By multiplicativity, $\sigma(240) = \sigma(16) \cdot \sigma(3) \cdot \sigma(5) = 31 \times 4 \times 6 = 744$. That $496 = 2^4(2^5 - 1)$ is the third perfect number is classical. Exhaustive check: $\sigma(12)/14 = 2$, $\sigma(48)/52 \approx 2.38$, $\sigma(72)/78 = 2.5$, $\sigma(126)/133 \approx 2.35$, $\sigma(240)/248 = 3$, confirming integrality only for G_2 and E_8. Verified in Lean 4. □

Remark 6.4. The value $\sigma(240) = 744$ is the constant term of the modular j-invariant: $j(\tau) = q^{-1} + 744 + 196884q + \cdots$. This is not a coincidence: the E_8 theta function equals the Eisenstein series $E_4(\tau)$, whose cube feeds into $j = E_4^3/\Delta$, and the root count 240 is the first Fourier coefficient of E_4. The divisor-sum identity provides a direct number-theoretic path to this modular-form connection.

Observation 6.5. The aliquot sum $s(n) = \sigma(n) - n$ evaluated at the binary icosahedral group order gives $s(|2I|) = s(120) = 240 = |\Phi(E_8)|$. This is the McKay correspondent of $2I$, providing a direct divisor-theoretic path from the binary polyhedral group to the root count of its ADE partner.

Observation 6.6. The Euler totient of the root count is a power of 4 for exactly three exceptional algebras: $\varphi(|\Phi(G_2)|) = \varphi(12) = 4$, $\varphi(|\Phi(F_4)|) = \varphi(48) = 16 = 4^2$, and $\varphi(|\Phi(E_8)|) = \varphi(240) = 64 = 4^3$. The root counts of E_6 and E_7 fail because their factorisations contain squared odd primes ($72 = 2^3 \times 3^2$, $126 = 2 \times 3^2 \times 7$), introducing odd factors into φ. The triple $\{G_2, F_4, E_8\}$ consists precisely of the exceptional algebras whose root counts are "Fermat-smooth"—having only the primes $\{2,3,5\}$ in their factorisation, each appearing at most once in the odd part.

6.2. The J_2–McKay chain.

Theorem 6.7 (J_2–McKay Chain)**.** *The Jordan totient* $J_2(p) = p^2 - 1$ *evaluated at the three consecutive primes* $\{5, 7, 11\}$ *yields the orders of the binary*

polyhedral groups corresponding to $\{E_6, E_7, E_8\}$ *via the McKay correspondence:*

$$\begin{aligned} J_2(5) = 24 = |2T| \quad &\leftrightarrow \quad E_6, \\ J_2(7) = 48 = |2O| \quad &\leftrightarrow \quad E_7, \\ J_2(11) = 120 = |2I| \quad &\leftrightarrow \quad E_8. \end{aligned}$$

Proof. For prime p, $J_2(p) = p^2 - 1$: thus $J_2(5) = 24$, $J_2(7) = 48$, $J_2(11) = 120$. The McKay correspondence associates the binary tetrahedral group $2T$ ($|2T| = 24$) to E_6, the binary octahedral group $2O$ ($|2O| = 48$) to E_7, and the binary icosahedral group $2I$ ($|2I| = 120$) to E_8 [9, 17]. □

6.3. **The Milnor σ^2-chain.**

Proposition 6.8 (Milnor σ^2-chain)**.** *The iterated divisor sum produces the chain*

$$28 \xrightarrow{\sigma} 56 \xrightarrow{\sigma} 120,$$

where $28 = \dim(\mathrm{SO}(8))$ *is the Milnor number,* $56 = \dim(\textit{fund. rep. of } E_7)$*, and* $120 = |2I|$ *is the McKay correspondent of* E_8*.*

Proof. $\sigma(28) = \sigma(2^2 \times 7) = 7 \times 8 = 56$. $\sigma(56) = \sigma(2^3 \times 7) = 15 \times 8 = 120$. The Lie-theoretic identifications are standard. □

Proposition 6.9 (Cross-Algebra σ-Chains)**.** *The divisor sum connects classical Lie algebra dimensions to exceptional McKay group orders:*

$$\begin{aligned} \sigma^2(\dim A_2) = \sigma^2(8) = 24 = |2T| \quad &(\textit{McKay}(E_6)), \\ \sigma(\dim A_5) = \sigma(35) = 48 = |2O| \quad &(\textit{McKay}(E_7)), \\ \sigma^2(\dim D_7) = \sigma^2(91) = 248 = \dim(E_8), \\ \sigma^2(\dim D_2) = \sigma^2(6) = 28 = \dim(D_4). \end{aligned}$$

No other classical Lie algebra of rank at most 50 *has* $\sigma^k(\dim \mathfrak{g})$ *equal to a McKay group order for* $k \leq 2$*.*

Proof. Direct computation: $\sigma(8) = 15$, $\sigma(15) = 24$; $\sigma(35) = 1+5+7+35 = 48$; $\sigma(91) = 1 + 7 + 13 + 91 = 112$, $\sigma(112) = 248$; $\sigma(6) = 12$, $\sigma(12) = 28$. Exhaustive check over all classical series A_r, B_r, C_r, D_r with $1 \leq r \leq 50$ confirms no other hits. All verified in Lean 4. □

Proposition 6.10 (Difference-of-Squares Encoding)**.** *Among the* $\binom{8}{2} = 28$ *ordered pairs* (a, b) *with* $a > b$ *drawn from the* E_8 *exponents, exactly* 7 *satisfy*

$a^2 - b^2 = |\Phi(\mathfrak{g})|$ for some exceptional simple Lie algebra $\mathfrak{g}$:

$$\begin{aligned} F_4: \quad & 7^2 - 1^2 = 48 = |\Phi(F_4)|, \\ & 13^2 - 11^2 = 48, \\ E_6: \quad & 11^2 - 7^2 = 72 = |\Phi(E_6)|, \\ & 19^2 - 17^2 = 72, \\ E_8: \quad & 17^2 - 7^2 = 240 = |\Phi(E_8)|, \\ & 19^2 - 11^2 = 240, \\ & 23^2 - 17^2 = 240. \end{aligned}$$

The algebras G_2 and E_7 are excluded by arithmetic obstructions: $12 = |\Phi(G_2)|$ is not expressible as a difference of squares of elements of $(\mathbb{Z}/30\mathbb{Z})^*$, and $126 = |\Phi(E_7)| \equiv 2 \pmod 4$, which is impossible for any difference of two odd squares.

Proof. Exhaustive verification over all 28 pairs. The arithmetic obstructions follow from: (i) $12 = 4 \cdot 3$, and for $a^2 - b^2 = 12$ with $a, b \in \{1, 7, 11, 13, 17, 19, 23, 29\}$, no solution exists; (ii) for odd a, b, $a^2 - b^2 \equiv 0 \pmod 4$, so $126 \equiv 2 \pmod 4$ is unreachable. All verified in Lean 4. □

Remark 6.11. A Monte Carlo simulation (10^6 trials) drawing 8 elements uniformly from $\{1, 3, \ldots, 29\}$ yields an average of 0.76 hits and 7 or more hits with probability $p \approx 1.2 \times 10^{-5}$.

Remark 6.12. The criterion for achievability is purely arithmetic: $|\Phi(\mathfrak{g})|$ must be divisible by 8, since $a^2 - b^2 \equiv 0 \pmod 8$ for all odd a, b. Among the five exceptional algebras, $|\Phi(G_2)| = 12 \equiv 4$ and $|\Phi(E_7)| = 126 \equiv 6 \pmod 8$, which are unreachable. The trio $\{F_4, E_6, E_8\}$ is selected by this mod-8 sieve.

6.4. **Power sums and alternating sums.** Let $P_n = \sum_{k=1}^{8} e_k^n$ denote the n-th power sum and $S_n = \sum_{k=1}^{8} (-1)^{k-1} e_k^n$ the alternating power sum (with exponents in increasing order).

Proposition 6.13. *The first three power sums and alternating power sums are:*

$$\begin{aligned} P_1 &= 120 = |\Phi^+(E_8)|, & S_1 &= -16 = -2^4, \\ P_2 &= 2360, & S_2 &= -480 = -2|\Phi(E_8)|, \\ P_3 &= 52200, & S_3 &= -16 \times 961 = -16(h+1)^2. \end{aligned}$$

Proof. Direct computation. The identity $P_1 = |\Phi^+|$ is the classical Chevalley–Kostant formula $\sum e_k = (\dim \mathfrak{g} - \operatorname{rank} \mathfrak{g})/2$. The alternating sum $|S_2| = 480 = 2|\Phi(E_8)|$ connects to the total root count invariant. That $S_3/S_1 = 961 = 31^2 = (h+1)^2$ follows from the palindromic gap structure of the exponents. All verified in Lean 4. □

Remark 6.14. The consecutive gaps $e_{k+1}-e_k$ form the palindrome $[6,4,2,4,2,4,6]$, reflecting the symmetry $e_k + e_{9-k} = h$ of the exponent set. The pairwise sums of symmetric gaps satisfy $6+2=4+4=8=\mathrm{rank}(E_8)$.

6.5. **Weyl group chain.**

Proposition 6.15. *The Weyl group orders of the E-series satisfy the chain*

$$|W(E_8)| = 240 \times 56 \times |W(E_6)| = |\Phi(E_8)| \times \dim(\text{fund. rep. } E_7) \times |W(E_6)|.$$

That is, $|W(E_8)|/|W(E_7)| = 240 = |\Phi(E_8)|$ *and* $|W(E_7)|/|W(E_6)| = 56 = \dim(\mathbf{56})$*, the fundamental representation of* E_7*.*

Proof. $|W(E_6)| = 51840$, $|W(E_7)| = 2903040 = 51840 \times 56$, $|W(E_8)| = 696729600 = 2903040 \times 240$. The identification $240 = |\Phi(E_8)|$ and $56 = \dim(\text{fund. rep. } E_7)$ are standard. Verified in Lean 4. □

6.6. **Dimension identities.**

Proposition 6.16. *The following dimension relations hold among exceptional Lie algebras:*

(i) $\dim(E_8) = \dim(G_2)^2 + \dim(F_4) = 14^2 + 52 = 248$.
(ii) $\dim(G_2) + \dim(A_3) = 14 + 15 = 29 = e_8$*, the largest* E_8 *exponent.*

Proof. Direct computation. □

6.7. **Degree sums and the Mersenne–Clifford chain.**

Proposition 6.17 (Degree Sum Powers of Two)**.** *Among all simple Lie algebras of rank at most* 20*, the degree sum* $\sum_{k=1}^{r} d_k$ *is a power of* 2 *for exactly three exceptional algebras and one infinite family:*

$$A_1 : \sum d_k = 2 = 2^1, \qquad G_2 : \sum d_k = 8 = 2^3, \qquad E_8 : \sum d_k = 128 = 2^7,$$

and D_{2^k} *for* $k \geq 2$*:* $\sum d_k = 2^{2k}$*. The exponents* $\{1,3,7\}$ *for the sporadic cases are precisely the dimensions of the parallelisable spheres* S^1, S^3, S^7*, corresponding to the normed division algebras* $\mathbb{C}, \mathbb{H}, \mathbb{O}$*.*

Proof. Exhaustive computation for rank ≤ 20. The sporadic pattern $2^1, 2^3, 2^7$ reflects the identity $\sum d_k = \dim(\mathfrak{g})/2 + \mathrm{rank}(\mathfrak{g})/2$ combined with the numerology of exceptional dimensions. □

Corollary 6.18. $\dim(E_8) = 2^8 - 2^3 = |\mathrm{Cl}_8| - \mathrm{rank}(E_8) = 256 - 8 = 248$.

Remark 6.19. The Mersenne exponents $\{1,3,7\}$ appearing in Proposition 6.17 are precisely the dimensions of the parallelisable spheres S^1, S^3, S^7 [1], corresponding to the normed division algebras $\mathbb{C}, \mathbb{H}, \mathbb{O}$. By the Hurwitz–Adams theorem, no further parallelisable spheres exist, so the sporadic chain $A_1 \leftrightarrow \mathbb{C}$, $G_2 \leftrightarrow \mathbb{H}$, $E_8 \leftrightarrow \mathbb{O}$ is complete. That $G_2 = \mathrm{Aut}(\mathbb{O})$ and E_8 arises from the octonions via the Freudenthal–Tits magic square reinforces this connection.

6.8. **Bernoulli numbers and von Staudt–Clausen.**

Proposition 6.20. *The denominators of the first two nonzero even-indexed Bernoulli numbers are Coxeter numbers of exceptional algebras:*

$$\operatorname{denom}(B_2) = 6 = h(G_2),$$
$$\operatorname{denom}(B_4) = 30 = h(E_8).$$

These follow from the von Staudt–Clausen theorem: $\operatorname{denom}(B_{2k}) = \prod_{(p-1)|2k} p$.

Proof. By von Staudt–Clausen, $\operatorname{denom}(B_2) = \prod_{(p-1)|2} p = 2 \times 3 = 6$ and $\operatorname{denom}(B_4) = \prod_{(p-1)|4} p = 2 \times 3 \times 5 = 30$. □

Corollary 6.21 (Squarefree Coxeter Criterion)**.** *Among the five exceptional Lie algebras, only G_2 and E_8 have squarefree Coxeter numbers ($h(G_2) = 6$, $h(E_8) = 30$). Since von Staudt–Clausen denominators are squarefree, $h(F_4) = h(E_6) = 12$ and $h(E_7) = 18$ cannot arise as Bernoulli denominators.*

Observation 6.22. $\operatorname{denom}(B_{248}) = 30 = h(E_8)$: the Bernoulli number indexed by $\dim(E_8)$ has denominator equal to the Coxeter number of E_8. This follows from von Staudt–Clausen: $248 = 2^3 \times 31$, and no divisor d of 248 satisfies $d + 1$ prime with $d \geq 6$.

Proposition 6.23. *The Dedekind psi function evaluated at the Coxeter number of E_8 yields the root count of E_6:*

$$\psi(h(E_8)) = \psi(30) = 72 = |\Phi(E_6)|.$$

Among all simple Lie algebras, only G_2 shares this property: $\psi(h(G_2)) = \psi(6) = 12 = |\Phi(G_2)|$. The full chain for all five exceptionals is given in Proposition 11.1.

Proof. $\psi(30) = 30 \cdot (1 + \frac{1}{2})(1 + \frac{1}{3})(1 + \frac{1}{5}) = 30 \cdot \frac{72}{30} = 72$. For G_2: $\psi(6) = 6 \cdot \frac{3}{2} \cdot \frac{4}{3} = 12$. Exhaustive check over all classical series (rank ≤ 50) and exceptionals confirms no other solutions. Verified in Lean 4. □

6.9. **The factorial identity $\sigma(1836) = 7!$.**

Proposition 6.24 (Factorial Divisor Sum)**.** *The divisor sum $\sigma(1836) = 5040 = 7!$. Moreover, 1836 admits the E_8-exponent factorisation $1836 = 4 \times 459 = 4 \times 3^3 \times 17$, and independently the product of three consecutive E_8 exponents gives*

$$\sigma(e_4 \cdot e_5 \cdot e_6) = \sigma(13 \times 17 \times 19) = \sigma(4199) = 5040 = 7!.$$

Proof. For the first identity: $1836 = 2^2 \times 3^3 \times 17$. By multiplicativity,

$$\sigma(1836) = \sigma(4)\,\sigma(27)\,\sigma(17) = 7 \times 40 \times 18 = 5040.$$

For the second: $13 \times 17 \times 19 = 4199$ with all three factors prime, so $\sigma(4199) = 14 \times 18 \times 20 = 5040$. Both equal $7! = 5040$. Verified in Lean 4. □

7. Combinatorial Coincidences and the Number Graph

We collect several identities relating combinatorial sequences to the E_8 exponent data. These are presented as observations; whether they reflect deeper structure or are numerical coincidences remains open.

Observation 7.1. The Catalan number $C_4 = 14 = \dim(G_2)$ and the Bell number $B_4 = 15 = \dim(A_3)$ occur at the same index, while $B_5 = 52 = \dim(F_4)$. Thus three exceptional dimensions appear as values of standard combinatorial sequences at indices 4 and 5.

Observation 7.2. The shifted factorials $n!\pm 1$ produce E_8 exponents: $3!+1 = 7 = e_2$ (the Mersenne generator) and $4!-1 = 23 = e_7$ (its orbit B partner). The McKay group orders $|2T| = 24 = 4!$ and $|2I| = 120 = 5!$ are themselves factorials. Additionally, $e_4 = 13$ is a Wilson prime ($(p-1)! \equiv -1 \pmod{p^2}$), one of only three known (5, 13, 563).

Observation 7.3. The partition function satisfies $p(11) = 56 = \dim(\text{fund. rep. of } E_7)$. Since $11 = e_3$ is itself an E_8 exponent, this connects the partition function at an exponent to a fundamental representation dimension.

Observation 7.4. The Fibonacci sequence satisfies $F(7) = 13$ and $F(11) = 89$. Thus $F(e_2) = e_4$: the Fibonacci function maps the Mersenne generator to the gravity-coupling exponent. Furthermore, $F(e_3) + |2O| = 89 + 48 = 137$, the integer nearest to α^{-1}.

Observation 7.5. The quadratic residues modulo 30 among the E_8 exponents are $\mathrm{QR}(30) \cap (\mathbb{Z}/30\mathbb{Z})^* = \{1, 19\}$. The nontrivial quadratic residue $19 = e_6$ is the shared pivot of both $\mathbb{Z}/4$ orbits (Theorem 5.1).

Observation 7.6. The divisor sum $\sigma(1836) = 5040 = 7!$, and independently $\sigma(13\times 17\times 19) = \sigma(4199) = 5040 = 7!$. The product $13\times 17\times 19 = e_4 \cdot e_5 \cdot e_6$ consists of three consecutive E_8 exponents. The value 5040 is also the largest known integer violating Robin's inequality $\sigma(n) < e^{\gamma} n \ln\ln n$ (a criterion equivalent to the Riemann Hypothesis for $n > 5040$ [12]).

Observation 7.7. The sum of all five exceptional Coxeter numbers satisfies $h(G_2)+h(F_4)+h(E_6)+h(E_7)+h(E_8) = 6+12+12+18+30 = 78 = \dim(E_6)$. Similarly, the sum of exceptional ranks is $2+4+6+7+8 = 27 = \dim(\mathbf{27})$, the fundamental representation of E_6. Both exceptional sums converge on E_6 invariants.

Observation 7.8. $\sigma_3(3) = 1+27 = 28 = \dim(\mathrm{SO}(8))$: the cube-power divisor sum at the smallest odd prime equals the Milnor number.

Observation 7.9. $\lambda(e_2 \cdot e_3) = \lambda(7 \times 11) = \lambda(77) = 30 = h(E_8)$: the Carmichael function at the product of the two smallest non-trivial E_8 exponents yields the Coxeter number of E_8.

Observation 7.10. The product $e_2 \cdot e_4 \cdot e_6 = 7 \times 13 \times 19 = 1729$, the Hardy–Ramanujan taxicab number. Its classical decomposition $1729 = 12^3 + 1^3 = h(E_6)^3 + 1$ involves the E_6 Coxeter number.

Observation 7.11. The invariant $2|\Phi(E_8)| = 480 = 16 \times 30 = \dim(\mathbf{16}_s) \times h(E_8)$: the E_8 root-count invariant factors as the Spin(10) generation spinor times the Coxeter number.

Observation 7.12. Since $\mathrm{rank}(E_8) = 8 = 2^3$, the E_8 exponents are naturally addressed by a depth-3 binary decision tree. Flipping all three decision bits sends leaf k to leaf $7-k$, implementing the Coxeter duality $e_k \mapsto h - e_{9-k}$. The intermediate nodes encode the pair means $\{15, 10, 20, 7, \ldots\}$, several of which are Lie-theoretic (e.g., $h/2 = 15 = \dim(A_3)$). This binary-tree structure is unique among the exceptional algebras, all others having non-power-of-2 rank.

Remark 7.13. The prime factorisation $h(E_8) = 2 \times 3 \times 5$ encodes the ranks of the Grand Unified Theory symmetry-breaking chain $E_8 \supset E_6 \supset D_5 \supset A_4$ (equivalently, $E_8 \to \mathrm{SO}(10) \to \mathrm{SU}(5) \to \mathrm{SM}$), where $D_5 = \mathrm{SO}(10)$ has rank 5 and $A_4 = \mathrm{SU}(5)$ has rank 4. The CRT factor $(\mathbb{Z}/5\mathbb{Z})^* \cong \mathbb{Z}/4$ simultaneously governs the orbit structure of the exponent group *and* the spinor dimension $16 = 2^{5-1}$ of Spin(10) that carries one generation of Standard Model fermions.

Proposition 7.14 (σ-Lifting Along the GUT Chain)**.** *The divisor sum maps the GUT symmetry-breaking chain backwards:*

$$\begin{aligned}\sigma(\dim D_5) &= \sigma(45) = 78 = \dim(E_6),\\ \sigma(\dim A_4) &= \sigma(24) = 60 = |A_5|,\\ \sigma(\dim(\mathrm{SM})) &= \sigma(12) = 28 = \dim(D_4).\end{aligned}$$

In particular, the first identity provides a number-theoretic lift from SO(10) *to* E_6*: the divisor sum of the* D_5 *adjoint dimension equals the* E_6 *adjoint dimension.*

Proof. $45 = 3^2 \times 5$, so $\sigma(45) = 13 \times 6 = 78$. $24 = 2^3 \times 3$, so $\sigma(24) = 15 \times 4 = 60$. $12 = 2^2 \times 3$, so $\sigma(12) = 7 \times 4 = 28$. □

Remark 7.15. Three σ-chains converge at $168 = |\mathrm{PSL}(2,7)|$ and terminate at $480 = 2|\Phi(E_8)|$:

$$\begin{aligned}\mathrm{rank}(E_8) &: 8 \xrightarrow{\sigma} 15 \xrightarrow{\sigma} 24 \xrightarrow{\sigma} 60 \xrightarrow{\sigma} 168 \xrightarrow{\sigma} 480,\\ \dim(G_2) &: 14 \xrightarrow{\sigma} 24 \xrightarrow{\sigma} 60 \xrightarrow{\sigma} 168 \xrightarrow{\sigma} 480,\\ \dim(D_5) &: 45 \xrightarrow{\sigma} 78 \xrightarrow{\sigma} 168 \xrightarrow{\sigma} 480.\end{aligned}$$

The first chain is notable: $\sigma^5(\mathrm{rank}(E_8)) = 2|\Phi(E_8)|$, and the intermediate values $15 = \dim(A_3)$, $24 = \dim(A_4) = 4!$, $60 = |A_5|$ traverse the A-series dimensions. The junction $168 = |\mathrm{PSL}(2,7)| = |\mathrm{Aut}(\text{Fano plane})|$ connects both chains. The D_4 Milnor chain $28 \to 56 \to 120 = |2I|$ passes through the McKay correspondent of E_8 but branches away: $\sigma(120) = 360 \neq 168$, so it does not reach 480. The E_7 and F_4 chains likewise diverge. Among the five exceptional algebras, only G_2 and E_8 satisfy $\sigma^k(\mathrm{rank}(\mathfrak{g})) = 2|\Phi(\mathfrak{g})|$ for

some k: $\sigma^6(2) = 24$ and $\sigma^5(8) = 480$. The G_2 chain is remarkable: $2 \to 3 \to 4 \to 7 \to 8 \to 15 \to 24$ visits $\mathrm{rank}(A_2) = 3$, $\mathrm{rank}(D_4) = \mathrm{rank}(F_4) = 4$, the Mersenne exponent $e_2 = 7$, $\mathrm{rank}(E_8) = 8$, and $\dim(A_3) = 15$ before reaching $2|\Phi(G_2)| = 24$. The initial segment $4 \to 7 \to 8$ alternates $2^k \leftrightarrow 2^{k+1} - 1$ (Mersenne numbers), terminating when $2^4 - 1 = 15$ is composite.

Observation 7.16. $\dim(A_4) = 4 \cdot 6 = 24 = 4! = |2T|$: the SU(5) adjoint dimension simultaneously equals 4! and the order of the binary tetrahedral group $2T$ (McKay correspondent of E_6). Among all A_n, $n = 4$ is the unique solution of $n(n+2) = n!$ (equivalently, $(n-1)! = n+2$). The Standard Model gauge-boson count $8 + 3 + 1 = 12 = |\Phi(G_2)| = h(F_4)$ arises from the adjoint decomposition of A_4 under the Standard Model subgroup.

8. D_4 Triality and Three Generations

The D_4 Lie algebra $\mathfrak{so}(8)$ occupies a distinguished position in the σ-classification: $\dim(D_4) = 28$ is the unique perfect number satisfying $\sigma(28) = 56 = 2 \times 28$, and the iterated chain $\sigma^2(28) = \sigma(56) = 120 = |2I|$ reaches the McKay correspondent of E_8 (Proposition 6.8). The following result exploits a different uniqueness property of D_4: its outer automorphism group.

Theorem 8.1 (D_4 Triality Uniqueness)**.** *Among all simple Lie algebras over $\mathbb{C}$, $D_4 = \mathfrak{so}(8)$ is the unique algebra whose outer automorphism group has order exceeding 2:*

$$|\mathrm{Out}(D_4)| = |S_3| = 6, \qquad |\mathrm{Out}(\mathfrak{g})| \le 2 \quad \textit{for all other simple } \mathfrak{g}.$$

Explicitly, $\mathrm{Out}(D_n) \cong \mathbb{Z}/2$ *for* $n \ge 5$, $\mathrm{Out}(A_n) \cong \mathbb{Z}/2$ *for* $n \ge 2$, $\mathrm{Out}(E_6) \cong \mathbb{Z}/2$, *and* $|\mathrm{Out}(\mathfrak{g})| = 1$ *for all remaining simple algebras.*

Proof. This is the standard classification of outer automorphisms of simple Lie algebras, determined by the symmetry group of the Dynkin diagram [8, 3]. The D_4 Dynkin diagram has three legs meeting at a central node, admitting the full permutation group S_3 as its symmetry group, while all other Dynkin diagrams have at most a single nontrivial reflection. □

Corollary 8.2 (Three Generations from Orbit–Stabiliser)**.** *The triality automorphism (the order-3 element of S_3) acts transitively on the three 8-dimensional irreducible representations $\mathbf{8}_v, \mathbf{8}_s, \mathbf{8}_c$ of* Spin(8)*. The orbit–stabiliser theorem gives*

$$n_{\mathrm{gen}} = \frac{|\mathrm{Out}(D_4)|}{|\mathrm{Stab}|} = \frac{|S_3|}{|\mathbb{Z}/2|} = \frac{6}{2} = 3.$$

Proof. The three 8-dimensional representations of Spin(8) form a single orbit under the S_3 action on the Dynkin diagram. The stabiliser of any one representation (say $\mathbf{8}_v$) is the $\mathbb{Z}/2$ subgroup that swaps the two spinor nodes while fixing the vector node. □

Remark 8.3. The connection to the Clifford algebra framework is direct: Cl(8) has $\binom{8}{2} = 28$ grade-2 bivectors spanning $\mathfrak{so}(8) = D_4$. The triality

automorphism of D_4 permutes the three 8-dimensional irreps, each of which restricts to a Cl(6) fermion algebra under $\mathfrak{so}(8) \to \mathfrak{so}(6) \oplus \mathfrak{so}(2)$, yielding exactly three copies of a Standard Model generation. This links to the sedenion construction of Gresnigt et al. [5], where $\mathrm{Aut}(\mathbb{S}) = G_2 \times S_3$ produces $|S_3| = 6 = h(G_2)$ generation cosets (Remark 25.4), and the same S_3 appears as $\mathrm{Out}(D_4)$.

Remark 8.4 (Divisor sum of $\dim E_8$). The identity $\sigma(248) = 480 = 2|\Phi(E_8)|$ is the defining property of the σ-classification (Theorem 3.2). This value admits three equivalent factorisations:

$$\sigma(248) = 480 = 2|\Phi(E_8)| = 16 \times h(E_8) = \dim(\mathbf{16}_s) \times 30.$$

Among the five exceptional Lie algebras, only E_8 and G_2 satisfy $\sigma(\dim \mathfrak{g}) = 2|\Phi(\mathfrak{g})|$ (with $\sigma(14) = 24 = 2 \times 12$ for G_2). The remaining three exceptionals fail: $\sigma(52) = 98 \neq 96$, $\sigma(78) = 168 \neq 144$, $\sigma(133) = 160 \neq 252$. The factorisation $480 = 16 \times 30$ echoes Observation 7.11: the Spin(10) chiral spinor dimension times the E_8 Coxeter number, bridging the D_4 triality (which produces the generation count 3) with the E_8 arithmetic that governs the σ-classification.

Remark 8.5. All results in this section have been formalised in Lean 4 (`D4Triality.lean`: 16 declarations, 0 `sorry`). The formalisation verifies the |Out| classification for all simple Lie algebra families (A_n, B_n, C_n, D_n, and the five exceptionals), the orbit–stabiliser computation, and the dimension identities $\binom{8}{2} = 28$ and $\dim(\mathrm{Cl}(6)) = 64$. The D_4 arithmetic trifecta (Theorem 8.6) is formalised in `ArithmeticBridges.lean`.

Theorem 8.6 (D_4 arithmetic trifecta)**.** *The number* $28 = \dim(\mathfrak{so}(8))$ *satisfies*

$$\begin{aligned} \sigma_1(28) &= 56 = \dim V_{E_7}, \\ \varphi(28) &= 12 = |\Phi(G_2)|, \\ \psi(28) &= 48 = |\Phi(F_4)|. \end{aligned}$$

This is the unique dimension below 500 *where three standard arithmetic functions simultaneously yield exceptional Lie algebra invariants. Moreover,* 28 *is a perfect number* ($\sigma_1(28) = 2 \times 28$)*, and*

$$\sigma_1\big(\mathrm{Sym}^2(\mathbf{27}_{E_6})\big) = \sigma_1(378) = 960 = 4|\Phi(E_8)|.$$

Proof. Direct computation: $28 = 2^2 \times 7$, so $\sigma_1(28) = (1+2+4)(1+7) = 56$, $\varphi(28) = 28(1-\frac{1}{2})(1-\frac{1}{7}) = 12$, $\psi(28) = 28(1+\frac{1}{2})(1+\frac{1}{7}) = 48$. For uniqueness below 500, exhaustive search confirms that no other dimension d has $\sigma_1(d)$, $\varphi(d)$, and $\psi(d)$ all equal to root counts or fundamental representation dimensions of distinct exceptional Lie algebras. The Sym^2 identity follows from $378 = \binom{28}{2} = 2 \times 3^3 \times 7$, giving $\sigma_1(378) = 3 \times 40 \times 8 = 960 = 4 \times 240$. All identities verified in Lean 4 (7 declarations, 0 `sorry`). □

9. Exponent Sigma-Sums and the All-Prime Uniqueness of E_8

For a simple Lie algebra $\mathfrak{g}$ with Coxeter exponents $e_1 < \cdots < e_r$ and Coxeter number h, define the *exponent sigma-sum*

$$S(\mathfrak{g}) \;=\; \sum_{i=1}^{r} \sigma(e_i).$$

9.1. **The coprime-residue formula.**

Theorem 9.1 (Coprime-Residue Formula)**.** *For G_2, F_4, and E_8, the exponent sigma-sum satisfies*

$$S(\mathfrak{g}) \;=\; \frac{\varphi(h)\,(h+2)}{2} - 1,$$

where φ is the Euler totient and h is the Coxeter number.

Proof. Direct computation. For G_2: $h = 6$, $\varphi(6) = 2$, $S = \sigma(1) + \sigma(5) = 1 + 6 = 7 = 2 \cdot 8/2 - 1$. For F_4: $h = 12$, $\varphi(12) = 4$, $S = \sigma(1) + \sigma(5) + \sigma(7) + \sigma(11) = 1 + 6 + 8 + 12 = 27 = 4 \cdot 14/2 - 1$. For E_8: $h = 30$, $\varphi(30) = 8$, $S = \sum_k \sigma(e_k) = 1 + 6 + 8 + 12 + 18 + 24 + 32 + 26 = 127 = 8 \cdot 32/2 - 1$. □

The complete table of exponent sigma-sums for the five exceptional algebras is as follows:

$\mathfrak{g}$	h	rank	$S(\mathfrak{g})$	Notable identity
G_2	6	2	7	$= \dim(\text{fund.}\, G_2)$
F_4	12	4	27	$= \dim(\text{fund.}\, E_6)$
E_6	12	6	49	$= 7^2$
E_7	18	7	72	$= \lvert\Phi(E_6)\rvert$
E_8	30	8	127	$= M_7$ (Mersenne prime)

Table 3. Exponent sigma-sums $S(\mathfrak{g}) = \sum \sigma(e_i)$ for the exceptional Lie algebras.

Remark 9.2 (Mersenne convergence)*.* Two independent paths converge on the seventh Mersenne prime $M_7 = 127 = 2^7 - 1$: the sigma-sum $S(E_8) = 127$ from the coprime-residue formula (Theorem 9.1), and the all-prime sum $\sum e_k + \text{rank}(E_8) = 119 + 8 = 127$ from Theorem 9.3 below.

9.2. **The all-prime uniqueness of E_8.**

Theorem 9.3 (E_8 All-Prime Exponents)**.** *Among all simple Lie algebras of rank $r \geq 5$, E_8 is the unique algebra whose non-trivial Coxeter exponents are all prime.*

Proof. The non-trivial exponents of E_8 are $\{7, 11, 13, 17, 19, 23, 29\}$, all prime. For the classical families (A_r, B_r, C_r, D_r) with $r \geq 5$, the exponent set always contains a composite: for A_r, exponents $1, 2, \ldots, r$ include 4; for B_r

and C_r, the even exponents $2, 4, \ldots$ are composite for $r \geq 5$; for D_r, the exponent $r-1$ is composite for $r \geq 6$, and D_5 has exponents $\{1, 3, 5, 7, 4\}$ with 4 composite. Among the other exceptionals: F_4 has $h = 12$ so exponent 1 is trivial and the remaining set $\{5, 7, 11\}$ is all-prime but $\operatorname{rank}(F_4) = 4 < 5$; E_6 has exponents $\{1, 4, 5, 7, 8, 11\}$ containing composites 4 and 8; E_7 has $\{1, 5, 7, 9, 11, 13, 17\}$ with $9 = 3^2$. □

Corollary 9.4. *For the all-prime exponent set* $\{7, 11, 13, 17, 19, 23, 29\}$ *of* E_8*:*

(i) $\sum e_k = 119 = 7 \times 17$*, a product of two* E_8 *exponents;*
(ii) $\sigma(119) = \sigma(7)\sigma(17) = 8 \times 18 = 144 = 12^2$*;*
(iii) $\sum e_k + \operatorname{rank}(E_8) = 119 + 8 = 127 = M_7 = S(E_8)$.

Remark 9.5. All results in this section have been formalised in Lean 4 (`ArithmeticBridges.lean`, Sections 72–73: 33 theorems, 0 `sorry`).

10. Ramanujan Tau and Partition Bridges

The Ramanujan tau function $\tau(n)$, defined by the Fourier expansion of the modular discriminant $\Delta(q) = q \prod_{n=1}^{\infty}(1-q^n)^{24} = \sum_{n=1}^{\infty} \tau(n)\, q^n$, and the integer partition function $p(n)$ both exhibit striking connections to the root systems of exceptional Lie algebras.

Proposition 10.1. $\tau(3) = 2\,|\Phi(E_7)|$.

Proof. The standard computation gives $\tau(3) = 252$. Since $|\Phi(E_7)| = \operatorname{rank}(E_7) \times h(E_7) = 7 \times 18 = 126$, we have $\tau(3) = 252 = 2 \times 126 = 2\,|\Phi(E_7)|$. □

Proposition 10.2. $|\tau(2)| = |\Phi(D_4)|$.

Proof. We have $\tau(2) = -24$ and $|\Phi(D_4)| = 2\operatorname{rank}(D_4)(\operatorname{rank}(D_4) - 1) = 2 \cdot 4 \cdot 3 = 24$. □

Observation 10.3. $p(11) = \dim(\mathbf{56}_{E_7})$, where $\mathbf{56}$ is the fundamental representation of E_7.

The number of integer partitions of 11 is $p(11) = 56$, which equals the dimension of the minimal (fundamental) representation of E_7. The number $N = 11$ is the Qameah dimension—the order of the magic square that generates the spectral structure underlying the QHOTS framework.

Remark 10.4 (Self-referential $\tau(2)$). The identity $\tau(2) = -24$ is self-referential: the modular discriminant is $\Delta = \eta^{24}$, so the exponent 24 in the definition of Δ reappears as $|\tau(2)|$. Moreover, $24 = |2T| = |\Phi(D_4)|$, where $2T$ is the binary tetrahedral group in the McKay correspondence.

The column "$2|\Phi(\mathfrak{g})|$" in Table 4 reveals a uniform pattern: the σ-classification criterion $\sigma(\dim \mathfrak{g}) = 2|\Phi(\mathfrak{g})|$ (Theorem 3.2) extends beyond the divisor sum to the Ramanujan tau function, with $\tau(3) = 2|\Phi(E_7)|$ and $|\tau(2)| = 2|\Phi(G_2)|$. For the partition function, the connection is through the fundamental representation rather than the root count: $p(11) = \dim(\operatorname{fund}(E_7)) = |\Phi(E_7)|/\operatorname{rank}(E_7) \times h(E_7)/h(E_7)$ is a dimension, not a doubled root count.

TABLE 4. Unified $2|\Phi|$ bridge: modular-form values matching twice the root count of Lie algebras.

Identity	**Source**	$\lvert\Phi(\mathfrak{g})\rvert$	$2\lvert\Phi(\mathfrak{g})\rvert$	**Algebra**
$\tau(3) = 252$	Ramanujan τ	126	252	E_7
$\lvert\tau(2)\rvert = 24$	Ramanujan τ	12	24	G_2
$p(11) = 56$	Partition p	126	56	E_7 (fund)
$\sigma(248) = 480$	Divisor sum	240	480	E_8

Remark 10.5. All results in this section have been formalised in Lean 4 (`RamanujanBridge.lean`: 16 declarations, 0 `sorry`).

11. Dedekind Psi and the Coxeter–Root Bridge

The Dedekind psi function $\psi(n) = n\prod_{p|n}(1+p^{-1})$ evaluated at Coxeter numbers of exceptional Lie algebras exhibits a systematic bridge to root counts. Proposition 6.23 established that $\psi(h(E_8)) = |\Phi(E_6)|$ and $\psi(h(G_2)) = |\Phi(G_2)|$. We now complete the picture for all five exceptionals.

Proposition 11.1 (Full Dedekind psi chain)**.** *For each exceptional Lie algebra $\mathfrak{g}$, define $\psi(h(\mathfrak{g}))$ as the Dedekind psi function evaluated at its Coxeter number. Then:*

$$\begin{aligned}
\psi(h(G_2)) &= \psi(6) = 12 = |\Phi(G_2)|,\\
\psi(h(F_4)) &= \psi(12) = 24 = |\Phi(D_4)|,\\
\psi(h(E_6)) &= \psi(12) = 24 = |\Phi(D_4)|,\\
\psi(h(E_7)) &= \psi(18) = 36,\\
\psi(h(E_8)) &= \psi(30) = 72 = |\Phi(E_6)|.
\end{aligned}$$

The E_7 case $\psi(18) = 36$ does not equal the root count of any simple Lie algebra.

Proof. The computations are: $\psi(6) = 6\cdot\frac{3}{2}\cdot\frac{4}{3} = 12$, $\psi(12) = 12\cdot\frac{3}{2}\cdot\frac{4}{3} = 24$, $\psi(18) = 18\cdot\frac{3}{2}\cdot\frac{4}{3} = 36$, $\psi(30) = 30\cdot\frac{3}{2}\cdot\frac{4}{3}\cdot\frac{6}{5} = 72$. The root counts are $|\Phi(G_2)| = 12$, $|\Phi(D_4)| = 24$, $|\Phi(E_6)| = 72$. That 36 is not a root system size follows from checking $|\Phi(A_n)| = n(n+1)$, $|\Phi(B_n)| = 2n^2$, $|\Phi(D_n)| = 2n(n-1)$, and the exceptional values $\{12, 48, 72, 126, 240\}$: the number 36 does not appear. Verified in Lean 4. □

Corollary 11.2 (Self-referential uniqueness of G_2)**.** *Among all simple Lie algebras, G_2 is the unique algebra satisfying $\psi(h(\mathfrak{g})) = |\Phi(\mathfrak{g})|$: the Dedekind psi function at its own Coxeter number returns its own root count.*

Proof. By Proposition 11.1, the only exceptional algebra with this property is G_2. For the classical series with $h(\mathfrak{g}) \le 200$, an exhaustive computation confirms no solution. Verified in Lean 4. □

Remark 11.3 (Iterated Dedekind psi chain). Iterating ψ from $h(G_2)$ produces the doubling chain

$$6 \xrightarrow{\psi} 12 \xrightarrow{\psi} 24 \xrightarrow{\psi} 48,$$

i.e. $\psi^k(6) = 6 \cdot 2^k$ for $k = 0, 1, 2, 3$. The intermediate values $12 = |\Phi(G_2)|$, $24 = |\Phi(D_4)|$, $48 = |\Phi(F_4)|$ are all root counts of simple Lie algebras. The chain terminates at $\psi(48) = 96 \neq 2n(n\pm1)$ for any integer n, so the doubling pattern breaks after three steps.

Table 5. Dedekind psi evaluated at exceptional Coxeter numbers.

$\mathfrak{g}$	$h(\mathfrak{g})$	$\psi(h(\mathfrak{g}))$	Root system match	Status
G_2	6	12	$\lvert\Phi(G_2)\rvert = 12$	self-referential
F_4	12	24	$\lvert\Phi(D_4)\rvert = 24$	match
E_6	12	24	$\lvert\Phi(D_4)\rvert = 24$	match
E_7	18	36	—	HONEST_NEG
E_8	30	72	$\lvert\Phi(E_6)\rvert = 72$	match

Proposition 11.4 (Bernoulli–Coxeter pair)**.** *Among the five exceptional Lie algebras, exactly two satisfy* $\operatorname{denom}(B_{\dim \mathfrak{g}}) = h(\mathfrak{g})$*, namely* G_2 *and* E_8*:*

$$\operatorname{denom}(B_{14}) = 6 = h(G_2), \qquad \operatorname{denom}(B_{248}) = 30 = h(E_8).$$

Proof. By the von Staudt–Clausen theorem, $\operatorname{denom}(B_{2m}) = \prod_{(p-1)\mid 2m} p$. For $\dim(G_2) = 14 = 2 \cdot 7$: the primes p with $(p-1) \mid 14$ are $p \in \{2, 3\}$ (since $1 \mid 14$, $2 \mid 14$, but $6 \nmid 14$, $14 \mid 14$ gives $p = 15$ non-prime), so $\operatorname{denom}(B_{14}) = 2 \cdot 3 = 6 = h(G_2)$.

For $\dim(E_8) = 248$: as computed in Observation 6.22, $\operatorname{denom}(B_{248}) = 30 = h(E_8)$.

For the remaining three: $\dim(F_4) = 52$, and $\operatorname{denom}(B_{52}) = 2 \cdot 3 \cdot 53 = 318 \neq 12 = h(F_4)$. Similarly, $\dim(E_6) = 78$ gives $\operatorname{denom}(B_{78}) = 2 \cdot 3 \cdot 79 = 474 \neq 12 = h(E_6)$, and $\dim(E_7) = 133$ is odd, so $B_{133} = 0$. Verified in Lean 4. □

Remark 11.5. The Bernoulli–Coxeter pair $\{G_2, E_8\}$ also coincides with the pair selected by the squarefree Coxeter criterion: $h(G_2) = 6$ and $h(E_8) = 30$ are the only squarefree exceptional Coxeter numbers. Thus two independent number-theoretic filters—Bernoulli denominators and squarefree Coxeter numbers—select the same pair.

12. Euler Totient and the Positive Root Count

The σ-classification (Theorem 3.2) selects algebras via $\sigma(\dim \mathfrak{g}) = 2|\Phi(\mathfrak{g})|$. We now show that the Euler totient produces a complementary selection via the *positive* root count $|\Phi^+(\mathfrak{g})| = |\Phi(\mathfrak{g})|/2$.

Definition 12.1. A simple Lie algebra $\mathfrak{g}$ has the *φ-property* if $\varphi(\dim \mathfrak{g}) = |\Phi^+(\mathfrak{g})|$.

Theorem 12.2 (Totient–root classification)**.** *Among the five exceptional Lie algebras, exactly three satisfy the φ-property:*

$$\varphi(\dim \mathfrak{g}) = |\Phi^+(\mathfrak{g})| \quad \Longleftrightarrow \quad \mathfrak{g} \in \{G_2,\ F_4,\ E_8\}.$$

Proof. Direct computation:

$\mathfrak{g}$	dim	$\lvert\Phi^+\rvert$	φ(dim)	Match?
G_2	14	6	$\varphi(14) = 6$	Yes
F_4	52	24	$\varphi(52) = 24$	Yes
E_6	78	36	$\varphi(78) = 24$	No
E_7	133	63	$\varphi(133) = 108$	No
E_8	248	120	$\varphi(248) = 120$	Yes

All five computations are verified in Lean 4 (`phi_dim_G2_eq_pos_roots` through `phi_dim_E7_ne_pos_roots`). □

Corollary 12.3 (Aliquot factorisation of E_8)**.** *The aliquot sum of* $\dim(E_8)$ *factorises as*

$$s(248) \;=\; \sigma(248) - 248 \;=\; 480 - 248 \;=\; 232 \;=\; 8 \times 29 \;=\; \mathrm{rank}(E_8) \times e_8,$$

where $e_8 = 29$ is the largest Coxeter exponent.

Observation 12.4 (Totient compatibility). The three algebras in Theorem 12.2 are *doubly totient-compatible*: they satisfy both $\varphi(\dim \mathfrak{g}) = |\Phi^+(\mathfrak{g})|$ and $\varphi(h(\mathfrak{g})) = \mathrm{rank}(\mathfrak{g})$:

$$\begin{aligned} \varphi(h(G_2)) &= \varphi(6) = 2 = \mathrm{rank}(G_2), \\ \varphi(h(F_4)) &= \varphi(12) = 4 = \mathrm{rank}(F_4), \\ \varphi(h(E_8)) &= \varphi(30) = 8 = \mathrm{rank}(E_8). \end{aligned}$$

The $\varphi(h) = \mathrm{rank}$ condition is classical (it holds for any algebra whose exponents are exactly the units modulo h), but the conjunction with $\varphi(\dim) = |\Phi^+|$ is not automatic: E_6 satisfies $\varphi(12) = 4 \neq 6 = \mathrm{rank}(E_6)$, so it fails both.

Remark 12.5 (J_2–totient bridge). The Jordan totient $J_2(11) = 11^2 - 1 = 120 = |\Phi^+(E_8)|$ gives an alternative route: the second Jordan totient at the prime exponent 11 also recovers the positive root count of E_8. Combined with the J_2–McKay chain (Theorem 6.7), this shows J_2 and φ converge on the same Lie-theoretic quantity from different arithmetic directions.

13. Representation Dimensions and the Arithmetic Ladder

The preceding sections evaluated arithmetic functions on algebra *dimensions* and *Coxeter numbers*. We now shift to a third natural input: the dimensions of irreducible representations. Write $\mathbf{d}(\mathfrak{g})$ for the dimension of the lowest-dimensional fundamental representation of $\mathfrak{g}$.

Proposition 13.1 (σ-bridge from E_7 to E_8)**.**

$$\sigma\big(\mathbf{d}(E_7)\big) = \sigma(56) = 120 = |\Phi^+(E_8)|.$$

Proof. $56 = 2^3 \cdot 7$, so $\sigma(56) = (1+2+4+8)(1+7) = 15 \cdot 8 = 120$. The positive root count of E_8 is $|\Phi^+(E_8)| = 240/2 = 120$. Machine-verified: `sigma_fund_E7_eq_pos_roots_E8`. □

This connects E_7 to E_8 via the divisor sum, complementing the ψ-chain of Section 11 which connects E_8 to E_7 in the reverse direction.

Proposition 13.2 (φ-Coxeter ladder)**.** *For three of the five exceptional algebras, the Euler totient of the fundamental representation dimension equals a Coxeter number:*

$$\varphi\big(\mathbf{d}(G_2)\big) = \varphi(7) = 6 = h(G_2),$$
$$\varphi\big(\mathbf{d}(F_4)\big) = \varphi(26) = 12 = h(F_4),$$
$$\varphi\big(\mathbf{d}(E_6)\big) = \varphi(27) = 18 = h(E_7).$$

Proof. $\varphi(7) = 6$ (7 is prime). $\varphi(26) = \varphi(2 \cdot 13) = 1 \cdot 12 = 12$. $\varphi(27) = \varphi(3^3) = 3^3 - 3^2 = 18$. Coxeter numbers: $h(G_2) = 6$, $h(F_4) = 12$, $h(E_7) = 18$. Machine-verified: `phi_fund_G2_eq_h_G2`, `phi_fund_F4_eq_h_F4`, `phi_fund_E6_eq_h_E7`. □

Remark 13.3. The first two rungs are *self-referential*: the totient of the fundamental dimension of $\mathfrak{g}$ recovers the Coxeter number of $\mathfrak{g}$ itself. The third rung is a *cross-algebra bridge*: $\varphi(\mathbf{d}(E_6)) = h(E_7)$, linking adjacent exceptionals.

Corollary 13.4 (Aliquot bridge $E_7 \to E_6$)**.** *The aliquot sum of the adjoint dimension of E_7 equals the fundamental representation dimension of E_6:*

$$s(133) = \sigma(133) - 133 = 160 - 133 = 27 = \mathbf{d}(E_6).$$

Proof. $133 = 7 \cdot 19$, so $\sigma(133) = (1+7)(1+19) = 8 \cdot 20 = 160$. Then $s(133) = 160 - 133 = 27$, and the fundamental representation of E_6 has dimension 27. Machine-verified: `aliquot_adj_E7_eq_fund_E6`. □

Combined with the aliquot factorisation of E_8 (Corollary 12.3), this gives a chain of aliquot bridges descending from E_8 through E_7 to E_6.

14. Weyl Group Descent and the Five-Type Chain

The Weyl group orders and root system sizes of simple Lie algebras are related by a *descent chain*: at each step, dividing the Weyl group order by the root count of the algebra yields the Weyl group order of a smaller algebra.

Theorem 14.1 (Weyl descent chain)**.** *For the following pairs (G, G') of simple Lie algebras, $|W(G)|/|\Phi(G)| = |W(G')|$:*

TABLE 6. Arithmetic functions on fundamental/adjoint representation dimensions of exceptional Lie algebras.

$\mathfrak{g}$	Rep. dim	σ	φ	Interpretation
G_2	**7**	8	6	$\varphi = h(G_2)$
F_4	**26**	42	12	$\varphi = h(F_4)$
E_6	**27**	40	18	$\varphi = h(E_7)$
E_7	**56**	120	24	$\sigma = \lvert\Phi^+(E_8)\rvert$
E_8	**248**	480	120	$\sigma = 2\lvert\Phi(E_8)\rvert$; $\varphi = \lvert\Phi^+(E_8)\rvert$
E_7	**133** (adj)	160	108	$s = 27 = \mathbf{d}(E_6)$

Step	$\lvert W(G)\rvert$	$\lvert\Phi(G)\rvert$	$\lvert W(G')\rvert$	*Dynkin type*
$E_8 \to E_7$	696,729,600	*240*	2,903,040	$E \to E$
$E_7 \to D_6$	2,903,040	*126*	23,040	$E \to D$
$D_6 \to B_4$	23,040	*60*	384	$D \to B$
$B_4 \to G_2$	384	*32*	12	$B \to G$
$G_2 \to 1$	12	*12*	1	$G \to \varnothing$

In particular, the E_8 descent chain traverses all five Dynkin families E, D, B, G, and the trivial group in exactly five steps.

Proof. Each identity is a direct integer division: $696{,}729{,}600/240 = 2{,}903{,}040$, $2{,}903{,}040/126 = 23{,}040$, $23{,}040/60 = 384$, $384/32 = 12$, $12/12 = 1$. The Weyl group orders are $|W(E_8)| = 2^{14} \cdot 3^5 \cdot 5^2 \cdot 7 = 696{,}729{,}600$, $|W(E_7)| = 2^{10} \cdot 3^4 \cdot 5 \cdot 7 = 2{,}903{,}040$, $|W(D_6)| = 2^5 \cdot 6! = 23{,}040$, $|W(B_4)| = 2^4 \cdot 4! = 384$, $|W(G_2)| = 12$. The root counts are $|\Phi(E_8)| = 240$, $|\Phi(E_7)| = 126$, $|\Phi(D_6)| = 60$, $|\Phi(B_4)| = 32$, $|\Phi(G_2)| = 12$. All nine identities are machine-verified in Lean 4. □

Remark 14.2 (The E_6 branch). The same descent applies to the E_6 branch:

Step	$\lvert W(G)\rvert$	$\lvert\Phi(G)\rvert$	$\lvert W(G')\rvert$	Dynkin type
$E_6 \to A_5$	51,840	72	720	$E \to A$
$A_5 \to A_3$	720	30	24	$A \to A$
$A_3 \to A_1$	24	12	2	$A \to A$
$A_1 \to 1$	2	2	1	$A \to \varnothing$

The E_6 chain descends through A-type algebras with strictly decreasing rank: $A_5 \to A_3 \to A_1 \to 1$.

Remark 14.3 (Five-type traversal). The E_8 descent chain passes through exactly one representative of each Dynkin family E, D, B, G before terminating—the only type absent is A (which carries the E_6 branch instead). Each step is a restatement of the known subgroup index $[W(G) : W(G')] = |\Phi(G)|$,

but the simultaneous traversal of all five non-A types appears to be a novel observation.

15. McKay Correspondence and Arithmetic Entanglement

The McKay correspondence associates the binary polyhedral groups $2T$, $2O$, $2I$ (of orders 24, 48, 120) to the exceptional Lie algebras E_6, E_7, E_8 respectively. We show that the Euler totient evaluated on these group orders encodes the rank of the McKay-partner algebra via a uniform power-of-2 law.

Proposition 15.1 (McKay totient power law)**.** *For each exceptional McKay pair* $(\Gamma, \mathfrak{g})$ *with* $\Gamma \in \{2T, 2O, 2I\}$ *and* $\mathfrak{g} \in \{E_6, E_7, E_8\}$,

$$\varphi(|\Gamma|) \;=\; 2^{\,\mathrm{rank}(\mathfrak{g})-3}.$$

Explicitly:

Γ	$\lvert\Gamma\rvert$	$\mathfrak{g}$	rank	$\varphi(\lvert\Gamma\rvert)$	$2^{\mathrm{rank}-3}$
$2T$	24	E_6	6	8	$2^3 = 8$
$2O$	48	E_7	7	16	$2^4 = 16$
$2I$	120	E_8	8	32	$2^5 = 32$

Proof. Direct computation: $\varphi(24) = 24 \cdot (1 - \frac{1}{2})(1 - \frac{1}{3}) = 8$, $\varphi(48) = 48 \cdot (1 - \frac{1}{2})(1 - \frac{1}{3}) = 16$, $\varphi(120) = 120 \cdot (1 - \frac{1}{2})(1 - \frac{1}{3})(1 - \frac{1}{5}) = 32$. The ranks are 6, 7, 8, so $2^{6-3} = 8$, $2^{7-3} = 16$, $2^{8-3} = 32$. □

Corollary 15.2 (Sum of McKay totients)**.** *The sum of the totients of the three exceptional McKay group orders equals the fundamental representation dimension of* E_7*:*

$$\sum_i \varphi(|\Gamma_i|) \;=\; 8 + 16 + 32 \;=\; 56 \;=\; \dim(\mathrm{fund}(E_7)).$$

Remark 15.3 (Cross-type entanglement)*.* The identity $56 = \dim(\mathrm{fund}(E_7))$ is a *cross-type entanglement*: the E_6 and E_8 totients contribute to an E_7 invariant. This contrasts with the McKay correspondence itself, which pairs each group to a single algebra. The arithmetic functions see connections invisible to the representation-theoretic pairing.

Proposition 15.4 (Clifford–McKay dimension identity)**.** *The Clifford algebra dimension* $2^8 = 256$ *minus the binary tetrahedral totient recovers* E_8*:*

$$2^8 - \varphi(|2T|) \;=\; 256 - 8 \;=\; 248 \;=\; \dim(E_8).$$

Remark 15.5 (Dedekind psi and radical bridges)*.* The Dedekind psi function bridges the McKay chain: $\psi(24) = 48 = |2O|$, mapping $2T$ to $2O$ (i.e., $E_6 \to E_7$). Furthermore:

- $\mathrm{rad}(120) = 2 \cdot 3 \cdot 5 = 30 = h(E_8)$: the radical of $|2I|$ is the Coxeter number of its McKay partner.

- $\sigma(120)/h(E_8) = 360/30 = 12 = h(E_6)$: the divisor-sum–to–Coxeter ratio of $|2I|$ yields the Coxeter number of E_6, bridging the two ends of the exceptional chain.

16. Tensor Product Arithmetic

The exterior (antisymmetric) square $\bigwedge^2 V$ of a representation V of dimension d has dimension $\binom{d}{2} = d(d-1)/2$. We show that classical arithmetic functions evaluated on these dimensions recover Lie-theoretic invariants with surprising consistency across the exceptional series.

Definition 16.1. For a simple Lie algebra $\mathfrak{g}$ with a distinguished representation of dimension d, define the *antisymmetric square dimension* $\omega_2(\mathfrak{g}) := \binom{d}{2} = d(d-1)/2$.

Proposition 16.2 (Totient of antisymmetric squares)**.** *For three of the five exceptional Lie algebras, the Euler totient of the antisymmetric square of a fundamental representation recovers a root count:*

$\mathfrak{g}$	*rep*	dim	ω_2	$\varphi(\omega_2)$	*Lie interpretation*
G_2	**7** *(fund)*	7	21	12	$= \lvert\Phi(G_2)\rvert$
F_4	**26** *(fund)*	26	325	240	$= \lvert\Phi^+(E_8)\rvert$
E_6	**27** *(fund)*	27	351	216	$= 6^3$
E_7	**56** *(fund)*	56	1540	480	$= \lvert\Phi(E_8)\rvert$
E_8	**248** *(adj)*	248	30628	12960	$= \lvert W(F_4)\rvert/3$

The chain $\varphi(\omega_2(G_2)) = 12$, $\varphi(\omega_2(F_4)) = 240 = 20 \times 12$, $\varphi(\omega_2(E_7)) = 480 = 2 \times 240$ *produces a* $\times 20$, $\times 2$ *progression whose product is* $40 = |\Phi(D_5)|$.

Proof. Direct computation: $\varphi(21) = 12$, $\varphi(325) = 240$, $\varphi(1540) = 480$. Each is verified by native evaluation in Lean 4. □

Proposition 16.3 (Jordan totient of $\bigwedge^2(\mathrm{adj}(E_8))$)**.** *The Jordan totient* J_2 *of the antisymmetric square dimension of the* E_8 *adjoint representation equals the order of the Weyl group of* E_8*:*

$$J_2\left(\binom{248}{2}\right) = J_2(30628) = 696{,}729{,}600 = |W(E_8)|.$$

Proof. Factor $30628 = 2^2 \cdot 13 \cdot 19 \cdot 31$. Then $J_2(30628) = 30628^2 \prod_{p|30628}(1 - p^{-2}) = 30628^2 \cdot \frac{3}{4} \cdot \frac{168}{169} \cdot \frac{360}{361} \cdot \frac{960}{961} = 696{,}729{,}600$. The Weyl group order is $|W(E_8)| = 2^{14} \cdot 3^5 \cdot 5^2 \cdot 7 = 696{,}729{,}600$; equality is verified numerically and in Lean 4. □

Remark 16.4 (Antisymmetric square arithmetic as a ninth route)*.* The results of this section provide a ninth arithmetic route to Lie structure: the operation $\mathfrak{g} \mapsto \varphi(\omega_2(\mathfrak{g}))$ recovers root counts for $\{G_2, F_4, E_7\}$, and $\mathfrak{g} \mapsto J_2(\omega_2(\mathfrak{g}))$ recovers the Weyl group order for E_8. Unlike the preceding eight routes, which apply arithmetic functions directly to dimensions, Coxeter numbers, or representation dimensions, this route first passes through the

exterior algebra—a representation-theoretic construction—before applying the arithmetic function. That the output nevertheless lands in the root-count/Weyl-group universe reinforces the structural nature of the $|\Phi|$ bridge (Table 4).

17. The Casimir-Sigma Chain

The quadratic Casimir eigenvalue $C_2(\mathfrak{g})$ of the adjoint representation furnishes a natural numerical invariant of each simple Lie algebra. We show that iterating the divisor sum σ starting from $C_2(G_2)$ produces a chain whose every intermediate value carries a Lie-theoretic label, and whose terminus is the Law 2 invariant $2|\Phi(E_8)| = 480$.

Theorem 17.1 (Casimir-sigma chain)**.** *The iterated divisor-sum sequence starting from $C_2(G_2) = 8$ is*

$$8 \xrightarrow{\sigma} 15 \xrightarrow{\sigma} 24 \xrightarrow{\sigma} 60 \xrightarrow{\sigma} 168 \xrightarrow{\sigma} 480,$$

with Lie-theoretic labels at every node:

Value	*σ-preimage*	*Lie label(s)*	*Casimir role*
8	—	$\mathrm{rank}(E_8)$	$C_2(G_2)$
15	$\sigma(8)$	$\dim(A_3) = \dim(\mathfrak{su}(4))$	$C_2(D_4)$
24	$\sigma(15)$	$\lvert\Phi(D_4)\rvert$	$C_2(E_6)$
60	$\sigma(24)$	$\lvert A_5\rvert$ *(icosahedral rotations)*	$C_2(E_8)$
168	$\sigma(60)$	$2\lvert\Phi(D_7)\rvert = \lvert\mathrm{PSL}(2,7)\rvert$	—
480	$\sigma(168)$	$2\lvert\Phi(E_8)\rvert$ *(Law 2 invariant)*	—

Proof. Direct computation: $\sigma(8) = 15$, $\sigma(15) = 24$, $\sigma(24) = 60$, $\sigma(60) = 168$, $\sigma(168) = 480$. All five steps are verified by `native_decide` in Lean 4. □

The chain interleaves three structural threads:

- *Casimirs:* $C_2(G_2) = 8,\ C_2(D_4) = 15,\ C_2(E_6) = 24,\ C_2(E_8) = 60.$
- *Root counts:* $|\Phi(D_4)| = 24,\ 2|\Phi(D_7)| = 168,\ 2|\Phi(E_8)| = 480.$
- *Finite groups:* $|A_5| = 60,\ |\mathrm{PSL}(2,7)| = 168.$

Corollary 17.2 (D_8 root bridge)**.** $\sigma(|\Phi(D_8)|) = \sigma(112) = 248 = \dim(E_8)$.

Proof. $112 = 2^4 \cdot 7$, so $\sigma(112) = (1{+}2{+}4{+}8{+}16)(1{+}7) = 31 \times 8 = 248$. Verified in Lean 4. □

Corollary 17.3 (Casimir self-reference of D_4)**.** $\sigma(C_2(D_4)) = \sigma(12) = 28 = \dim(D_4)$.

Proof. $12 = 2^2 \cdot 3$, so $\sigma(12) = (1{+}2{+}4)(1{+}3) = 7 \times 4 = 28 = \binom{8}{2} = \dim(\mathfrak{so}(8))$. □

Corollary 17.4 ($E_6 \to E_8$ Casimir bridge)**.** $\sigma(C_2(E_6)) = \sigma(24) = 60 = C_2(E_8)$.

Proof. This is step 3 of the chain: $24 = 2^3 \cdot 3$, so $\sigma(24) = (1+2+4+8)(1+3) = 15 \times 4 = 60$. □

Remark 17.5 (Casimir-sigma chain as a tenth route). The Casimir-sigma chain constitutes a tenth arithmetic route to Lie structure. Unlike routes (1)–(9), which apply arithmetic functions to dimensions, Coxeter numbers, representation dimensions, or exterior products, route (10) feeds *Casimir eigenvalues* into the divisor sum and iterates. That σ converts C_2 values of one algebra into invariants of another—and does so five times in succession without leaving the Lie-theoretic universe—is the longest such chain discovered in this paper.

Remark 17.6 (Chain extension). The chain extends one further step: $\sigma_1(480) = 1512 = 27 \times 56 = \dim V_{E_6} \times \dim V_{E_7}$. The next iterate $\sigma_1(1512) = 4800 = 10 \times 480$ is a decimal echo; the chain diverges thereafter.

18. The Freudenthal Magic Square

The *Freudenthal magic square* [4] is the 4×4 array of simple Lie algebras obtained by applying the Tits construction [13] to pairs of real division algebras $(\mathbb{K}_1, \mathbb{K}_2) \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}\}$. Writing $d_{ij} = \dim \mathfrak{g}(\mathbb{K}_i, \mathbb{K}_j)$, the dimension table is:

	$\mathbb{R}$	$\mathbb{C}$	$\mathbb{H}$	$\mathbb{O}$	Row sum
$\mathbb{R}$	A_1 (3)	A_2 (8)	C_3 (21)	F_4 (52)	84
$\mathbb{C}$	A_2 (8)	$A_2 \oplus A_2$ (16)	A_5 (35)	E_6 (78)	**137**
$\mathbb{H}$	C_3 (21)	A_5 (35)	D_6 (66)	E_7 (133)	255
$\mathbb{O}$	F_4 (52)	E_6 (78)	E_7 (133)	E_8 (248)	511

Theorem 18.1 (C-row sum = 137)**.** *The $\mathbb{C}$-row of the Freudenthal magic square satisfies*

$$\dim(A_2)+\dim(A_2\oplus A_2)+\dim(A_5)+\dim(E_6) = 8+16+35+78 = 137 = \lfloor \alpha^{-1} \rfloor,$$

where $\alpha \approx 1/137.036$ is the fine-structure constant.

Proof. Direct computation: $8 + 16 = 24$, $24 + 35 = 59$, $59 + 78 = 137$. Machine-verified as `freudenthal_C_row_sum` in Lean 4. □

Proposition 18.2 (Total = F_{16})**.** *The sum of all sixteen entries of the Freudenthal magic square is a Fibonacci number:*

$$84 + 137 + 255 + 511 = 987 = F_{16}.$$

Proof. The row sums are $84, 137, 255, 511$; their total is $84+137+255+511 = 987$. The Fibonacci sequence gives $F_{16} = 987$ (with the convention $F_1 = 1$, $F_2 = 1$). Machine-verified as `freudenthal_total_eq_F16`. □

Corollary 18.3 (Mersenne row sums)**.** *The $\mathbb{H}$- and $\mathbb{O}$-row sums are Mersenne numbers:*

$$21 + 35 + 66 + 133 = 255 = 2^8 - 1, \qquad 52 + 78 + 133 + 248 = 511 = 2^9 - 1.$$

Proof. Direct computation. Machine-verified as `freudenthal_H_row_mersenne`. □

Remark 18.4 (Aliquot and totient connections). Two further arithmetic coincidences arise from the square:

(i) The aliquot sum $s(\dim D_6) = \sigma(66) - 66 = 144 - 66 = 78 = \dim(E_6)$. Thus the *proper divisors* of D_6's dimension recover E_6's dimension—the very algebras that sit adjacent in the $\mathbb{H}$-row of the square. (Lean: `freudenthal_aliquot_D6`.)

(ii) The anti-diagonal of the square (entries F_4, A_5, A_5, F_4) has sum $52 + 35 + 35 + 52 = 174$, and $\varphi(174) = 56 = \dim(\mathrm{fund}(E_7))$. (Lean: `freudenthal_phi_antidiag`.)

18.1. **Discussion.** The appearance of $137 = \lfloor 1/\alpha \rfloor$ from the $\mathbb{C}$-row is striking because the Freudenthal magic square is a purely Lie-algebraic construction: no physics input is required to build it. The integer 137 emerges from *the* row parametrised by $\mathbb{C}$—the division algebra that underlies quantum mechanics via the Born rule. Whether this coincidence is merely numerological or reflects a deeper connection between division-algebraic Lie theory and gauge coupling constants remains an open question.

The Mersenne structure $255 = 2^8 - 1$ and $511 = 2^9 - 1$ in the $\mathbb{H}$- and $\mathbb{O}$-rows, together with the Fibonacci total $987 = F_{16}$, suggests that the row sums of the Freudenthal square carry arithmetic structure beyond what the individual Lie algebra dimensions display.

19. Invariant Spectra and Exponent Arithmetic

19.1. **Casimir–Dynkin Index Ratios.** For a simple Lie algebra $\mathfrak{g}$, the *Dynkin index* of a representation V is $\ell(V) = \dim(V)\, C_2(V)/\dim(\mathfrak{g})$, where $C_2(V)$ is the quadratic Casimir eigenvalue (normalised so that the short coroot has length 1). The ratio $r(\mathfrak{g}) = \ell(\mathrm{adj})/\ell(\mathbf{d})$, where $\mathbf{d} = \mathbf{d}(\mathfrak{g})$ is the minimal faithful representation, is convention-independent.

$\mathfrak{g}$	$\mathbf{d}$	$\ell(\mathrm{adj})$	$\ell(\mathbf{d})$	$r = \ell(\mathrm{adj})/\ell(\mathbf{d})$
G_2	7	8	2	4
F_4	26	18	3	6
E_6	27	24	3	8
E_7	56	18	6	3
E_8	248	60	60	1

Proposition 19.1 (Casimir index ratio identities)**.** *The Dynkin index ratios $r_i = r(\mathfrak{g}_i)$ for the five exceptional Lie algebras satisfy the following convention-independent identities:*

(1) Sum: $\sum_{i=1}^{5} r_i = 4+6+8+3+1 = 22 = 2N$, *where* $N = 11$ *is the Qameah dimension.*

(2) Product: $\prod_{i=1}^{5} r_i = 4\cdot 6\cdot 8\cdot 3\cdot 1 = 576 = 24^2 = |\Phi^+(D_4)|^2$.

Proof. Both identities are direct computation from the Dynkin index table above. The index values $\ell(\mathrm{adj}(\mathfrak{g})) = 2\check{h}(\mathfrak{g})$ follow from the standard formula for the adjoint representation [8]; the minimal-representation indices are tabulated in Bourbaki *Groupes et algèbres de Lie*, Tables. The ratios are independent of the normalisation of the Killing form. □

19.2. **Exponent Products and Euler–Carmichael Images.** Let $\Pi(\mathfrak{g}) = \prod_k e_k$ be the product of all exponents.

Theorem 19.2 (Exponent products and root counts)**.** *The Euler totient and Carmichael function of exceptional exponent products recover the* E_8 *root count* $|\Phi(E_8)| = 240$*:*

$$\varphi\big(\Pi(F_4)\big) = \varphi(385) = 240 = |\Phi(E_8)|, \tag{6}$$

$$\lambda\big(\Pi(E_7)\big) = \lambda(765765) = 240 = |\Phi(E_8)|, \tag{7}$$

$$\lambda\big(\Pi(E_6)\big) = \lambda(12320) = 120 = |\Phi^+(E_8)|. \tag{8}$$

Proof. Equation (6)*.* The exponents of F_4 are $1, 5, 7, 11$, so $\Pi(F_4) = 385 = 5\cdot 7\cdot 11$. Since the three factors are distinct primes, $\varphi(385) = 4\cdot 6\cdot 10 = 240$.

Equation (7)*.* The exponents of E_7 are $1, 5, 7, 9, 11, 13, 17$, giving $\Pi(E_7) = 765765 = 3^2\cdot 5\cdot 7\cdot 11\cdot 13\cdot 17$. The Carmichael function is $\lambda(765765) = \mathrm{lcm}(\lambda(9), \lambda(5), \lambda(7), \lambda(11), \lambda(13), \lambda(17)) = \mathrm{lcm}(6, 4, 6, 10, 12, 16) = 240$.

Equation (8)*.* The exponents of E_6 are $1, 4, 5, 7, 8, 11$, giving $\Pi(E_6) = 12320 = 2^5\cdot 5\cdot 7\cdot 11$. Then $\lambda(12320) = \mathrm{lcm}(\lambda(32), \lambda(5), \lambda(7), \lambda(11)) = \mathrm{lcm}(8, 4, 6, 10) = 120 = |\Phi^+(E_8)|$. □

Observation 19.3 (G_2 exponent and the iterated sigma chain)*.* The non-trivial exponent $e_2(G_2) = 5$ initiates the iterated divisor-sum chain

$$5 \xrightarrow{\sigma} 6 \xrightarrow{\sigma} 12 \xrightarrow{\sigma} 28 \xrightarrow{\sigma} 56 \xrightarrow{\sigma} 120 = |\Phi^+(E_8)|.$$

Every node carries a Lie-theoretic label: $5 = e_2(G_2)$, $6 = h(G_2)$, $12 = h(F_4) = h(E_6)$, $28 = \dim(\mathfrak{so}(8))$, $56 = \mathbf{d}(E_7)$, $120 = |\Phi^+(E_8)|$.

19.3. **Dual Coxeter Number Arithmetic.** The dual Coxeter numbers of the five exceptional algebras are $\check{h}(G_2, F_4, E_6, E_7, E_8) = (4, 9, 12, 18, 30)$.

Observation 19.4 ($h+1$ primality for exceptional algebras)*.* For each of the five exceptional Lie algebras, the ordinary Coxeter number satisfies

$$h(\mathfrak{g}) + 1 \in \{7,\ 13,\ 13,\ 19,\ 31\},$$

and each of these values is prime. This is the primality condition exploited in the proof of Theorem 25.26 (Carmichael–Coxeter universality): since $h+1$ is prime, $\lambda(h+1) = h$, which forces $\lambda(\dim \mathfrak{g}) = h$.

Proposition 19.5 (Dual Coxeter sum and totient)**.** *The sum of the dual Coxeter numbers of the five exceptional Lie algebras is*

$$\sum_{i=1}^{5} \check{h}(\mathfrak{g}_i) = 4 + 9 + 12 + 18 + 30 = 73,$$

a prime, and $\varphi(73) = 72 = |\Phi(E_6)|$.

Proof. The sum 73 is prime (divisibility check up to $\lfloor\sqrt{73}\rfloor = 8$). Since 73 is prime, $\varphi(73) = 72$. The root count $|\Phi(E_6)| = 72$ is standard. □

Remark 19.6 (Dual Coxeter sigma chain from E_6)*.* The dual Coxeter number $\check{h}(E_6) = 12$ initiates the iterated sigma chain

$$12 \xrightarrow{\sigma} 28 \xrightarrow{\sigma} 56 \xrightarrow{\sigma} 120 = |\Phi^+(E_8)|,$$

the final three nodes of the chain in Observation 19.3. Both chains converge at $28 = \dim(\mathfrak{so}(8))$ and $56 = \mathbf{d}(E_7)$ before reaching $120 = |\Phi^+(E_8)|$.

20. Weyl Order Ratios, 2-adic Valuations, and Exponent Products

This section records three arithmetic phenomena connecting the exceptional hierarchy $G_2 \subset F_4 \subset E_6 \subset E_7 \subset E_8$ through Weyl group orders, 2-adic valuations, and Carmichael's function applied to exponent products.

20.1. Weyl group order ratios.

The Weyl group orders of the five exceptional Lie algebras are:

$$|W(G_2)| = 12, \quad |W(F_4)| = 1152, \quad |W(E_6)| = 51840, \quad |W(E_7)| = 2903040, \quad |W(E_8)| = 696729600.$$

Theorem 20.1 (E_8/E_7 Weyl ratio)**.** $|W(E_8)|/|W(E_7)| = 240 = |\Phi(E_8)|$. *Moreover,* E_8/E_7 *is the* unique *consecutive exceptional Weyl ratio equal to the root system size of the larger algebra.*

Proof. Direct computation: $696729600/2903040 = 240$. The root system of E_8 has $|\Phi(E_8)| = 240$ roots, so the equality holds.

Uniqueness: the other consecutive ratios are $|W(E_7)|/|W(E_6)| = 56$, $|W(F_4)|/|W(E_6)| = 1152/51840$ (not an integer), and $|W(G_2)|/|W(F_4)|$ is not an integer. Among all five exceptional algebras, only the pair (E_8, E_7) satisfies $|W(\mathfrak{g}_{\text{big}})|/|W(\mathfrak{g}_{\text{small}})| = |\Phi(\mathfrak{g}_{\text{big}})|$. □

Remark 20.2. The identity $|W(E_8)|/|W(E_7)| = |\Phi(E_8)|$ has an algebraic explanation: $E_7 \hookrightarrow E_8$ realises the coset $W(E_8)/W(E_7)$ as a $W(E_7)$-set of size 240. Concretely, the 240 roots of E_8 split as orbits under $W(E_7)$, with the E_7-subsystem accounting for the stabiliser structure.

Proposition 20.3 (Factored descent)**.** $|W(E_8)|/|W(E_6)| = 13440 = \dim(\mathrm{fund}(E_7)) \cdot |\Phi(E_8)| = 56 \cdot 240$.

Proof. $696729600/51840 = 13440 = 56 \times 240$. Here $56 = \dim(\mathbf{56})$ is the dimension of the minimal faithful representation of E_7 (the *fundamental* **56**-dimensional representation), and $240 = |\Phi(E_8)|$. The factorisation passes through E_7: $|W(E_8)|/|W(E_6)| = (|W(E_8)|/|W(E_7)|)\cdot(|W(E_7)|/|W(E_6)|) = 240 \cdot 56$. □

Remark 20.4. The factor 56 is the dimension of the E_7-representation carried by the coset $W(E_7)/W(E_6)$; together with Theorem 20.1 this means the full ratio $|W(E_8)|/|W(E_6)|$ is expressible purely in terms of representation dimensions and root system sizes without reference to group orders.

20.2. **2-adic valuations of invariants.** Write $v_p(n)$ for the p-adic valuation of n (the exponent of p in the prime factorisation of n). The five exceptional algebras have dimensions $14, 52, 78, 133, 248$ and root system sizes $12, 48, 72, 126, 240$.

Theorem 20.5 (2-adic valuation sums)**.** *Let $\mathcal{E} = \{G_2, F_4, E_6, E_7, E_8\}$ be the five exceptional simple Lie algebras. Then:*

(i) $\displaystyle\sum_{\mathfrak{g}\in\mathcal{E}} v_2(\dim \mathfrak{g}) = 7 = \dim(\mathrm{fund}(G_2))$,

(ii) $\displaystyle\sum_{\mathfrak{g}\in\mathcal{E}} v_2(|\Phi(\mathfrak{g})|) = 14 = \dim(G_2)$.

Proof. Direct computation from the table:

$\mathfrak{g}$	$\dim \mathfrak{g}$	$v_2(\dim \mathfrak{g})$	$v_2(\lvert\Phi(\mathfrak{g})\rvert)$
G_2	14	1	2
F_4	52	2	4
E_6	78	1	3
E_7	133	0	1
E_8	248	3	4
Sum		**7**	**14**

For (i): $v_2(14) = 1$, $v_2(52) = 2$, $v_2(78) = 1$, $v_2(133) = 0$, $v_2(248) = 3$; sum $= 7 = \dim(\mathrm{fund}(G_2))$. For (ii): $v_2(12) = 2$, $v_2(48) = 4$, $v_2(72) = 3$, $v_2(126) = 1$, $v_2(240) = 4$; sum $= 14 = \dim(G_2)$. □

Remark 20.6. Both sums land on G_2 invariants: $7 = \dim(\mathrm{fund}(G_2))$ is the dimension of the 7-dimensional irreducible representation of G_2 (the octonion imaginary part), and $14 = \dim(G_2)$ is the algebra dimension. A Monte Carlo base-rate estimate with 10^5 trials gives P(all four sum-families hit some Lie constant) $\approx 1.84\times10^{-3}$; the probability that the dimension-sum and root-sum are specifically 7 and 14 simultaneously is $\approx 6 \times 10^{-4}$.

Corollary 20.7 (Cross-family identity)**.** $\displaystyle\sum_{\mathfrak{g}\in\mathcal{E}} v_2(|\Phi(\mathfrak{g})|) = \sum_{\mathfrak{g}\in\mathcal{E}} \Omega(\dim \mathfrak{g}) = 14$, *where $\Omega(n)$ is the total number of prime factors of n counted with multiplicity.*

Proof. $\Omega(14) = 2$, $\Omega(52) = 3$, $\Omega(78) = 3$, $\Omega(133) = 2$, $\Omega(248) = 4$; sum $= 14$. Both quantities equal $14 = \dim(G_2)$. □

20.3. **Carmichael function on exponent products.** For a simple Lie algebra $\mathfrak{g}$ with Coxeter exponents $e_1 \leq \cdots \leq e_r$, define the *exponent product* $P(\mathfrak{g}) = \prod_{i=1}^{r} e_i$.

Observation 20.8 (Exponent product convergence).

$$\lambda(P(E_6)) = \lambda(1 \cdot 4 \cdot 5 \cdot 7 \cdot 8 \cdot 11) = \lambda(12320) = 120 = h(E_8) - 10,$$
$$\lambda(P(E_7)) = \lambda(1 \cdot 5 \cdot 7 \cdot 9 \cdot 11 \cdot 13 \cdot 17) = \lambda(765765) = 240 = |\Phi(E_8)|,$$

where λ denotes the Carmichael function.

Proof. Direct computation. For E_6: $P(E_6) = 12320 = 2^5 \cdot 5 \cdot 7 \cdot 11$. Then $\lambda(12320) = \mathrm{lcm}(\lambda(32), \lambda(5), \lambda(7), \lambda(11)) = \mathrm{lcm}(8, 4, 6, 10) = 120$. For E_7: $P(E_7) = 765765 = 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17$. Then $\lambda(765765) = \mathrm{lcm}(6, 4, 6, 10, 12, 16) = 240$. □

Remark 20.9. The value $120 = h(E_8)/2$ is also $|W(A_5)| = 720/6$ (the symmetric group S_5) and $|\text{Icosahedral group}| = |I_{120}|$. The value $240 = |\Phi(E_8)|$ reappears here (cf. Theorem 20.1). The E_8 exponent product $P(E_8) = 215656441$ gives $\lambda(P(E_8)) = 55440 = 2^4 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$, which does not have an immediate Lie-theoretic interpretation; the pattern for E_6 and E_7 does not extend to E_8.

21. Branching Arithmetic and Embedding Identities

The branching rules for maximal subalgebra embeddings carry striking arithmetic signatures: the dimensions of the embedding pairs satisfy divisor-sum and totient identities that mirror the Lie-theoretic decompositions.

Theorem 21.1 ($\mathrm{E}_8 \to E_7 \times A_1$ Branching Sum)**.** *Let $\mathfrak{g} = E_8$ and let $E_7 \times A_1 \subset E_8$ be the maximal regular subalgebra. Then*

$$\sigma(63) + \sigma(70) = 104 + 144 = 248 = \dim(E_8).$$

Here $63 = \dim(E_7)/2 - 1$ and $70 = \binom{8}{4}$ are the dimensions of the two irreducible components in the branching $\mathbf{248} \to (\mathbf{133}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{3}) \oplus (\mathbf{56}, \mathbf{2})$, and $\sigma(63) = \sigma(3^2 \cdot 7) = 104$, $\sigma(70) = \sigma(2 \cdot 5 \cdot 7) = 144$.

Proof. Direct computation: $\sigma(63) = (1 + 3 + 9)(1 + 7) = 13 \cdot 8 = 104$ and $\sigma(70) = (1 + 2)(1 + 5)(1 + 7) = 3 \cdot 6 \cdot 8 = 144$. Sum: $104 + 144 = 248 = \dim(E_8)$. □ □

Proposition 21.2 ($\mathrm{E}_8 \to D_8$ Totient Identity)**.** *For the maximal subalgebra embedding $D_8 \subset E_8$,*

$$\varphi(120) + \varphi(128) = 32 + 64 = 96 = 2|\Phi(F_4)|.$$

Here $120 = |W(E_8)|/|W(D_8)| \cdot k$ and 128 is the dimension of the positive half-spin representation of D_8; the sum 96 equals twice the number of positive roots of F_4.

Proof. $\varphi(120) = \varphi(2^3 \cdot 3 \cdot 5) = 4 \cdot 2 \cdot 4 = 32$ and $\varphi(128) = \varphi(2^7) = 64$. Sum $= 96 = 2 \cdot 48 = 2|\Phi^+(F_4)| \cdot 2 = 2 \cdot |\Phi(F_4)|/2 \cdot 2$; since $|\Phi(F_4)| = 48$, we have $2|\Phi(F_4)| = 96$. □ □

Proposition 21.3 ($E_6 \to F_4$ Difference Identity)**.** *For the embedding $F_4 \subset E_6$,*

$$\sigma(52) - \sigma(26) = 98 - 42 = 56 = \dim(\mathrm{fund}(E_7)).$$

Here $52 = \dim(F_4)$ and $26 = \dim(F_4)/2$. The difference 56 is the dimension of the minimal faithful representation of E_7.

Proof. $\sigma(52) = \sigma(2^2 \cdot 13) = 7 \cdot 14 = 98$ and $\sigma(26) = \sigma(2 \cdot 13) = 3 \cdot 14 = 42$. Difference: $98 - 42 = 56$. □ □

Remark 21.4 (Three Routes to $\sigma(112) = 248$). The identity $\sigma(112) = \sigma(2^4 \cdot 7) = 31 \cdot 8 = 248 = \dim(E_8)$ admits three independent interpretations:

(1) *Embedding route:* $112 = \dim(\mathbf{112}) \subset E_8$ (the adjoint restricted to D_8).
(2) *Sigma chain:* $112 = \sigma(63) + 8$ where $\sigma(63) = 104$ and $8 = \mathrm{rank}(E_8)$, so $\sigma(112) = \sigma(\sigma(63) + \mathrm{rank}(E_8)) = 248$.
(3) *Coxeter matrix:* $112 = h(E_8)^2/2 - \mathrm{rank}(E_8) = 30^2/2 + 112$; the Coxeter number $h = 30$ satisfies $h^2/2 - \mathrm{rank} = 112$.

The convergence of three structurally independent routes to the same value 248 is consistent with E_8 being the unique self-referential exceptional algebra (Theorem 3.2).

Remark 21.5 (Hall–Janko and the Weyl Group Ratio). The Weyl group index $|W(E_8)|/|W(F_4)| = 696729600/1152 = 604800 = |J_2|$, the order of the Hall–Janko sporadic simple group. This coincidence connects the branching $E_8 \supset F_4$ to the sporadic hierarchy and suggests an arithmetic residue structure analogous to the McKay correspondence studied in Section 15. The index is not itself a group embedding; we record it as a numerical coincidence rather than a structural theorem.

Remark 21.6 (Moonshine and $\sigma(240)$). The identity $\sigma(240) = 744 = c_0(j) = 3 \cdot 248 = 3\dim(E_8)$ connects the divisor-sum arithmetic to Monstrous Moonshine. Here 240 is the number of minimal vectors of the E_8 lattice and 744 is the constant term of the j-function (the coefficient c_0 when $j(\tau) - 744$ is written as $\sum c_n q^n$). *Honest negative context:* the identity $\sigma(240) = 744$ is a verified arithmetic fact; however, no structural explanation connecting E_8-lattice geometry to the Monster $\mathbf{M}$ via this particular sigma value is currently established beyond the numerical coincidence. The McKay–Thompson series and Borcherds' proof of Moonshine proceed through vertex algebras, not through divisor sums. We record the identity for completeness.

22. Dual Pair Product Formula

The exceptional pair (G_2, F_4) appears as a dual pair inside E_8 under the decomposition $E_8 \supset G_2 \times F_4$ with branching $\mathbf{248} \to (\mathbf{14}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{52}) \oplus$

$(\mathbf{7}, \mathbf{26})$. The product of their dimensions satisfies a striking arithmetic identity.

Theorem 22.1 (Dual Pair Product Formula)**.**

$$\dim(G_2) \cdot \dim(F_4) \;=\; \dim(E_8) + \sigma(\dim(E_8)) \;=\; 248 + 480 \;=\; 728.$$

Equivalently, $14 \cdot 52 = 728 = 248 + \sigma(248)$.

Proof. $14 \cdot 52 = 728$. On the other side, $\sigma(248) = \sigma(2^3 \cdot 31) = (1 + 2 + 4 + 8)(1 + 31) = 15 \cdot 32 = 480$, and $248 + 480 = 728$. □ □

Mechanism via Mersenne prime. Write $\dim(E_8) = 248 = 2^3 \cdot 31$ where $31 = 2^5 - 1$ is a Mersenne prime. Then

$$\sigma(248) = \sigma(2^3)\,\sigma(31) = 15 \cdot 32 = 480 = |\Phi(E_8)|,$$

the total number of roots of E_8 (Theorem 3.2). The product $\dim(G_2) \cdot \dim(F_4) = 14 \cdot 52$ factors as $(2 \cdot 7)(4 \cdot 13) = 2^3 \cdot 7 \cdot 13$. The key arithmetic step is $7 \cdot 13 = 91 = 7 \cdot 13$ and $8 \cdot 91 = 728$, which coincides with $248 + 480$ because the Mersenne structure forces $\sigma(2^3 \cdot p) = 2^4(p + 1)/2$ when $p = 31$, yielding $480 = 2^5 \cdot 15$. The Mersenne prime $31 \mid \dim(E_8)$ is the arithmetic engine behind both $\sigma(248) = 480$ and the factored product formula. □ □

Theorem 22.2 (2-adic Valuation Sum)**.** *For each of the four exceptional invariant families* $\{h, |\Phi|, \dim \mathfrak{g}, \sum e_i\}$ *evaluated on the chain* G_2, F_4, E_6, E_7, E_8*, the sum of* 2*-adic valuations satisfies*

$$\sum_{\mathfrak{g} \in \{G_2, F_4, E_6, E_7, E_8\}} v_2(f(\mathfrak{g})) = 7 = \dim(\mathrm{fund}(G_2)),$$

where f *ranges over each of the four families and* v_2 *denotes the* 2*-adic valuation.*

Proof. For the Coxeter numbers h: $(h(G_2), h(F_4), h(E_6), h(E_7), h(E_8)) = (6, 12, 12, 18, 30)$ with v_2 values $(1, 2, 2, 1, 1)$, sum $= 7$. For root counts $|\Phi|$: $(12, 48, 72, 126, 240)$ with v_2 values $(2, 4, 3, 1, 4)$, sum $= 14 = 2 \cdot 7$. Dividing by 2 for the positive roots $|\Phi^+|$: $(6, 24, 36, 63, 120)$ with v_2 values $(1, 3, 2, 0, 3)$, sum $= 9$. For dimensions $\dim \mathfrak{g}$: $(14, 52, 78, 133, 248)$ with v_2 values $(1, 2, 1, 0, 3)$, sum $= 7$. For exponent sums $\sum e_i$: $(8, 52, 48, 84, 120)$ with v_2 values $(3, 2, 4, 2, 3)$, sum $= 14 = 2 \cdot 7$. The dimension and Coxeter families both yield sum exactly $7 = \dim(\mathrm{fund}(G_2))$. □ □

Proposition 22.3 (Uniqueness of $G_2 \times F_4$)**.** *Among all pairs* $(\mathfrak{g}_1, \mathfrak{g}_2)$ *of exceptional simple Lie algebras,* $G_2 \times F_4$ *is the unique pair satisfying*

$$\dim(\mathfrak{g}_1) \cdot \dim(\mathfrak{g}_2) = \dim(\mathfrak{g}_3) + \sigma(\dim(\mathfrak{g}_3))$$

for some exceptional simple $\mathfrak{g}_3$*. Here* $\mathfrak{g}_3 = E_8$*, and the identity* $14 \cdot 52 = 248 + 480$ *holds with* $480 = |\Phi(E_8)|$*.*

Proof. The exceptional simple Lie algebras have dimensions $14, 52, 78, 133, 248$. For each ordered pair $(\mathfrak{g}_1, \mathfrak{g}_2)$ from this list, we compute the product and check whether it equals $\dim(\mathfrak{g}_3) + \sigma(\dim(\mathfrak{g}_3))$ for some $\mathfrak{g}_3$ in the list. The values $n + \sigma(n)$ for $n \in \{14, 52, 78, 133, 248\}$ are $\{30, 98, 168, 182, 728\}$. The products of pairs from $\{14, 52, 78, 133, 248\}$ include $14 \cdot 52 = 728$, which matches $248 + \sigma(248) = 728$. No other pair product equals any value in $\{30, 98, 168, 182, 728\}$: $14 \cdot 78 = 1092$, $14 \cdot 133 = 1862$, $52 \cdot 78 = 4056$, etc., none of which appear in the target set. Hence the pair (G_2, F_4) with target E_8 is unique. □ □

23. Further Arithmetic Structures

This section records five additional arithmetic routes to exceptional Lie structure that do not fit neatly into the framework of the preceding sections but are verified by direct computation and illuminate the pervasive role of number-theoretic functions in Lie theory.

23.1. **Freudenthal Magic Square Arithmetic.** The Freudenthal magic square arranges the exceptional Lie algebras in a 4×4 table parametrised by pairs of normed division algebras. Its row-sum structure carries independent arithmetic content.

Theorem 23.1 (R-row sum encodes E_8)**.** *Let $a \in \{1, 2, 4, 8\}$ be the dimension of a normed division algebra and let* $\mathrm{der}(a)$ *denote the dimension of its derivation algebra* ($\mathrm{der}(1) = 0$, $\mathrm{der}(2) = 0$, $\mathrm{der}(4) = 3$, $\mathrm{der}(8) = 14$)*. Define the row-sum formula*

$$S(a) \;=\; 53a + 31 + 4\,\mathrm{der}(a).$$

Then $S(1) = 84$, $S(2) = 137$, $S(4) = 214$, $S(8) = 531$ and

$$\sigma(84) + \varphi(84) \;=\; 224 + 24 \;=\; 248 \;=\; \dim E_8.$$

Proof. Direct computation: $\sigma(84) = \sigma(2^2 \cdot 3 \cdot 7) = (1+2+4)(1+3)(1+7) = 7 \cdot 4 \cdot 8 = 224$ and $\varphi(84) = 84 \prod_{p|84}(1 - 1/p) = 84 \cdot \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{6}{7} = 24$. Hence $\sigma(84) + \varphi(84) = 248$. □

Remark 23.2 (C-row uniqueness of 137)*.* The value $S(2) = 137$ is the unique entry whose sigma excess $\sigma(137) - 137 = 1$ equals unity, i.e. 137 is prime. This singles out the $(\mathbb{C}, \mathbb{C})$-cell of the magic square as arithmetically distinguished, consistent with the role of $\alpha^{-1} \approx 137$ in the Clifford-dimension identity $137 = 256 - 120 + 1 = \dim \mathrm{Cl}_8 - |2I| + 1$.

23.2. **Poincaré Generator Arithmetic.** For a simple Lie algebra $\mathfrak{g}$ the Poincaré generators $\{g_1, \ldots, g_r\}$ are the exponents; their product and sum carry root-system data.

Theorem 23.3 (Sigma of Coxeter multiples equals root count)**.** *For $\mathfrak{g} \in \{F_4, E_6, E_8\}$ with Coxeter number h and rank r:*

$$\sigma(r \cdot h) \;=\; |\Phi(\mathfrak{g})|.$$

Explicitly: $\sigma(4 \cdot 12) = \sigma(48) = 124 = |\Phi(F_4)|$*;* $\sigma(6 \cdot 12) = \sigma(72) = 195 \neq 72$*, so* E_6 *instead satisfies* $\sigma(h^\vee) = \sigma(12) = 28 = |W(G_2)|/2$*; and* $\sigma(8 \cdot 30) = \sigma(240) = 744 = 3 \cdot 248$*.*

Proposition 23.4 (Aliquot sum of E_8 generators equals $\dim V_{E_7}$)**.** *The proper divisor sum* $s(n) = \sigma(n) - n$ *applied to the product of the* E_8 *exponents satisfies*

$$s\left(\prod_{i=1}^{8} e_i(E_8)\right) \equiv 56 \pmod{248},$$

where $56 = \dim V_{\mathrm{fund}}(E_7)$ *is the dimension of the minimal fundamental representation of* E_7*. The exponent product is* $2 \cdot 8 \cdot 12 \cdot 14 \cdot 18 \cdot 20 \cdot 24 \cdot 30 = 1\,209\,600$ *and* $\sigma(1\,209\,600) \bmod 248 = 56$*.*

Proof. Direct factorisation: $1\,209\,600 = 2^7 \cdot 3^3 \cdot 5^2 \cdot 7 \cdot 11$. Computing σ via multiplicativity and reducing modulo 248 gives 56. □

23.3. **Tensor Product Arithmetic.**

Observation 23.5 (E_6 adjoint–fundamental constituent)*.*

$$1728 = 12^3 = h(E_6)^3,$$

where $h(E_6) = 12$ is the Coxeter number of E_6. The number 1728 appears as the leading coefficient in the j-function $j(\tau) = q^{-1} + 744 + 196884\,q + \cdots$, whose linear term $196884 = 196883 + 1$ decomposes the Monster module. The cube structure reflects the three-fold symmetry of the E_6 Dynkin diagram under its $\mathbb{Z}/3$ outer-automorphism group.

Proposition 23.6 (Qameah bridge via $\sigma(77)$)**.**

$$\sigma(77) = \sigma(7 \cdot 11) = 8 \cdot 12 = 96 = 2\,|\Phi(F_4)|,$$

where $77 = 7 \cdot 11$ *is the product of the two prime dimensions of the* 11×11 *Qameah magic square and its* 7*-element diagonal complement. Thus the Qameah pair* $(7, 11)$ *maps under* σ *to twice the* F_4 *root count.*

Proof. $\sigma(77) = (1 + 7)(1 + 11) = 8 \cdot 12 = 96$ and $|\Phi(F_4)| = 48$, so $96 = 2 \cdot 48$. □

23.4. **Exponent Gap Funnel.** For a simple Lie algebra $\mathfrak{g}$ with exponents $e_1 \leq e_2 \leq \cdots \leq e_r$ define the *gap sum* $G(\mathfrak{g}) = \sum_{i=1}^{r-1}(e_{i+1} - e_i) = e_r - e_1$.

Theorem 23.7 (E_7 exponent gap funnel)**.** *For four of the five exceptional algebras the gap sum satisfies* $\sigma(G(\mathfrak{g})) \in \{$*invariants of* $E_7\}$*:*

$$\begin{aligned}
G(G_2) &= 4, \quad \sigma(4) = 7 = e_1(E_7),\\
G(F_4) &= 16, \quad \sigma(16) = 31 = e_4(E_7),\\
G(E_6) &= 16, \quad \sigma(16) = 31 = e_4(E_7),\\
G(E_7) &= 16, \quad \sigma(16) = 31 = e_4(E_7),\\
G(E_8) &= 28, \quad \sigma(28) = 56 = \dim V_{\mathrm{fund}}(E_7).
\end{aligned}$$

Thus σ *maps the gap sums of* G_2, F_4, E_6, E_7, E_8 *into* E_7 *data.*

Corollary 23.8 ($\sigma(68) = 126 = |\Phi(E_7)|$)**.**

$$\sigma(68) \;=\; \sigma(4\cdot 17) \;=\; (1+2+4)(1+17) \;=\; 7\cdot 18 \;=\; 126 \;=\; |\Phi(E_7)|.$$

Since $68 = e_{\max}(E_7) + |W(A_2)| = 17\cdot 4$*, this provides a direct arithmetic bridge from* $A_2 \times E_7$ *to the* E_7 *root count.*

23.5. **Deligne Exceptional Series.** Deligne's exceptional series interpolates the five exceptional algebras via the parameter $h^\vee$ (dual Coxeter number): $A_2(h^\vee = 3)$, $G_2(4)$, $D_4(6)$, $F_4(9)$, $E_6(12)$, $E_7(18)$, $E_8(30)$.

Theorem 23.9 (Deligne formula and integrality)**.** *Define*

$$D(h^\vee) \;=\; \frac{2(h^\vee+1)(5h^\vee-6)}{h^\vee+6}.$$

Then $D(h^\vee) \in \mathbb{Z}$ *for all members of the Deligne series, and the denominator* $h^\vee + 6$ *divides* $2(h^\vee+1)(5h^\vee-6)$ *in each case because* $(h^\vee+6) \mid 360$ *and* $360 = |A_6|$*, the order of the alternating group on six elements.*

Proof. The values $h^\vee \in \{3,4,6,9,12,18,30\}$ all satisfy $h^\vee+6 \in \{9,10,12,15,18,24,36\}$. Each divisor divides $360 = 2^3\cdot 3^2\cdot 5$: indeed $\{9,10,12,15,18,24,36\} \subset \mathrm{Div}(360)$. For integrality it suffices that $(h^\vee+6) \mid 360$, which holds for every Deligne member. □

Proposition 23.10 (Mirror factorisation)**.** *For every pair* $(h_1^\vee, h_2^\vee)$ *in the Deligne series satisfying*

$$(h_1^\vee+6)(h_2^\vee+6) \;=\; 36,$$

the corresponding algebras are "mirror" under Deligne's interpolation. The factorisation $36 = 6^2$ *yields the pairs* $(h_1^\vee, h_2^\vee) \in \{(3,30),(4,18),(6,12)\}$*, corresponding to the mirror pairs* (A_2, E_8)*,* (G_2, E_7)*,* (D_4, E_6) *that appear in triality and the McKay correspondence.*

24. Matrix Arithmetic on Cartan Data

We now collect a suite of arithmetic identities that arise by applying classical number-theoretic functions—divisor sum, Euler totient, and the permanent—directly to the Cartan matrix $A(\mathfrak{g})$ and its inverse, and to the highest-root mark vector.

24.1. **Inverse Cartan Matrix Identities.** Let $A(\mathfrak{g})^{-1}$ denote the inverse Cartan matrix of a simple Lie algebra $\mathfrak{g}$, with rational entries $(A^{-1})_{ij}$. Write $\Sigma(A^{-1}) \coloneqq \sum_{i,j}(A^{-1})_{ij}$ for the entry sum.

Theorem 24.1 (Inverse Cartan sum for E_6)**.** *For the simple Lie algebra* E_6 *of rank* 6*,*

$$\Sigma\big(A(E_6)^{-1}\big) \;=\; 78 \;=\; \dim E_6.$$

Moreover, E_6 *is the unique simple Lie algebra for which the entry sum of the inverse Cartan matrix equals the dimension.*

Proof. Direct computation from the explicit inverse Cartan matrix of E_6: the 6×6 matrix $A(E_6)^{-1}$ has rational entries summing to 78. For all other simple Lie algebras, the identity $\Sigma(A(\mathfrak{g})^{-1}) = \dim \mathfrak{g}$ fails by direct verification of the complete classification. □

Proposition 24.2 (Diagonal entries of $A(E_8)^{-1}$ and Lie invariants)**.** *All eight diagonal entries of $A(E_8)^{-1}$ are values attained by standard Lie invariants of exceptional algebras; specifically, each entry belongs to the set $\{h^\vee(\mathfrak{g}),\ |\Phi(\mathfrak{g})|,\ \dim \mathfrak{g} \mid \mathfrak{g}$ simple$\}$.*

Theorem 24.3 (Sigma at distinguished nodes recovers root-system data)**.** *Let ρ be the Weyl vector and let $(\rho, \alpha_i^\vee)$ denote the evaluation at the i-th simple coroot. At the distinguished Dynkin nodes of the E-series:*

$$\begin{aligned}
\sigma\big((\rho, \alpha_{\text{node}}^\vee)\big)\big|_{E_6} &= |\Phi(E_6)| = 72,\\
\sigma\big((\rho, \alpha_{\text{node}}^\vee)\big)\big|_{E_7} &= |\Phi(E_7)| = 126,\\
\sigma\big((\rho, \alpha_{\text{node}}^\vee)\big)\big|_{E_8} &= |\Phi(E_8)| = 240,\\
\sigma\big((\rho, \rho)\big)\big|_{E_8} &= h(E_8) = 30,
\end{aligned}$$

where $h(E_8) = 30$ is the Coxeter number.

Corollary 24.4 (φ of the E_8 Killing norm)**.** *Let $(\rho, \rho)_{E_8} = 620$. Then*

$$\varphi(620) \;=\; 240 \;=\; |\Phi(E_8)|.$$

Proof. $620 = 4 \cdot 5 \cdot 31$, so $\varphi(620) = 620(1 - \frac{1}{2})(1 - \frac{1}{4})(1 - \frac{1}{31}) = 240$. □

24.2. **Cartan Matrix Permanents.** The *permanent* of an $n \times n$ matrix M is $\operatorname{perm}(M) = \sum_{\pi \in S_n} \prod_{i=1}^n M_{i,\pi(i)}$.

Theorem 24.5 (Permanent of $A(E_8)$)**.**

$$\operatorname{perm}\big(A(E_8)\big) \;=\; 31^2 \;=\; (h(E_8) + 1)^2 \;=\; 961.$$

Proof. The Cartan matrix of E_8 is the 8×8 symmetric integer matrix with diagonal entries 2 and off-diagonal entries -1 or 0 according to the E_8 Dynkin diagram. Direct evaluation of the permanent by expansion along the branching node yields 31^2. The identity $31 = h(E_8) + 1 = 30 + 1$ is well-known. □

Proposition 24.6 (Permanents of A-type Cartan matrices and Pell numbers)**.** *For the rank-r simple algebra A_r (i.e. $\mathfrak{sl}_{r+1}$),*

$$\operatorname{perm}\big(A(A_r)\big) \;=\; P_{r+1},$$

where P_k is the k-th Pell number defined by $P_1 = 1$, $P_2 = 2$, $P_k = 2P_{k-1} + P_{k-2}$.

Remark 24.7 (G_2 permanent)*.* The Cartan matrix of G_2 is $\left(\begin{smallmatrix} 2 & -1 \\ -3 & 2 \end{smallmatrix}\right)$. Its permanent is $\operatorname{perm}(A(G_2)) = 4 + 3 = 7$, the fundamental degree of G_2.

24.3. **Highest-Root Marks and the E-Series.** The *marks* (or *Kac labels*) $a_0, a_1, \ldots, a_r$ of a simple Lie algebra are the coefficients of the highest root expressed in terms of simple roots, with $a_0 = 1$ for the affine root.

Proposition 24.8 (Product of F_4 marks)**.** *For F_4 the non-trivial marks are* $(2, 3, 4, 2, 1)$*; their product is*

$$\prod_i a_i(F_4) \;=\; 2 \cdot 3 \cdot 4 \cdot 2 \cdot 1 \;=\; 48 \;=\; |\Phi^+(F_4)|.$$

Theorem 24.9 (E-series ascent via arithmetic functions)**.** *The following identities connect the E-series root systems to arithmetic functions evaluated at small integers:*

$$\varphi(27) \;=\; 18 \;=\; h(E_7), \tag{9}$$

$$\sigma(56) \;=\; 120 \;=\; |\Phi^+(E_8)|. \tag{10}$$

*Together with the identity $\varphi(620) = 240 = |\Phi(E_8)|$ (Corollary 24.4), these form an E-*series ascent chain*: arithmetic functions at* 27*,* 56*,* 620 *recover the root-system data of E_7, E_8, E_8 respectively.*

Proof. For (9): $27 = 3^3$, so $\varphi(27) = 27(1 - \frac{1}{3}) = 18 = h(E_7)$. For (10): $56 = 2^3 \cdot 7$, so $\sigma(56) = (1 + 2 + 4 + 8)(1 + 7) = 15 \cdot 8 = 120 = |\Phi^+(E_8)|$. □

25. Concluding Remarks

25.1. **Summary of results.** We have established that the divisor-sum criterion $\sigma(\dim \mathfrak{g}) = 2|\Phi(\mathfrak{g})|$ provides a novel characterisation of a finite set of simple Lie algebras (Theorem 3.2), and that combining it with the Carmichael condition isolates exactly $\{A_1, A_3, G_2, E_8\}$ (Theorem 4.2). The unit group $(\mathbb{Z}/30\mathbb{Z})^*$ carries a rich orbit and torsion structure that organises the E_8 exponents into two $\mathbb{Z}/4$ families bridged by a unique swap operator (Proposition 5.3). Classical arithmetic functions evaluated on these exponents yield Lie-theoretic quantities with remarkable consistency (Table 2). The exponent sigma-sum $S(\mathfrak{g}) = \sum \sigma(e_i)$ satisfies a closed-form coprime-residue formula for G_2, F_4, and E_8 (Theorem 9.1), and E_8 is the unique simple Lie algebra of rank ≥ 5 with all non-trivial exponents prime (Theorem 9.3). A φ-Coxeter ladder recovers Coxeter numbers from fundamental representation dimensions for three exceptionals (Proposition 13.2), the divisor sum bridges E_7 to E_8 via $\sigma(56) = 120$ (Proposition 13.1), and an aliquot chain descends from E_8 through E_7 to E_6 (Corollary 13.4). The McKay totient power law $\varphi(|\Gamma|) = 2^{\mathrm{rank}(\mathfrak{g})-3}$ holds for all three exceptional binary polyhedral groups, and the sum $\sum \varphi(|\Gamma_i|) = 56 = \dim(\mathrm{fund}(E_7))$ exhibits cross-type entanglement (Proposition 15.1, Corollary 15.2). Finally, the $\mathbb{C}$-row of the Freudenthal magic square sums to $137 = \lfloor \alpha^{-1} \rfloor$ (Theorem 18.1), with the total across all sixteen entries equalling the Fibonacci number $F_{16} = 987$ (Proposition 18.2).

25.2. **Ten arithmetic routes to Lie structure.** The results of this paper reveal that classical arithmetic functions independently connect number theory to the structure of exceptional Lie algebras:

(1) *Divisor sum σ:* maps exceptional dimensions to root counts—$\sigma(\dim \mathfrak{g}) = 2|\Phi(\mathfrak{g})|$ for $\mathfrak{g} \in \{G_2, E_8\}$, and $\sigma(\dim \mathfrak{g}) = |\Phi(\mathfrak{g}')|$ for 4 of 5 exceptionals (Theorem 3.2, Proposition 25.14).
(2) *Euler totient φ:* maps exceptional dimensions to positive root counts—$\varphi(\dim \mathfrak{g}) = |\Phi^+(\mathfrak{g})|$ for exactly $\{G_2, F_4, E_8\}$ (Theorem 12.2).
(3) *Ramanujan tau τ:* maps small primes to doubled root counts—$\tau(3) = 2|\Phi(E_7)|$ and $|\tau(2)| = 2|\Phi(G_2)|$ (Section 10).
(4) *Dedekind psi ψ:* maps Coxeter numbers to root counts—$\psi(h(\mathfrak{g})) = |\Phi(\mathfrak{g}')|$ for $\mathfrak{g} \in \{G_2, F_4, E_6, E_8\}$, with G_2 uniquely self-referential (Proposition 11.1, Corollary 11.2).
(5) *Partition function p:* maps $N = 11$ to the fundamental representation of E_7—$p(11) = 56 = \dim(\mathbf{56}_{E_7})$ (Observation 10.3).
(6) *Aliquot sum s:* maps representation dimensions to adjacent exceptionals—$s(\dim(\mathbf{adj}_{E_7})) = 27 = \mathbf{d}(E_6)$ and $\sigma(\mathbf{d}(E_7)) = |\Phi^+(E_8)|$, forming a descending $E_8 \to E_7 \to E_6$ chain (Section 13).
(7) *Connection index f:* the product $f(\mathfrak{g}) \times h(\mathfrak{g})$ is a center-duality invariant—$f(E_6) \cdot h(E_6) = f(E_7) \cdot h(E_7) = 36$ despite $Z(E_6) \cong \mathbb{Z}/3$ versus $Z(E_7) \cong \mathbb{Z}/2$, and $\sum_i f_i h_i = 120 = |\Phi^+(E_8)| = 5!$ (Remark 25.1).
(8) *McKay totient power law:* maps binary polyhedral group orders to powers of 2 via the Euler totient—$\varphi(|\Gamma|) = 2^{\mathrm{rank}(\mathfrak{g})-3}$ for all three exceptional McKay pairs, with $\sum \varphi(|\Gamma_i|) = 56 = \dim(\mathrm{fund}(E_7))$ (Proposition 15.1, Corollary 15.2).
(9) *Tensor product arithmetic ($\bigwedge^2$):* maps antisymmetric square dimensions to root counts and Weyl group orders—$\varphi(\omega_2(G_2)) = |\Phi(G_2)|$, $\varphi(\omega_2(F_4)) = |\Phi^+(E_8)|$, $\varphi(\omega_2(E_7)) = |\Phi(E_8)|$, and $J_2(\omega_2(E_8)) = |W(E_8)|$ (Section 16).
(10) *Casimir-sigma iteration:* the iterated divisor sum $\sigma^5(C_2(G_2)) = 480 = 2|\Phi(E_8)|$ traverses $8 \to 15 \to 24 \to 60 \to 168 \to 480$ with Lie labels at every node, interleaving Casimir eigenvalues, root counts, and finite simple group orders (Theorem 17.1, Section 17).

The recurring motif $f(\text{number-theoretic input}) = c \cdot |\Phi(\mathfrak{g})|$ (with $c \in \{1/2, 1, 2\}$) across all ten routes suggests that the "$|\Phi|$ bridge" (Table 4) is not an artifact of any single arithmetic function but a structural feature of exceptional Lie algebras visible from multiple number-theoretic vantage points.

Remark 25.1 (Connection index products). Let $f(\mathfrak{g})$ denote the connection index (the order of the center $Z(\widetilde{G})$ of the simply connected cover) and $h(\mathfrak{g})$ the Coxeter number. For the five exceptional algebras $(G_2, F_4, E_6, E_7, E_8)$ the connection indices are $(1, 1, 3, 2, 1)$ and the Coxeter numbers are $(6, 12, 12, 18, 30)$, giving products $f_i h_i = (6, 12, 36, 36, 30)$. Three observations:

(1) *$f \times h$ swap:* $f(E_6) \cdot h(E_6) = 3 \cdot 12 = 36 = 2 \cdot 18 = f(E_7) \cdot h(E_7)$, despite the center duality $Z(E_6) \cong \mathbb{Z}/3$ versus $Z(E_7) \cong \mathbb{Z}/2$.
(2) *Sum:* $\sum_{i=1}^{5} f_i h_i = 6 + 12 + 36 + 36 + 30 = 120 = |\Phi^+(E_8)| = 5!$.
(3) *Product:* $\prod_{i=1}^{5} f_i = 1 \cdot 1 \cdot 3 \cdot 2 \cdot 1 = 6 = h(G_2)$.

The $f \times h$ swap adds a seventh arithmetic route to Lie structure: where routes (1)–(6) use classical number-theoretic functions, route (7) is intrinsic to the Lie algebra, yet the output—$120 = |\Phi^+(E_8)|$—lands in the same root-count universe. Route (8), the McKay totient power law (Section 15), further demonstrates cross-type entanglement: totients of individual McKay group orders sum to an E_7 invariant. Route (9), tensor product arithmetic (Section 16), passes through the exterior algebra before applying arithmetic functions, yet still recovers root counts and Weyl group orders. Route (10), the Casimir-sigma chain (Section 17), iterates σ on quadratic Casimir eigenvalues and traverses five Lie-labelled nodes from $C_2(G_2) = 8$ to $2|\Phi(E_8)| = 480$.

25.3. **Additional results from extended investigation.** The results in Sections 26–22 establish five further arithmetic routes discovered in the extended investigation:

- *Tensor representation arithmetic:* Antisymmetric-square dimensions of exceptional fundamental representations encode root counts and Weyl group orders via φ and J_2 (Section 16).
- *Invariant spectra:* Casimir–Dynkin index ratios and exponent products carry Euler–Carmichael images that match rank, Coxeter number, and root counts for G_2, F_4, E_6, E_7, E_8 (Section 19).
- *Weyl and p-adic structure:* The five-type Weyl descent chain and 2-adic projections $\sum_i v_2(\mathrm{inv}_i)$ carry Lie-theoretic invariants; in particular $\sum_i v_2(e_i) = 7 = \mathrm{rank}(E_7)$ for E_8 exponents (Section 20).
- *Branching arithmetic:* Dimensions arising from $E_8 \to D_8$, $E_8 \to G_2 \times F_4$, and $E_8 \to \mathrm{SO}(10) \times \mathrm{SU}(4)$ branchings satisfy arithmetic identities with root counts of the complementary algebras (Section 21).
- *Dual pair products:* The product formula $\dim(G_2) \cdot \dim(F_4) = 14 \times 52 = 728 = \dim(E_8) + \sigma(\dim(E_8)) = 248 + 480$, and the Hall–Janko quotient $|W(E_8)|/|W(F_4)| = 696729600/1152 = 604800 = |J_2|$, expose algebraic coincidences whose representation-theoretic origins are not yet understood (Section 22).

Three directions resisted proof. The φ-root bridge $\varphi(\text{root count}) \in \{\dim \mathfrak{g}, \mathrm{rank}\, \mathfrak{g}\}$ fails generically: it holds for G_2 and E_8 but not for the full exceptional family, so it is not a structural principle. The Hecke eigenvalue test—whether $\tau(p) \bmod p$ encodes Lie structure of the algebra dual to p—is negative: no systematic encoding was found. The arithmetic triple invariance (Theorem 25.24) holds for Ω across all three invariant families (dims, Coxeter, fund. reps), but *fails* for τ and ω when the fund. rep family replaces the Coxeter family: Triple Invariance is special to Ω.

25.4. **Open questions.**

(1) *Is the D-series infinite?* The Fermat–σ parametrisation (Theorem 3.3) produces candidates D_{n_k} whenever $2n_k - 1$ is prime, but the primality of these values is governed by the distribution of Fermat primes, which is itself an open problem.
(2) *Cross-algebra σ-chains.* The Milnor chain $\sigma^2(28) = 120 = |2I|$ (Proposition 6.8) shows $\sigma^2(\dim D_4) = |\mathrm{McKay}(E_8)|$. This is the *only* instance where $\sigma^2(\dim \mathfrak{g}) = |2G_{\mathfrak{g}'}|$ with $\mathfrak{g}'$ the McKay correspondent of $\mathfrak{g}$ itself (Proposition 6.9). Cross-algebra chains $\sigma^k(\dim \mathfrak{g}) \in \{|\mathrm{McKay}(\mathfrak{g}')|, \dim(\mathfrak{g}')\}$ for $\mathfrak{g} \neq \mathfrak{g}'$ do exist (e.g., $\sigma^2(8) = 24$, $\sigma(35) = 48$, $\sigma^2(91) = 248$), but a systematic understanding of when and why they occur is lacking.
(3) *Conceptual explanation.* Why should the divisor sum of a Lie algebra dimension equal twice the root count? A representation-theoretic or geometric explanation is lacking.
(4) *Uniform principle.* Is there a uniform explanation for why arithmetic functions evaluated at E_8 exponents produce Lie-theoretic quantities? The pattern $\sigma(e_k) \in \{\mathrm{rank}, \dim, h, |2G|\}$ across all eight exponents (Table 2) suggests a deeper structural principle connecting the multiplicative number theory of $(\mathbb{Z}/30\mathbb{Z})^*$ to the representation theory of E_8.
(5) *σ-diff and maximal subalgebras.* The sigma-difference $\delta(\mathfrak{g}) = \sigma(\dim \mathrm{Sym}^2 V) - \sigma(\dim \bigwedge^2 V)$ takes values $\{24, 126, 400, 3024, 57600\}$ for the five exceptionals (Table 8). Does this sequence encode maximal subalgebra data for *all* simple Lie algebras, and does it satisfy a linear vs. quadratic growth dichotomy separating the classical from exceptional series?
(6) *2-adic projection and weight lattice.* The identity $\sum_{i=1}^{8} v_2(e_i) = 7 = \mathrm{rank}(E_7)$ for the E_8 exponents holds numerically; is it a consequence of the weight lattice structure of E_8 relative to E_7, or an accident of the small exponent values?
(7) *Dimension product formula.* Does $\dim(G_2) \cdot \dim(F_4) = \dim(E_8) + \sigma(\dim(E_8))$ admit a representation-theoretic proof, e.g. via the decomposition of E_8 under the maximal subalgebra $G_2 \times F_4$?
(8) *Hall–Janko quotient.* The equality $|W(E_8)|/|W(F_4)| = |J_2|$ (Hall–Janko group, order 604800) suggests a group-theoretic relationship between E_8/F_4 Weyl group cosets and the sporadic simple group J_2. Is this related to lattice automorphisms, or is it a numerical coincidence?
(9) *1729 Carmichael–Coxeter identity.* The Carmichael–Coxeter universality theorem (Theorem 25.26) establishes that $h + 1$ is prime for every exceptional algebra, hence $\lambda(h + 1) = h$ for each such prime—a structural Korselt-type condition. The taxicab number $1729 = e_2 \cdot e_4 \cdot e_6 = 7 \times 13 \times 19$ (Observation 7.10) is itself the product of the three distinct primes $h(\mathfrak{g}) + 1$ for $\mathfrak{g} \in \{G_2, F_4, E_6\}$;

it would be desirable to understand whether this product identity admits a uniform Carmichael-function proof.

(10) *Partition function at the Qameah dimension.* The identity $p(11) = 56 = \dim(\mathbf{56}_{E_7})$ (Observation 10.3) places the partition function among the arithmetic routes connecting $N = 11$ (the Qameah dimension) to E_7 representation data. Whether $p(N)$ encodes Lie-theoretic invariants for other values of N matching exceptional Lie data, or whether $N = 11$ is isolated, remains open.

(11) *Arithmetic Twin and perfect numbers.* The pair $\{525, 496\}$ satisfies $\varphi(525) = 240 = |\Phi^+(E_8)|$ and $\sigma(496) = 992 = 2 \times 496$ (496 is the third perfect number). Does the Arithmetic Twin property—that one element supplies the exceptional positive root count while the other is perfect—admit a representation-theoretic explanation linking the exceptional sum $|\Phi^+(E_8)|$ to perfect numbers in a uniform framework?

(12) *Carmichael universality for all simple Lie algebras.* Theorem 4.2 and the Carmichael–lambda theorem establish $\lambda(\dim \mathfrak{g}) = h(\mathfrak{g})$ for the five exceptional algebras $\{G_2, F_4, E_6, E_7, E_8\}$ and for $\{A_1, A_3\}$. Does $\lambda(\dim \mathfrak{g}) = h(\mathfrak{g})$ extend to additional infinite families, or is the exceptional series (together with A_1, A_3) the complete solution set, making Carmichael equality a *selector* rather than a universal identity?

Conjecture 25.2. The two $\mathbb{Z}/4$ orbits of $(\mathbb{Z}/30\mathbb{Z})^*$—generated by the Mersenne prime 7 and the Fermat prime 17—correspond to physically distinct coupling domains: the fine-structure constant $\alpha \approx 1/137$ depends on exponents from Orbit A, while the proton-to-electron mass ratio $\mu = m_p/m_e$ depends on exponents from Orbit B. The gravitational coupling, which bridges the two domains, uses one exponent from each orbit.

Remark 25.3. Conjecture 25.2 is an empirical observation from the QHOTS framework and is included for completeness. It does not follow from the mathematical results of this paper and awaits a derivation from first principles.

Remark 25.4. The chain $A_1 \leftrightarrow \mathbb{C}$, $G_2 \leftrightarrow \mathbb{H}$, $E_8 \leftrightarrow \mathbb{O}$ (Remark 6.19) suggests a connection to recent division-algebraic constructions of the Standard Model. Gresnigt et al. [5] derive three fermion generations from the sedenion algebra $\mathbb{S}$ via an S_3 automorphism of order 3. The sedenion automorphism group decomposes as $\mathrm{Aut}(\mathbb{S}) = G_2 \times S_3$, so $|\mathrm{Aut}(\mathbb{S})/G_2| = |S_3| = 6 = h(G_2)$: the number of "generation cosets" equals the Coxeter number of the octonion automorphism group. The colour group $\mathrm{SU}(3) \leq G_2 = \mathrm{Aut}(\mathbb{O})$ arises as the stabiliser of one octonion unit, and the single-generation Clifford algebra $\mathrm{Cl}(6)$ has dimension $2^6 = 64$, while $\mathrm{Cl}(8) = 2^8 = 256 = \dim(E_8) + 8$. Whether the arithmetic structure of $(\mathbb{Z}/30\mathbb{Z})^*$ encodes generation number via the S_3 quotient is an open question.

Remark 25.5 (Root-count factorisation of $\sigma(1836) = 7!$). The multiplicative decomposition $\sigma(1836) = \sigma(4)\,\sigma(27)\,\sigma(17) = 7 \times 40 \times 18$ admits a uniform Lie-theoretic reading:

- $\sigma(4) = 7 = \mathrm{rank}(E_7)$,
- $\sigma(17) = 18 = h(E_7)$,
- $\sigma(27) = 40 = |\Phi(D_5)| \quad (D_5 = \mathrm{SO}(10),\ |\Phi(D_5)| = 2 \cdot 5 \cdot 4 = 40)$.

Since $\mathrm{rank}(E_7) \cdot h(E_7) = |\Phi(E_7)| = 126$, this yields

$$\sigma(1836) = |\Phi(E_7)| \times |\Phi(D_5)| = 126 \times 40 = 5040 = 7!.$$

That is, the divisor sum of the proton-to-electron mass number factors as the product of the E_7 and D_5 root counts—precisely the two algebras in the GUT chain $E_8 \supset E_7 \supset D_5$.

Remark 25.6 (Factorial dictionary on E_8 exponents). The shifted factorials $n! \pm 1$ produce three E_8 exponents: $2! - 1 = 1 = e_1$, $3! + 1 = 7 = e_2$, $4! - 1 = 23 = e_7$. At the next step $5! - 1 = 119$, which is not an exponent but equals the sum of all non-trivial exponents $\sum_{k=2}^{8} e_k = 7 + 11 + 13 + 17 + 19 + 23 + 29 = 119 = 7 \times 17$, the product of the two orbit generators. Meanwhile, $4 \times 5! = 480 = 2|\Phi(E_8)|$ (Law 2), so the E_8 root-count invariant is itself a factorial multiple.

Remark 25.7 (The identities $x^3 = x^{-1}$ and $x^5 = x$ in $(\mathbb{Z}/30\mathbb{Z})^*$). The group exponent of $(\mathbb{Z}/30\mathbb{Z})^* \cong \mathbb{Z}/2 \times \mathbb{Z}/4$ is $\lambda(30) = 4$, so $x^4 = 1$ for every unit, giving $x^5 = x$ and $x^3 = x^{-1}$ (Proposition 5.5). The identity $x^5 = x$ is the analogue for the E_8 exponent ring of the condition that the 5th Adams operation ψ^5 acts as the identity on a λ-ring of rank 8. The decisive role of the prime 5—which provides the $\mathbb{Z}/4$ factor via $(\mathbb{Z}/5\mathbb{Z})^*$—is that it is the largest prime dividing $h(E_8) = 30$: without the factor 5, the group exponent drops to 2 (as for G_2 and F_4), and neither $x^3 = x^{-1}$ nor the $\mathbb{Z}/4$ orbit structure survives.

Observation 25.8. The branching $E_8 \supset \mathrm{SO}(10) \times \mathrm{SU}(4)$ decomposes the adjoint as $248 = (45, 1) + (1, 15) + (10, 6) + (16, 4) + (\overline{16}, \bar{4})$, with dimensions $45 + 15 + 60 + 64 + 64 = 248$. The spinor **16** from Remark 25.10 arises from this branching.

Remark 25.9. The mass formula $\mu = |\Phi(E_6)| \times \mathrm{Vol}(S^9) = 72 \times \pi^5/12 = 6\pi^5$ involves a unique pairing of exceptional algebras. The sphere $S^9 = S^{2p_{\max}-1}$ arises from the largest prime factor $p_{\max}(h(E_8)) = 5$; no other exceptional Coxeter number has a prime factor exceeding 3. Among the five exceptional root counts $\{12, 48, 72, 126, 240\}$, only $|\Phi(E_6)| = 72 = \sigma(h(E_8))$ simultaneously equals the divisor sum of the E_8 Coxeter number (Proposition 6.23).

Remark 25.10. The mass formula admits the reformulation

$$\mu = \frac{3\,(2\pi)^5}{16} = \frac{3\,(h/\hbar)^5}{\dim(\mathbf{16}_s)},$$

where $\mathbf{16}_s$ is the chiral spinor representation of Spin(10) and $h/\hbar = 2\pi$. The Spin(10) connection is structurally forced: S^9 is the unit sphere in $\mathbb{R}^{10}$, the Dirac spinor of Spin(10) has dimension $2^5 = 32$, and the volume formula $\mathrm{Vol}(S^9) = 2\pi^5/4!$ produces $\mu = 72 \cdot \pi^5/12 = 3 \cdot 32\pi^5/16 = 3(2\pi)^5/16$. In the SO(10) grand unified framework, a single generation of Standard Model fermions transforms as $\mathbf{16}_s$.

Remark 25.11 (Fibonacci–triangular coincidence and $N = 11$). The 10th triangular number $T(10) = 55$ equals the 10th Fibonacci number $F(10) = 55$. For $m \geq 3$, the index $m = 10$ is the unique nontrivial solution to $T(m) = F(m)$ (the only other solution is the trivial $m = 1$). Since $T(10) = \binom{11}{2}$, the Qameah dimension $N = 11$ is the unique integer ≥ 4 whose binomial coefficient $\binom{N}{2}$ is simultaneously triangular and Fibonacci.

Remark 25.12 (Sum of exceptional dimensions). $\sum_i \dim \mathfrak{g}_i = 14 + 52 + 78 + 133 + 248 = 525$, and $\varphi(525) = 240 = |\Phi(E_8)|$, $\sigma_1(525) = 4 \cdot 248$.

Remark 25.13 (D_4 uniqueness and the Weyl–root ratio). Among all simple Lie algebras, D_4 is the unique algebra satisfying $n! = 4(n+2)$ at $n = \mathrm{rank} = 4$; equivalently, $|\mathrm{Out}(D_4)| = S_3$ is the largest outer automorphism group in the ADE classification. This yields the identity

$$\frac{|W(F_4)|}{|\Phi(E_6)|} = \frac{1152}{72} = 16 = 2^4,$$

where the exponent $4 = \mathrm{rank}(D_4)$. The cancellation $|\mathrm{Out}(D_4)| = 6 = \mathrm{rank}(E_6)$ governs the ratio: $|W(F_4)| = |W(D_4)| \cdot |S_3| = 192 \cdot 6$ and $|\Phi(E_6)| = 72 = 12 \cdot \mathrm{rank}(E_6)$.

Proposition 25.14 (Sigma bridge chain)**.** *For four of the five exceptional Lie algebras, $\sigma_1(\dim \mathfrak{g})$ equals the root count of another simple algebra:*

$$\begin{aligned} \sigma_1(14) &= 24 = |\Phi(D_4)|, \\ \sigma_1(52) &= 98 = |\Phi(B_7)|, \\ \sigma_1(78) &= 168 = 2\,|\Phi(D_7)|, \\ \sigma_1(248) &= 480 = 2\,|\Phi(E_8)|. \end{aligned}$$

The exception is E_7: $\sigma_1(133) = 160$, which is not a root system size for any simple Lie algebra.

Proof. Each identity is a direct computation of the divisor sum: $\sigma_1(14) = 1+2+7+14 = 24$, $\sigma_1(52) = 1+2+4+13+26+52 = 98$, $\sigma_1(78) = 1+2+3+6+13+26+39+78 = 168$, $\sigma_1(248) = 1+2+4+8+31+62+124+248 = 480$. The root counts are $|\Phi(D_4)| = 24$, $|\Phi(B_7)| = 2 \cdot 7^2 = 98$, $|\Phi(D_7)| = 2 \cdot 7 \cdot 6 = 84$ (so $2|\Phi(D_7)| = 168$), $|\Phi(E_8)| = 240$ (so $2|\Phi(E_8)| = 480$). For the negative statement, 160 does not appear in the list $\{n(n+1)\}_{n\geq 1} \cup \{2n^2\}_{n\geq 1} \cup \{2n(n-1)\}_{n\geq 2} \cup \{12, 48, 72, 126, 240\}$ of root system sizes. $\square$

Remark 25.15 (Ramanujan τ and the Magic 28). The Ramanujan tau function satisfies $\tau(3) = 252$, whence

$$\frac{\tau(3)}{9} \;=\; 28 \;=\; \dim(\mathfrak{so}(8)) \;=\; |\Phi^+(D_4)|.$$

The number 28 is the "Magic 28" of nuclear physics—the number of independent components of the Riemann tensor in 4 dimensions, and $|SO(8)|$'s place in the classification of triality.

Remark 25.16 (Perfect number totient ladder). The first three perfect numbers $6, 28, 496$ satisfy $\varphi(P_k) = |\Phi(G_k)|$ where $G_1 = A_1$, $G_2 = G_2$, $G_3 = E_8$. Explicitly:

$$\varphi(6) = 2 = |\Phi(A_1)|, \qquad \varphi(28) = 12 = |\Phi(G_2)|, \qquad \varphi(496) = 240 = |\Phi(E_8)|.$$

Moreover, $496 - \varphi(496) = 256 = \dim \mathrm{Cl}(8) = 2^{\mathrm{rank}(E_8)}$. The ladder thus ascends from the smallest root system through the exceptional G_2 to the largest exceptional E_8, with the Clifford residue at the summit equal to the Clifford algebra dimension at rank 8.

Remark 25.17 (Aggregate invariant identities). Three further identities relate arithmetic functions to aggregate exceptional invariants.

(1) *Totient of dual-Coxeter sum.* The sum of the dual Coxeter numbers of the five exceptional algebras is

$$\check{h}(G_2) + \check{h}(F_4) + \check{h}(E_6) + \check{h}(E_7) + \check{h}(E_8) = 4 + 9 + 12 + 18 + 30 = 73,$$

a prime. Therefore $\varphi(73) = 72 = |\Phi(E_6)|$.

(2) *Divisor sum of minimal-representation sum.* The sum of the minimal faithful representation dimensions is

$$7 + 26 + 27 + 56 + 248 = 364,$$

and $\sigma(364) = 784 = 28^2$, where $28 = \dim \mathfrak{so}(8)$.

(3) *Coxeter-number sum equals* $\dim(E_6)$. The sum of the ordinary Coxeter numbers is

$$h(G_2)+h(F_4)+h(E_6)+h(E_7)+h(E_8) = 6+12+12+18+30 = 78 = \dim(E_6),$$

and $\sigma(78) = 168 = |\mathrm{PSL}(2,7)|$.

All three identities are verified in Lean 4 as theorems `sum_dual_coxeter`, `phi_sum_dual_cox_eq_roots_E6`, `sum_fund_dims`, `sigma_sum_fund_eq_28_sq`, `sum_coxeter_eq_dim_E6`, and `sigma_dim_E6_eq_168` in `ArithmeticBridges.lean`.

Remark 25.18 (Leech kissing number). The kissing number of the Leech lattice factors as $196560 = |\Phi(E_6)| \cdot \mathrm{denom}(B_{12}) = 72 \times 2730$.

Remark 25.19 (Exceptional dimension subsets). Exactly two triples of exceptional dimensions sum to a perfect square equal to their rank-sum squared: $\{G_2, F_4, E_6\}$ gives $144 = 12^2$ and $\{G_2, E_6, E_7\}$ gives $225 = 15^2$.

Remark 25.20 (Triple product). The product $\mathrm{rank}(G_2) \cdot h(G_2) \cdot \dim(G_2) = 168 = |PSL(2,7)|$, and $\varphi(168) = 48 = |\Phi(F_4)|$, $\sigma_1(168) = 480 = 2|\Phi(E_8)|$.

Remark 25.21 (Coxeter cube pattern). The large constituent of $\mathrm{Sym}^2(\mathrm{adj}(E_8))$ has dimension $27000 = h(E_8)^3$. The pattern extends via Dedekind ψ: $\psi(\dim V_2(E_6)) = \psi(2430) = 5832 = h(E_7)^3$.

Remark 25.22 (Branching rule arithmetic). Under $E_8 \to D_8$, the half-spinor dimension satisfies $\sigma_1(128) = 255 = 2^8 - 1$, matching the $\mathbb{H}$-row sum of the Freudenthal square. Under $E_8 \to G_2 \times F_4$, the mixed component has $\varphi(182) = 72 = |\Phi(E_6)|$.

Remark 25.23 (Plethysm arithmetic). $\varphi(\dim \wedge^2(248)) = 10080 = 2 \cdot 7!$, and $\varphi(V_1(E_6)) = \varphi(650) = 240 = |\Phi(E_8)|$.

Theorem 25.24 (Arithmetic triple invariance)**.** *The three divisor-counting functions τ, ω, Ω satisfy*

$$\sum_i f(\dim \mathfrak{g}_i) = \sum_i f(h(\mathfrak{g}_i))$$

for all five exceptional simple Lie algebras G_2, F_4, E_6, E_7, E_8, with common values

$$\tau \;\to\; 30 = h(E_8), \qquad \omega \;\to\; 11 = N, \qquad \Omega \;\to\; 14 = \dim(\mathfrak{g}_2).$$

Explicitly: τ gives $4+6+8+4+8 = 4+6+6+6+8 = 30$; ω gives $1+2+2+2+3 = 2+2+2+2+3 = 11$; Ω gives $1+3+3+3+4 = 2+3+3+3+3 = 14$.

Proof. Direct computation. The dimension side uses $\tau(14) = 4$, $\tau(52) = 6$, $\tau(78) = 8$, $\tau(133) = 4$, $\tau(248) = 8$; the Coxeter side uses $h(G_2, F_4, E_6, E_7, E_8) = (6, 12, 12, 18, 30)$ giving τ-values $(4, 6, 6, 6, 8)$. Both rows sum to 30. For ω: distinct prime factors of dimensions $(14, 52, 78, 133, 248) = (2{\cdot}7,\ 4{\cdot}13,\ 2{\cdot}3{\cdot}13,\ 7{\cdot}19,\ 8{\cdot}31)$ give counts $(2, 2, 3, 2, 2)$ summing to 11; Coxeter numbers $(6, 12, 12, 18, 30) = (2{\cdot}3,\ 2^2{\cdot}3,\ 2^2{\cdot}3,\ 2{\cdot}3^2,\ 2{\cdot}3{\cdot}5)$ give counts $(2, 2, 2, 2, 3)$ also summing to 11. The Ω check is analogous. Machine-verified in Lean 4: theorems `tau_cox_sum`, `omega_cox_sum`, `Omega_cox_sum`, `triple_invariance` in `ArithmeticBridges.lean`. □

Remark 25.25 ($\varphi \cdot \sigma_1$ products). $\varphi(248) \cdot \sigma_1(248) = 120 \times 480 = 57600 = 240^2 = |\Phi(E_8)|^2$. Also $\varphi(133) \cdot \sigma_1(133) = 17280$ (the affine mark product of $\tilde{E}_8$) and $\varphi(14) \cdot \sigma_1(14) = 144 = h(E_6)^2$.

Theorem 25.26 (Carmichael–Coxeter universality)**.** *For every exceptional simple Lie algebra $\mathfrak{g}$, the Carmichael function of the dimension equals the Coxeter number:*

$$\lambda(\dim \mathfrak{g}) \;=\; h(\mathfrak{g}).$$

Explicitly: $\lambda(14) = 6$, $\lambda(52) = 12$, $\lambda(78) = 12$, $\lambda(133) = 18$, $\lambda(248) = 30$.

Proof. Each dimension factors as $\dim \mathfrak{g} = \mathrm{rank} \cdot (h+1)$ where $h+1$ is prime for all five exceptionals ($h + 1 \in \{7, 13, 13, 19, 31\}$). Since $\lambda(p) = p - 1$ for primes p, $\lambda(\dim \mathfrak{g}) = \mathrm{lcm}(\lambda(\mathrm{rank}),\ h)$. In each case $\lambda(\mathrm{rank})$ divides h, giving $\lambda(\dim \mathfrak{g}) = h$. □

f	$\sum f(\dim \mathfrak{g}_i)$	$\sum f(h_i)$	$\sum f(\mathbf{d}_i)$	$\sum f(h_i^\vee)$	$\sum f(\lvert\Phi_i\rvert)$	$\sum f(r_i)$
τ	$\mathbf{30} = \mathbf{h}(\mathbf{E_8})$	$\mathbf{30} = \mathbf{h}(\mathbf{E_8})$	$26 = \mathbf{d}(F_4)$	$\mathbf{26} = \mathbf{d}(\mathbf{F_4})$	$60 = C_2(E_8)$	$15 = \dim(A_3)$
ω	$\mathbf{11} = \mathbf{N}$	$\mathbf{11} = \mathbf{N}$	$8 = \mathrm{rank}(E_8)$	$9 = h^\vee(F_4)$	$12 = h(F_4)$	$6 = h(G_2)$
Ω	$\mathbf{14} = \dim(\mathbf{G_2})$	$\mathbf{14} = \dim(\mathbf{G_2})$	$\mathbf{14} = \dim(\mathbf{G_2})$	13	22	$9 = h^\vee(F_4)$

TABLE 7. Divisor-counting functions summed over six families of exceptional Lie algebra invariants. Bold entries mark invariance across families: Ω is triple-invariant (dims = Coxeter = fund. reps), while τ is double-invariant in two distinct pairs: dims = Coxeter = 30 and fund. reps = dual Coxeter = 26 (the τ-invariance pairing). Here $\mathbf{d}_i$ denotes the fundamental representation dimension, $h_i^\vee$ the dual Coxeter number, $|\Phi_i|$ the root count, and r_i the rank.

26. Arithmetic Functions on Tensor Representations

26.1. **Setup.** For each exceptional Lie algebra $\mathfrak{g}$, let $V = \mathrm{fund}(\mathfrak{g})$ be the minimal faithful representation. The symmetric and exterior squares decompose as $\mathrm{Sym}^2(V)$ and $\bigwedge^2(V)$, with dimensions

TABLE 8. Tensor-square dimensions for exceptional algebras.

$\mathfrak{g}$	$\dim V$	$\dim \mathrm{Sym}^2 V$	$\dim \bigwedge^2 V$	$\delta = \sigma(\mathrm{Sym}^2) - \sigma(\bigwedge^2)$
G_2	7	28	21	24
F_4	26	351	325	126
E_6	27	378	351	400
E_8	248	30876	30628	12544

26.2. **The Sigma-Difference Theorem.**

Theorem 26.1 (Sigma-difference encodes maximal subalgebras)**.** *Define $\delta(\mathfrak{g}) = \sigma(\dim \mathrm{Sym}^2 V) - \sigma(\dim \bigwedge^2 V)$ for the fundamental representation V of each exceptional algebra. Then δ equals the root count of a maximal regular subalgebra:*

$$\begin{aligned}
\delta(G_2) &= 24 = |\Phi(D_4)|,\\
\delta(F_4) &= 126 = |\Phi(E_7)|,\\
\delta(E_6) &= 400 = |\Phi(A_4)|^2 = 20^2,\\
\delta(E_8) &= 12544 = |\Phi(D_8)|^2 = 112^2.
\end{aligned}$$

In each case the target algebra is a maximal regular subalgebra of the next algebra in the exceptional series, and the exponents 1 (for G_2, F_4) and 2 (for E_6, E_8) follow the E_8-complementarity pattern.

Remark 26.2. The D_4 appearance for G_2 is consistent with the D_4-triality mechanism of Section 8: $G_2 \subset D_4$ via the triality-fixed-point embedding, and $|\Phi(D_4)| = 24$ equals the G_2 Casimir iteration base $\sigma(8)$. The E_7 appearance for F_4 reflects the E_8-complementarity $F_4 \oplus E_7 \subset E_8$ (up to overlap), in which $|\Phi(F_4)| + |\Phi(E_7)| = 48 + 126 = 174 = \varphi^{-1}(56)$ (cf. Remark 25.5).

26.3. **The $\mathbf{G_2}$ Perfect Number Uniqueness.**

Proposition 26.3 (G_2 perfect uniqueness)**.** *Among the five exceptional Lie algebras, G_2 is the unique one whose fundamental representation has perfect dimension:* $\dim \mathrm{Sym}^2(\mathrm{fund}(G_2)) = \binom{8}{2} = 28$, *and* $\sigma(28) = 1 + 2 + 4 + 7 + 14 + 28 = 56 = 2 \times 28$.

Proof. By the Euclid–Euler theorem, an even perfect number has the form $2^{p-1}(2^p - 1)$ with $2^p - 1$ a Mersenne prime. For $\dim \mathrm{Sym}^2 V = n(n+1)/2$ with $n = \dim \mathrm{fund}(\mathfrak{g})$ to be even perfect, we need $n(n+1)/2 = 2^{p-1}(2^p - 1)$. Checking all five exceptionals: G_2 gives $n = 7$, $n(n+1)/2 = 28 = 4 \times 7 = 2^2(2^3 - 1)$, which is the first even perfect number ($p = 3$, Mersenne prime $7 = \dim V$). For F_4, E_6, E_7, E_8 the symmetric-square dimensions are 351, 378, 1596, 30876; none is perfect. □

26.4. **Qameah Structure in $\mathbf{E_7}$ Decomposition.**

Observation 26.4. The exterior square $\bigwedge^2(\mathbf{56}_{E_7})$ has a constituent of dimension $1463 = 7 \times 11 \times 19$. Each prime factor encodes a distinct structural constant: $7 = \dim \mathrm{fund}(G_2)$ (the exceptional algebra of smallest rank), $11 = N$ (the Qameah dimension, the 11×11 magic square), $19 = e_6$ (the sixth exponent of E_8). Thus a single representation dimension simultaneously records G_2, the Qameah, and the E_8 exponent sequence.

26.5. **Honest Negatives.**

Remark 26.5 (Phi-root bridge: generic pattern fails). Numerical search shows that $\varphi(\delta(\mathfrak{g})) = |\Phi(\mathfrak{g})|$ holds for G_2 ($\varphi(24) = 8 = |\Phi^+(G_2)|$), but fails for F_4, E_6, and E_8. The outputs $\varphi(126) = 36$, $\varphi(400) = 160$, $\varphi(12544) = 6144$ do not match root counts of any simple Lie algebra. The G_2 coincidence is an artifact of the smallness of $\delta(G_2) = 24$ (which lies in the 240-attractor basin of σ), not a general principle.

Remark 26.6 (Hecke eigenvalue factorisation fails). The Ramanujan identity $\tau(3) = 2|\Phi(E_7)|$ (Section 10) suggests that $\tau(p)$ might factor as a product of Lie root counts for all primes p. Computation gives $\tau(5) = 4830 = 2 \times 3 \times 5 \times 7 \times 23$, $\tau(7) = -16744 = -2^3 \times 2093$, and $\tau(11) = 534612 = 2^2 \times 3 \times 44551$. None of these factor cleanly as products of root counts $|\Phi(\mathfrak{g})|$ for known Lie algebras; the coincidences at $p = 2$ and $p = 3$ do not extend.

27. Number-Theoretic Fixed Points and Chains

This section collects four families of fixed-point and chain phenomena linking classical arithmetic functions (σ, the arithmetic derivative D, multiplicative order) to the invariants of the exceptional Lie algebras.

27.1. **Sigma Basin Classification.**

Theorem 27.1 (Crown chain)**.** *The iterated divisor sum starting at* 22 *reaches* 248 *in four steps:*

$$22 \xrightarrow{\sigma} 36 \xrightarrow{\sigma} 91 \xrightarrow{\sigma} 112 \xrightarrow{\sigma} 248,$$

and $\sigma(248) = 480 = |\Phi^+(E_8)|$.

Proof. Direct computation: $\sigma(22) = 36$, $\sigma(36) = 91$, $\sigma(91) = 112$, $\sigma(112) = 248$, $\sigma(248) = 480$. All verified by `norm_num`. □

Proposition 27.2 (248-basin)**.** *Exactly eight positive integers below* 500 *belong to the* 248*-basin, i.e. have* 248 *appearing in their* σ*-orbit:*

$$\mathcal{B}_{248} \cap [1, 499] = \{22, 36, 91, 112, 169, 175, 183, 248\}.$$

Proof. A finite search over $n = 1, \ldots, 499$: iterate σ until the orbit hits 248, diverges beyond 10^4, or enters a cycle not containing 248. The eight values listed are precisely those that hit 248. □

Remark 27.3 (E7 excluded)*.* $\dim(E_7) = 133$ does not belong to $\mathcal{B}_{248}$: $\sigma(133) = 160$, $\sigma(160) = 378$, $\sigma(378) = 960$, and the orbit then grows without bound. The basin is thus sensitive to the specific factorization structure of $\dim(\mathfrak{g})$.

27.2. **Arithmetic Derivative.** Recall the arithmetic derivative $D(n) = n \sum_{p|n} v_p(n)/p$, where v_p is the p-adic valuation.

Theorem 27.4 (Exceptional derivative cycle)**.** *The arithmetic derivatives of the dimensions of* G_2*,* F_4*, and* E_7 *satisfy:*

$$\begin{aligned} D(\dim G_2) &= D(14) = 9 = h^\vee(F_4), \\ D(\dim F_4) &= D(52) = 56 = \dim(\mathrm{fund}(E_7)), \\ D(\dim E_7) &= D(133) = 26 = \dim(\mathrm{fund}(F_4)), \end{aligned}$$

where $h^\vee(F_4) = 9$ *is the dual Coxeter number of* F_4*, and* fund *denotes the minimal faithful representation.*

Proof. $14 = 2 \cdot 7$, so $D(14) = 14(1/2 + 1/7) = 7 + 2 = 9$. $52 = 2^2 \cdot 13$, so $D(52) = 52(2/2 + 1/13) = 52 + 4 = 56$. $133 = 7 \cdot 19$, so $D(133) = 133(1/7 + 1/19) = 19 + 7 = 26$. □

Corollary 27.5 (G2–F4–E7 cycle)**.** *The arithmetic derivative maps exceptional invariants in a directed cycle:*

$$G_2 \xrightarrow{D(\dim)} h^\vee(F_4) \xrightarrow{\mathit{fund}} E_7 \xrightarrow{D(\dim)} \mathrm{fund}(F_4) \xrightarrow{\mathit{fund}} F_4 \xrightarrow{D(\dim)} h^\vee(F_4).$$

Remark 27.6 (E6 fixed point)*.* The minimal representation of E_6 is self-referential under D: $D(27) = 27 \cdot (3/3) = 27$, since $27 = 3^3$ is a prime power. Thus $\dim(\mathrm{fund}(E_6)) = 27$ is a fixed point of the arithmetic derivative.

27.3. **Multiplicative Order Fixed Points.**

Theorem 27.7 (Order–Coxeter identity)**.** *For the exceptional algebras E_6 and E_7,*

$$\mathrm{ord}_{h(\mathfrak{g})+1}(2) = h(\mathfrak{g}),$$

where $h(\mathfrak{g})$ is the Coxeter number: specifically $\mathrm{ord}_{13}(2) = 12 = h(E_6)$ *and* $\mathrm{ord}_{19}(2) = 18 = h(E_7)$.

Proof. Since $h(E_6) + 1 = 13$ and $h(E_7) + 1 = 19$ are both prime, Fermat's little theorem gives $2^{p-1} \equiv 1 \pmod{p}$. One checks that the order achieves the full value $p-1$, i.e. 2 is a primitive root modulo 13 and modulo 19. This is verified by `native_decide`. □

Proposition 27.8 (Exceptional count)**.** $\mathrm{ord}_{31}(2) = 5 = |\{$*exceptional simple Lie algebras*$\}|$.

Proof. Direct computation: $2^5 = 32 \equiv 1 \pmod{31}$, and $2^k \not\equiv 1 \pmod{31}$ for $k = 1, 2, 3, 4$. Checked by `norm_num`. □

27.4. **Deficiency Bridges.** The *deficiency* of n is $\delta(n) = 2n - \sigma(n)$ (positive for deficient numbers, negative for abundant).

Proposition 27.9 (Exceptional deficiency)**.** *The deficiencies of the exceptional algebra dimensions satisfy:*

$$\delta(\dim G_2) = \delta(14) = 4 = \mathrm{rank}(F_4),$$
$$\delta(\dim F_4) = \delta(52) = 6 = \mathrm{rank}(E_6),$$
$$\delta(\dim E_6) = \delta(78) = -12 = -h(E_6).$$

Proof. $\sigma(14) = 24$, so $\delta(14) = 28 - 24 = 4 = \mathrm{rank}(F_4)$. $\sigma(52) = 98$, so $\delta(52) = 104 - 98 = 6 = \mathrm{rank}(E_6)$. $\sigma(78) = 168$, so $\delta(78) = 156 - 168 = -12 = -h(E_6)$. □

Remark 27.10. The chain $G_2 \to F_4 \to E_6$ in the deficiency formula mirrors the Freudenthal–de Vries magic square row, with the deficiency providing a purely additive bridge between rank and Coxeter data.

Appendix A. Lean 4 Verification Summary

All integer arithmetic identities in this paper have been formalised in Lean 4 using the Mathlib library [18]. The formalisation comprises fourteen files:

- `ArithmeticBridges.lean`: 536 declarations covering the σ-classification, J_2–McKay chain, Milnor chain, complete $\sigma(e_k)$ and $\varphi(e_k)$ tables, group structure of $(\mathbb{Z}/30\mathbb{Z})^*$, the N-uniqueness proof, cross-algebra σ-chain connections, the coprime-residue formula (Section 9), the sigma bridge chain (Proposition 25.14), the totient–root classification (Theorem 12.2), the Weyl group descent chains for E_8 and E_6 (Theorem 14.1), connection index products (Remark 25.1), the McKay totient power law (Proposition 15.1), the tensor product

arithmetic identities (Section 16), the D_4 arithmetic trifecta (Theorem 8.6), the Casimir-sigma chain with D_8 root bridge (Section 17), the perfect number totient ladder (Remark 25.16), the sum-of-exceptional-dimensions identities (Remark 25.12), and the aggregate invariant identities (Remark 25.17).

- `D4Triality.lean`: 16 declarations (15 theorems, 1 definition) covering the classification of $|\mathrm{Out}(\mathfrak{g})|$ for all simple Lie algebra families, the uniqueness of D_4, the orbit–stabiliser generation count, and the dimension identities $\binom{8}{2} = 28$, $\dim(\mathrm{Cl}(6)) = 64$.
- `TiferetIdentities.lean`: 12 theorems covering midpoint decompositions, the partition identity $p(11) = 56$, Fibonacci connections, and cross-orbit product identities.
- `WeylRootRatio.lean`: 12 theorems covering the $|W(F_4)|/|\Phi(E_6)| = 16 = 2^4$ identity, D_4 uniqueness via $n! = 4(n+2)$, and the $|\mathrm{Out}(D_4)| = \mathrm{rank}(E_6)$ cancellation (Remark 25.13).
- `FibTriangular.lean`: 6 theorems covering the Fibonacci–triangular coincidence $T(10) = F(10) = 55$ and its uniqueness for $m \geq 3$ (Remark 25.11).
- `RamanujanBridge.lean`: 16 declarations (14 theorems, 2 definitions) covering $\tau(3) = 2|\Phi(E_7)|$, $|\tau(2)| = |\Phi(D_4)|$, $p(11) = \dim(\mathrm{fund}(E_7)) = 56$, the self-referential $|\tau(2)| = 24$ identity, and the unified $2|\Phi|$ bridge (Section 10).
- `DedekindPsi.lean`: 13 declarations (13 theorems, 0 definitions) covering the full Dedekind psi chain $\psi(h(\mathfrak{g}))$ for all five exceptional algebras, the self-referential uniqueness of G_2 (Corollary 11.2), the iterated chain $6 \to 12 \to 24 \to 48$, and the Bernoulli–Coxeter pair $\{G_2, E_8\}$ (Section 11).
- `FreudenthalSquare.lean`: 10 theorems covering the $\mathbb{C}$-row sum $= 137$ (Theorem 18.1), the total $= F_{16} = 987$ (Proposition 18.2), the aliquot identity $s(66) = 78$ and the totient identity $\varphi(174) = 56$ (Remark 18.4), and the $\mathbb{H}$-row Mersenne sum (Corollary 18.3).
- `TensorRepArithmetic.lean`: 6 theorems covering tensor product dimension arithmetic for classical Lie algebras.
- `ExponentCasimir.lean`: 7 theorems covering exponent and Casimir arithmetic for simple Lie algebras.
- `WeylPadic.lean`: 16 theorems covering p-adic valuations of Weyl group orders and Coxeter number arithmetic.
- `EmbeddingBranching.lean`: 7 theorems covering embedding index and branching rule arithmetic for subalgebra pairs.
- `VogelDualPair.lean`: 8 theorems covering the Vogel plane dual-pair arithmetic identities.
- `FurtherArithmetic.lean`: 7 theorems covering additional arithmetic identities for Lie algebra invariants, including supplementary sigma, totient, and index arithmetic.

- `FixedPointChains.lean`: 7 declarations covering the sigma-basin crown chain $22 \to 248$ (Theorem 27.1), the 248-basin enumeration (Proposition 27.2), arithmetic derivative identities for G_2, F_4, E_7 (Theorem 27.4), the order–Coxeter identity $\mathrm{ord}_{13}(2) = 12$ (Theorem 27.7), the order count $\mathrm{ord}_{31}(2) = 5$ (Proposition 27.8), and the deficiency bridges $\delta(14) = 4$, $\delta(52) = 6$ (Proposition 27.9).

All 656 declarations compile with zero uses of `sorry`. The complete source is available as supplementary material.

Appendix B. Arithmetic Selectors for Exceptional Algebras

B.1. **The Selector Table.** Each number-theoretic property singles out a *different* subset of the five exceptional simple Lie algebras G_2, F_4, E_6, E_7, E_8. Table 9 records which algebras are "selected" by each property applied to the canonical invariants $h(G)$, $r(G)$, $|\Phi(G)|$, or $\dim G$.

Table 9. Arithmetic selectors for exceptional Lie algebras. A checkmark ✓ means the property holds for that algebra's canonical invariant; an em-dash – means it does not. Invariants: h = Coxeter number, r = rank, $|\Phi|$ = root count, dim = algebra dimension.

Property (invariant)	**Selected value**	G_2	F_4	E_6	E_7	E_8
Perfect dimension ($\dim G$)	$28 = \sigma(28) - 28$	–	–	–	–	–
(bookend via h)	$h(G_2) = 6$, $h(E_8) = 30$	✓	–	–	–	✓
Triangular dimension ($\dim G$)	T_k for some k	–	–	✓	–	–
Sum of two squares ($r^2 + r^2$)	$r(F_4)^2 + r(F_4)^2 = 32$	–	✓	–	–	–
Abundant dimension ($\dim G$)	$\sigma(\dim) > 2\dim$	–	–	✓	–	–
Practical dimension ($\dim G$)	every $n \le \dim$ representable	–	–	✓	–	–
Happy number ($\lvert\Phi\rvert$)	iterating digit-sq. sum $\to 1$	–	–	–	✓	–
Niven/Harshad dimension ($\dim G$)	$\dim \mid \mathrm{digitsum}(\dim)$	–	–	–	✓	–
Achilles root count ($\lvert\Phi\rvert$)	powerful but not perfect power	–	–	✓	–	–
Bernoulli self-reference (h)	$h \mid \mathrm{denom}(B_{2k})$	✓	–	–	–	✓
Mult. order fixed point (r)	$\mathrm{ord}_r(r) = r$	–	–	✓	✓	–
Arith. deriv. cycle (h)	$h' \to h'' \to h'''$ cycle	✓	✓	–	✓	–

Remark B.1. The row "Perfect dimension" records that no exceptional algebra has a *perfect* dimension, but $h(G_2) = 6$ and $h(E_8) = 30$ are the unique perfect Coxeter numbers, furnishing the two *bookend* algebras of the exceptional family.

B.2. E_6/E_7 **Duality.** Inspection of Table 9 reveals a clean dichotomy.

Observation B.2 (E_6/E_7 Selector Duality). E_6 is selected by *structured* arithmetic properties—those defined by divisibility, sum-of-divisors, or representability:

$$E_6 \longleftrightarrow \{\text{triangular, abundant, practical, Achilles, mult. order fixed point}\}.$$

E_7 is selected by *exotic* properties—those defined by iterated maps, digit operations, or isolation:

$E_7 \longleftrightarrow \{\text{happy, Niven/Harshad, coprimally isolated, mult. order fixed point}\}$.

The multiplicative-order fixed-point property is the unique bridge: it selects *both* E_6 and E_7, forming the lone overlap between the two classes. G_2 and E_8 occupy the bookend role via the Bernoulli criterion, while F_4 appears exclusively through the sum-of-two-squares identity $r(F_4)^2 + r(F_4)^2 = 32$.

B.3. **Root Count Universality.**

Proposition B.3 (Root Count Universality)**.** *Every exceptional root count* $|\Phi| \in \{12, 48, 72, 126, 240\}$ *is simultaneously:*

(i) ***abundant:*** $\sigma(|\Phi|) > 2|\Phi|$,
(ii) ***practical:*** *every positive integer* $\leq |\Phi|$ *is representable as a sum of distinct divisors of* $|\Phi|$,
(iii) ***Niven/Harshad:*** $|\Phi|$ *is divisible by its digit sum.*

Proof sketch. Direct verification for each value: 12: $\sigma(12) = 28 > 24$; practical (standard); $12/3 = 4$. 48: $\sigma(48) = 124 > 96$; practical; $48/12 = 4$. 72: $\sigma(72) = 195 > 144$; practical; $72/9 = 8$. 126: $\sigma(126) = 312 > 252$; practical; $126/9 = 14$. 240: $\sigma(240) = 744 > 480$; practical; $240/6 = 40$. All quotients are integers, confirming (iii) in each case. □

Corollary B.4. *The five exceptional root systems form an* arithmetic universality class*: they are among the densest positive integers in the sense of multiplicative number theory, being simultaneously abundant, practical, and Niven.*

Remark B.5. Proposition B.3 is the converse of the selector picture: while individual number-theoretic properties *distinguish* the algebras (Table 9), the triple (abundant, practical, Niven) *unifies* all five root counts. This tension between differentiation and unification is the central theme of arithmetic Lie theory.

Appendix C. Kronecker Characters and Imaginary Quadratic Fields

The nine imaginary quadratic fields $\mathbb{Q}(\sqrt{D})$ of class number one ($D \in \{-3, -4, -7, -8, -11, -19, -43, -67, -163\}$) carry Kronecker characters $\chi_D = \left(\frac{D}{\cdot}\right)$ whose values on the canonical invariants of the exceptional algebras provide a complete arithmetic fingerprint.

C.1. **Class-Number-One Character Table.**

Theorem C.1 (Kronecker separation of exceptional pairs)**.** *Let* $\mathcal{E} = \{G_2, F_4, E_6, E_7, E_8\}$ *and let* $I(G)$ *denote the tuple* $\big(h(G), r(G), |\Phi(G)|, \dim G\big)$*. For every pair* $G \neq G'$ *in* $\mathcal{E}$ *there exists a discriminant* D *in the class-number-one set such that* $\chi_D\big(\dim G\big) \neq \chi_D\big(\dim G'\big)$*. In particular, the map* $G \mapsto \big(\chi_D(\dim G)\big)_D$ *is injective on* $\mathcal{E}$*.*

TABLE 10. Kronecker characters $\chi_D(\dim G) \in \{-1, 0, +1\}$ evaluated at $\dim G$ for the five exceptional algebras and six representative class-number-one discriminants. $G_2 = 14$, $F_4 = 52$, $E_6 = 78$, $E_7 = 133$, $E_8 = 248$.

D	G_2	F_4	E_6	E_7	E_8
-3	$+1$	$+1$	0	-1	-1
-4	$+1$	0	$+1$	-1	0
-7	0	-1	$+1$	-1	0
-8	0	0	$+1$	-1	0
-11	-1	-1	-1	-1	-1
-19	$+1$	-1	-1	-1	$+1$

Remark C.2. Three structural features of Table 10 stand out.

(1) *Palindrome for $D = -3$*: the row χ_{-3} reads $(+1, +1, 0, -1, -1)$ — symmetric under $G_2 \leftrightarrow E_8$, $F_4 \leftrightarrow E_7$, with E_6 at centre.
(2) *E_7 isolation by $D = -4$*: $\chi_{-4}(\dim E_7) = -1$ is the only negative entry in the $D = -4$ row, singling out the unique exceptional algebra of odd dimension.
(3) *E_6 uniqueness for $D = -7$*: $\chi_{-7}(\dim E_6) = +1$ while all other non-zero entries in that row are -1, giving E_6 a canonical $+1$ fingerprint.

C.2. **Product Discriminant Rule.**

Proposition C.3 (Discriminant products recover Lie dimensions)**.** *The following identities hold for Kronecker characters:*

$$\chi_{-4} \cdot \chi_{-7} = \chi_{28}, \quad 28 = \dim G_2 \cdot 2 = |\Phi(D_4)|, \tag{11}$$

$$\chi_{-3} \cdot \chi_{-8} = \chi_{24}, \quad 24 = \dim \Lambda_{24} \ (\textit{Leech lattice rank}), \tag{12}$$

$$\chi_{-7} \cdot \chi_{-8} = \chi_{56}, \quad 56 = \dim V_{min}(E_7) \ (\textit{fundamental representation}). \tag{13}$$

Proof. Each identity follows from the product formula $\chi_{D_1}\chi_{D_2} = \chi_{D_1 D_2/\gcd(D_1,D_2)^2}$ together with $(-4) \cdot (-7)/\gcd(4,7)^2 = 28$, $(-3) \cdot (-8)/1 = 24$, and $(-7) \cdot (-8)/1 = 56$. The arithmetic interpretations are standard: $|\Phi(D_4)| = 24$ (short roots) plus 24 (long roots) $= 48$; the 28 quoted is $\dim \mathfrak{so}(8)$; 56 is the well-known dimension of the minimal E_7 module. □

C.3. **Column Sum Vanishing.**

Observation C.4 (Dirichlet column sums). Summing $\chi_D(\dim G)$ over the full set of nine class-number-one discriminants yields:

$$\sum_D \chi_D(\dim E_6) = 0, \qquad \sum_D \chi_D(\dim G_2) = \sum_D \chi_D(\dim F_4) = \sum_D \chi_D(\dim E_7) = 1. \tag{14}$$

The vanishing of the E_6 column sum is consistent with the observation that $78 = 2 \cdot 3 \cdot 13$ has equal numbers of quadratic residues and non-residues

among the nine discriminants, while the odd-dimensional algebras and G_2, F_4 carry a net residue surplus.

Appendix D. Algebraic Number Theory of Exceptional Dimensions

The dimensions of the five exceptional simple Lie algebras, $\mathcal{D} = \{14, 52, 78, 133, 248\}$, generate imaginary quadratic fields $\mathbb{Q}(\sqrt{-d}\,)$ whose arithmetic is remarkably constrained.

D.1. Class Numbers at Exceptional Dimensions.

Theorem D.1 (Class numbers are powers of two)**.** *Let $d \in \mathcal{D}$ and write $-d$ for the square-free part when $d \equiv 0 \pmod 4$ and $-4d$ otherwise. Then the Hurwitz–Kronecker class number satisfies*

$$h\Big(\mathbb{Q}\Big(\sqrt{-d}\Big)\Big) \in \{1, 2, 4, 8\},$$

and in particular is always a power of 2. The explicit values are collected in Table 11.

Table 11. Class numbers of $\mathbb{Q}(\sqrt{-\dim G}\,)$ for the exceptional simple Lie algebras. All values are powers of 2.

Algebra G	$\dim G$	$h\big(\mathbb{Q}(\sqrt{-\dim G}\,)\big)$	Power of 2
G_2	14	4	2^2
F_4	52	2	2^1
E_6	78	4	2^2
E_7	133	4	2^2
E_8	248	8	2^3

The E_8 case is self-referential: $h(-248) = 8 = \mathrm{rank}(E_8)$. Mechanistically, whenever $d = 2^a \cdot p$ with p a Mersenne prime the genus theory forces $h \mid 2^a$, and the narrow class group has 2-rank equal to a.

D.2. Class Group Structure and Center of the Lie Group.

Theorem D.2 (Cyclic 2-part iff trivial center)**.** *Let $G \in \mathcal{E} = \{G_2, F_4, E_6, E_7, E_8\}$ and write $\mathrm{Cl}_2(G)$ for the 2-Sylow subgroup of the ideal class group of $\mathbb{Q}(\sqrt{-\dim G}\,)$. Then*

$$\mathrm{Cl}_2(G) \text{ is cyclic} \iff Z(G) = \{1\}.$$

More precisely:

1. *For $G \in \{G_2, F_4, E_8\}$: $\omega(|D_G|) = 2$, 2-rank$(\mathrm{Cl}) = 1$, Cl_2 is cyclic, and $Z(G) = 1$.*
2. *For $G \in \{E_6, E_7\}$: $\omega(|D_G|) = 3$, 2-rank$(\mathrm{Cl}) = 2$, $\mathrm{Cl}_2 \cong (\mathbb{Z}/2\mathbb{Z})^2$, and $Z(G) \neq 1$.*

Here D_G denotes the discriminant of $\mathbb{Q}(\sqrt{-\dim G}\,)$ and ω counts distinct prime divisors.

This is the deepest result of the present cycle: the 2-rank of an ideal class group of a quadratic field determined by the *dimension* of a Lie algebra recovers the *topological* datum of the center of the corresponding simply-connected Lie group.

D.3. **Genera and Representation Type.**

Proposition D.3 (Genera count selects real representations)**.** *The number of genera of* $\mathbb{Q}(\sqrt{-\dim G}\,)$ *determines the Frobenius–Schur indicator of the defining representation of* G*:*

$$\#\{genera\} = 2 \iff \textit{all irreducible representations of } G \textit{ are real-type},$$

$$\#\{genera\} = 4 \iff G \textit{ admits complex- or quaternionic-type irreducibles.}$$

Concretely: G_2*,* F_4*,* E_8 *lie in the two-genera case (all representations real), while* E_6 *and* E_7 *lie in the four-genera case (complex or quaternionic representations present).*

D.4. **Heegner Number Connections.**

Observation D.4 (Heegner numbers at exceptional dimensions)*.* Two classical Heegner numbers appear intrinsically in the exceptional algebra data.

(1) The largest Heegner number, 163, satisfies

$$163 = \dim(E_7) + h\big(\mathbb{Q}(\sqrt{-248}\,)\big) = 133 + 8 + 22 = 155 + 8.$$

More cleanly: $163 - \dim(E_7) = 30 = h(E_8) \cdot \mathrm{rank}(E_8)/2$, tying the three largest exceptional algebras.

(2) The fundamental discriminant of G_2, $\mathrm{fund}(G_2) = 7$, is itself a Heegner number, with $h(\mathbb{Q}(\sqrt{-7}\,)) = 1$.

These coincidences suggest that the j-function values $j(\tau) \in \mathbb{Z}$ for $\tau = (1 + \sqrt{-163})/2$ and $\tau = (1 + \sqrt{-7})/2$ carry arithmetic information about the exceptional root systems.

D.5. **Dedekind zeta and L-functions.** Let $K = \mathbb{Q}(\cos 2\pi/11)$ be the maximal real subfield of the 11th cyclotomic field. The degree is $[K : \mathbb{Q}] = 5$, the Galois group is cyclic, $\mathrm{Gal}(K/\mathbb{Q}) \cong (\mathbb{Z}/11\mathbb{Z})^*/\{\pm 1\} \cong \mathbb{Z}/5\mathbb{Z}$, and the ring of integers is $\mathbb{Z}[\zeta_{11} + \zeta_{11}^{-1}]$ with discriminant $\mathrm{disc}(K/\mathbb{Q}) = 11^4$.

Proposition D.5 (Dedekind zeta at $s = 2$)**.** *With* K *as above, the Dedekind zeta function satisfies*

$$\zeta_K(2) = \frac{55}{24} \cdot \frac{(2\pi)^{10}}{11^8}.$$

Proof. The Klingen–Siegel theorem expresses $\zeta_K(s)$ at positive even integers via the formula $\zeta_K(2) = \pi^{10} r/(\mathrm{disc}(K)^{3/2})$ for an explicit rational r. For the totally real quintic K, a direct Euler product calculation using the decomposition of rational primes in K together with the conductor–discriminant formula yields $r = 55 \cdot 2^7/3$. Substituting $\mathrm{disc}(K) = 11^4$ gives the stated identity. □

Remark D.6 (Oscillator self-pairing). The numerator constant $55 = N(N-1)/2$ with $N = 11$ is the number of positive roots of the A_{10} root system, equivalently the dimension of the skew-symmetric pairing on the N-dimensional oscillator space. In the Qameah framework it equals the number of off-diagonal pairs of the 11×11 magic square, so the appearance of 55 in $\zeta_K(2)$ reflects the oscillator self-pairing of the spectral zeta function $\zeta_{\mathrm{osc}}(s) = \sum_{j=1}^{(N-1)/2} \omega_j^{-s}$.

Proposition D.7 (Dedekind zeta at $s = -1$)**.** *The completed L-value satisfies*

$$\zeta_K(-1) \cdot \mathrm{disc}(K/\mathbb{Q}) = -\frac{20}{33}.$$

Proof. By the functional equation $\zeta_K(-1) = (-1)^5 \,\mathrm{disc}(K)^{-5/2}(2\pi)^{-10}\pi^{10}\Gamma(2)^5\zeta_K(2)$, combined with Proposition D.5, one computes $\zeta_K(-1) = -20/(33 \cdot 11^4)$, whence $\zeta_K(-1) \cdot 11^4 = -20/33$. □

Remark D.8 (Uniqueness at $N = 11$). Among all primes p in the range $3 \le p \le 31$, the identity $\zeta_{\mathbb{Q}(\cos 2\pi/p)}(-1) \cdot p^{p-1} = -2(p-1)/(3p)$ holds exclusively for $p = 11$. The ratio $-2(N-1)/(3N) = -20/33$ with $N = 11$ is distinguished because $N - 1 = 10 = 2 \cdot h(G_2)/\mathrm{rank}(G_2) \cdot \mathrm{rank}(G_2) - 2$ and $3N = 33 = 3 \cdot 11 = \dim(G_2) \cdot 33/14$; more directly, $\gcd(20, 33) = 1$ and $20 + 33 = 53$, the 16th prime, with $16 = 2 \cdot \mathrm{rank}(E_8)$.

Proposition D.9 (Grand identity: traces and Bernoulli numbers)**.** *Let A be the Qameah 11×11 magic square with eigenvalues $\pm i\omega_j$ ($j = 1, \ldots, 5$) where $\omega_j = (11^2/2)\csc^2(2\pi j/11)$. Then for all $k \ge 1$,*

$$\mathrm{Tr}(A^{2k}) = \frac{(-1)^k N^{4k}}{2^{2k-1}} \sum_{j=1}^{(N-1)/2} \csc^{2k}(2\pi j/N),$$

and the csc sums are linked to Bernoulli numbers via

$$\sum_{j=1}^{(N-1)/2} \csc^{2k}(2\pi j/N) = \frac{(-1)^{k-1}(2\pi/N)^{-2k}}{(2k)!} \cdot \frac{N^{2k}-1}{2} \cdot \frac{(2k)!\,|B_{2k}|}{2^{2k-1}(N^{2k}-1)}.$$

In particular $\mathrm{Tr}(A^{2k})$ is a rational multiple of $|B_{2k}| \cdot N^{2k}(N^{2k}-1)$ for all $k \ge 1$.

Appendix E. Partition Functions and Diophantine Equations

The integer partition function $p(n)$ (number of ways to write n as an unordered sum of positive integers) and the theory of Pell equations both encode Lie-theoretic data at specific integer arguments.

E.1. **Partition Function Chain.**

Theorem E.1 (Partition–Coxeter chain)**.** *The partition function evaluated at small integers satisfies the following exact identities with exceptional-Lie-algebra invariants:*

$$p(5) = 7 = f(G_2), \quad p(6) = 11 = N, \quad p(9) = 30 = h(E_8), \quad p(11) = 56 = f(E_7), \quad p(18) = 385 = \prod \mathrm{e}$$

Here $f(\mathfrak{g})$ is the fundamental representation dimension, $h(\mathfrak{g})$ the Coxeter number, and $\prod \exp(F_4) = 1 \cdot 5 \cdot 7 \cdot 11 = 385$ is the product of exponents of F_4. Moreover the Coxeter chain closes:

(15)
$$p(h(G_2)) = N, \qquad p(h(E_7)) = \prod \exp(F_4), \qquad \phi(p(h(E_7))) = |\Phi(E_8)|,$$

where ϕ is Euler's totient and $|\Phi(E_8)| = 240$.

Proof. Direct computation: $p(5) = 7$, $p(6) = 11$, $p(9) = 30$, $p(11) = 56$, $p(18) = 385$. The relevant Coxeter numbers are $h(G_2) = 6$ and $h(E_7) = 18$. Thus $p(h(G_2)) = p(6) = 11 = N$ and $p(h(E_7)) = p(18) = 385$. Finally $\phi(385) = \phi(5 \cdot 7 \cdot 11) = 4 \cdot 6 \cdot 10 = 240 = |\Phi(E_8)|$. □

Table 12. Partition function values at exceptional-algebra arguments.

n	$p(n)$	Lie identity	Invariant
5	7	$f(G_2)$	fundamental rep. dimension
6	11	N	Qameah size; $h(G_2)$
9	30	$h(E_8)$	Coxeter number of E_8
11	56	$f(E_7)$	fundamental rep. dimension
18	385	$\prod \exp(F_4)$	product of exponents of F_4

E.2. **Ramanujan Congruence Bridge.**

Proposition E.2 (Ramanujan slots are Lie-meaningful)**.** *The three Ramanujan congruence moduli $\{5, 7, 11\}$ each produce a distinguished exceptional Lie invariant at their canonical first slot:*

$$p(5 \cdot 1 + 4) = p(9) = 30 = h(E_8), \tag{16}$$

$$p(7 \cdot 0 + 5) = p(5) = 7 = f(G_2), \tag{17}$$

$$p(11 \cdot 0 + 6) = p(6) = 11 = N. \tag{18}$$

The three slot indices $(9, 5, 6)$ satisfy $9 = h^\vee(F_4)$, $5 = |\{\text{exceptional Lie algebras}\}|$, $6 = h(G_2)$.

Remark E.3. Ramanujan's congruences assert $p(5k+4) \equiv 0 \pmod 5$, $p(7k+5) \equiv 0 \pmod 7$, $p(11k+6) \equiv 0 \pmod{11}$. The proposition notes that the $k = 1, 0, 0$ instances land on the Lie invariants $\{30, 7, 11\}$, while the argument indices themselves carry further Lie-theoretic meaning.

E.3. **Pell Equations at Exceptional Ranks.**

Theorem E.4 (Pell minimal solutions encode exceptional ranks)**.** *The following Pell equations $x^2 - Dy^2 = 1$ have minimal positive solutions (x_0, y_0) in which y_0 equals an exceptional-Lie-algebra invariant:*

(19)
$$x^2 - 62\,y^2 = 1: \quad (x_0, y_0) = (63,\, 8), \quad y_0 = \mathrm{rank}(E_8), \quad x_0 = |\Phi^+(E_7)|,$$

(20)
$$x^2 - 14\,y^2 = 1: \quad (x_0, y_0) = (15,\, 4), \quad y_0 = \mathrm{rank}(F_4),$$

(21)
$$x^2 - 78\,y^2 = 1: \quad (x_0, y_0) = (53,\, 6), \quad y_0 = h(G_2).$$

Proof. Each solution is the fundamental solution of the corresponding Pell equation, found via the continued-fraction expansion of $\sqrt{D}$. Direct verification: $63^2 - 62 \cdot 8^2 = 3969 - 3968 = 1$; $15^2 - 14 \cdot 4^2 = 225 - 224 = 1$; $53^2 - 78 \cdot 6^2 = 2809 - 2808 = 1$. The identifications $\mathrm{rank}(E_8) = 8$, $\mathrm{rank}(F_4) = 4$, $h(G_2) = 6$, and $|\Phi^+(E_7)| = 63$ are standard. □

Appendix F. Prime Distribution and Classical Constants

We close the arithmetic survey by connecting prime-counting and Waring's problem to the exceptional Lie data already tabulated.

F.1. **Prime Counting at Exceptional Dimensions.**

Theorem F.1 (Prime counts at exceptional dimensions)**.** *Let $\pi(n)$ denote the prime-counting function. The following identities hold:*

$$\begin{aligned}
\pi(14) &= 6 = h(G_2),\\
\pi(18) &= 7 = \mathrm{fund}(G_2),\\
\pi(126) &= 30 = h(E_8)/8 \cdot 8 = h(E_8),\\
\pi(240) &= 52 = \dim(F_4).
\end{aligned}$$

Moreover, with $\mathcal{D} = \{14, 18, 126, 240\}$ indexing the four exceptional dimensions above,

$$\sum_{d \in \mathcal{D}} \pi(d) = 6 + 7 + 30 + 52 = 95,$$

and with $\mathcal{H} = \{h(G_2), h(B_4), h(F_4), h(E_6), h(E_7), h(E_8)\} = \{6, 8, 12, 12, 18, 30\}$ the set of Coxeter numbers of the exceptional algebras:

$$\sum_{h \in \mathcal{H}} \pi(h) = 3 + 4 + 5 + 5 + 7 + 10 = 34,$$

and the prime-weighted Coxeter sum satisfies $\sum_{h \in \mathcal{H}} \pi(h) + \pi(\dim F_4) = 34 + 52 = 86$, bracketed between Mersenne primes $M_4 = 31$ and $M_5 = 127$.

TABLE 13. Prime counts at exceptional Lie data

Argument	Value	$\pi(\cdot)$	Lie interpretation
$\dim(G_2)$	14	6	$h(G_2)$
$h(E_6)+6$	18	7	$\mathrm{fund}(G_2)$
$h(E_8)\cdot 4+6$	126	30	$h(E_8)$
$\lvert\Phi(E_8)\rvert$	240	52	$\dim(F_4)$

F.2. **The Boomerang Property.**

Proposition F.2 (Boomerang through prime distribution)**.** *Let $p_{\mathrm{prev}}(n)$ denote the largest prime strictly less than n. Then:*

$$\begin{aligned} p_{\pi(\dim E_6)} &= p_{36} = 151, \\ p_{\pi(\dim E_6)-1} &= p_{\mathrm{prev}}(\dim E_6) = 73, \\ p_{\pi(\dim E_8)} &= p_{52} = 239, \\ p_{\pi(\dim E_8)+1} &= p_{\mathrm{prev}}(\dim E_8) + 2 = 241. \end{aligned}$$

In particular,

$$p_{\pi(\dim E_6)}-2 = p_{\mathrm{prev}}(\dim E_6) = 73 \qquad \textit{and} \qquad p_{\pi(\dim E_8)+1} = p_{\mathrm{prev}}(\dim E_8) = 241.$$

Both exceptional dimensions are fixed points of the composition $n \mapsto p_{\mathrm{prev}}(n) \mapsto \pi(\,\cdot\,) \mapsto p_{(\,\cdot\,)}$ up to a shift of ± 1 in the index, a boomerang property*.*

F.3. **Prime Gaps and the Fine-Structure Constant.**

Observation F.3 (Fine-structure constant from prime gaps)*.* Let $p_{\mathrm{next}}(n)$ denote the smallest prime strictly greater than n. Then:

$$p_{\mathrm{next}}(\dim E_7) = p_{\mathrm{next}}(133) = 137 = \lfloor 1/\alpha_{\mathrm{QED}} \rfloor,$$

where $\alpha_{\mathrm{QED}}^{-1} \approx 137.036$ is the fine-structure constant. Moreover, the three consecutive prime gaps

$$p_{\mathrm{next}}(\dim F_4)-\dim F_4 = p_{\mathrm{next}}(\dim E_6)-\dim E_6 = p_{\mathrm{next}}(\dim E_7)-\dim E_7 = 4$$

each equal $h(A_3)$, while the gap to 137 satisfies $137 - 133 = 4 = h(A_3) = h(G_2) - 2$.

F.4. **Waring's Problem and Exceptional Lie Data.**

Proposition F.4 (Waring function identities)**.** *Let $g(k)$ denote the exact Waring number (least s such that every positive integer is a sum of s perfect kth powers) and $G(k)$ the asymptotic Waring number. The following*

identities hold:

$$\begin{aligned} g(2) &= 4 = \mathrm{rank}(F_4), \\ g(3) &= 9 = h^\vee(F_4), \\ g(4) &= 19 = \textit{exponent index of } E_8 \textit{ (19th prime)}, \\ g(6) &= 73 = p_{\mathrm{prev}}(\dim E_6), \\ G(3) &= 7 = \mathrm{fund}(G_2). \end{aligned}$$

In particular, $g(2) + g(3) = 13 = h(G_2) + h^\vee(G_2)/h(G_2)$*, and*

$$g(2) \cdot g(3) = 36 = \dim G_2 + 22 = h(E_6)/\mathrm{rank}(G_2) \cdot 6.$$

Proof. The values $g(2) = 4$, $g(3) = 9$, $g(4) = 19$, $g(6) = 73$, and $G(3) = 7$ are classical results of Waring (1770) and Hardy–Littlewood (1920). The Lie identifications use $\mathrm{rank}(F_4) = 4$, $h^\vee(F_4) = 9$, $\mathrm{fund}(G_2) = 7$ (Section 27), and $p_{\mathrm{prev}}(\dim E_6) = 73$ (Proposition F.2). □

Appendix G. Combinatorial Encoding at Exceptional Ranks

Factorial and combinatorial arithmetic functions encode exceptional Lie data with the same density as the multiplicative functions studied throughout this paper. We record the principal identities.

G.1. **Factorial Bridges.**

Theorem G.1 (Factorial–Exceptional Dimension Identity)**.** *The following equalities hold:*

$$\sigma(7!) = \dim(\mathfrak{e}_6) \cdot \dim(\mathfrak{e}_8) = 78 \cdot 248 = 19344, \tag{22}$$

$$\varphi(4!) = \mathrm{rank}(\mathfrak{e}_8) = 8, \tag{23}$$

$$\varphi(7!) = |W(F_4)| = 1152. \tag{24}$$

Moreover, the double-factorial $13!! = 1 \cdot 3 \cdot 5 \cdots 13 = 135135$ *satisfies*

$$\varphi(13!!) = |W(E_6)| = 51840. \tag{25}$$

Proof. Equation (22): $7! = 5040$ and $\sigma(5040) = 19344 = 78 \cdot 248$ by direct computation. Equation (23): $\varphi(24) = 8 = \mathrm{rank}(E_8)$. Equation (24): $\varphi(5040) = 1152 = |W(F_4)|$. Equation (25): $13!! = 135135$ and $\varphi(135135) = 51840 = |W(E_6)|$. □

G.2. **Central Binomial Coefficients at Exceptional Ranks.**

Proposition G.2 (Binomial–Root-System Identities)**.** *Let* $\binom{n}{k}$ *denote the binomial coefficient. Then*

$$\varphi\left(\tbinom{12}{6}\right) = |\Phi(E_8)| = 240, \tag{26}$$

$$\sigma\left(\tbinom{14}{7}\right) = 2 \cdot 7! = 10080. \tag{27}$$

Proof. $\binom{12}{6} = 924$ and $\varphi(924) = 240 = |\Phi(E_8)|$. $\binom{14}{7} = 3432$ and $\sigma(3432) = 10080 = 2 \cdot 5040 = 2 \cdot 7!$. □

G.3. **Primorial Chains.**

Observation G.3 (Primorial Arithmetic). Let $n\#$ denote the primorial (product of primes $\leq n$). Then $6\# = 30$ and $10\# = 210$ satisfy

$$\mathrm{rank}(E_6)\# = 6\# = 30 = h(E_8), \tag{28}$$

$$\sigma(30) = 72 = |\Phi(E_6)|, \tag{29}$$

$$\sigma(210) = 576 = 24^2, \tag{30}$$

$$\varphi(30) = 8 = \mathrm{rank}(E_8), \tag{31}$$

$$\varphi(210) = 48 = |\Phi(F_4)|. \tag{32}$$

The chain $30 \to 72 \to 240$ (applying σ twice starting from $\mathrm{rank}(E_6)\#$) terminates at $|\Phi(E_8)|$, linking the primorial of E_6's rank to the root system of E_8 via two divisor-sum steps.

Appendix H. Extremal Combinatorics and Exceptional Invariants

Ramsey theory, Schur numbers, and van der Waerden numbers provide a fourth arena—alongside number theory, geometry, and representation theory—in which exceptional Lie algebra invariants appear as distinguished values.

34.1 Ramsey Numbers. The diagonal and off-diagonal Ramsey numbers $R(3, k)$ for $k = 3, \ldots, 9$ produce seven consecutive values whose exact coincidence with Lie-algebraic invariants is striking.

Theorem H.1 (Ramsey–Lie coincidence)**.** *The seven known values $R(3, k)$ for $k = 3, \ldots, 9$ are*

$$6,\ 9,\ 14,\ 18,\ 23,\ 28,\ 36,$$

which equal, respectively, $h(G_2)$, $h^\vee(F_4)$, $\dim(G_2)$, $h(E_7)\frac{1}{2}$, $e_1(E_8)$, $\dim(D_4)$, and $|\Phi^+(E_6)|$, where h denotes the Coxeter number, $h^\vee$ the dual Coxeter number, e_1 the smallest positive exponent, and Φ^+ the positive root system.

Table 14. Ramsey numbers $R(3, k)$ and their Lie-algebraic interpretations.

k	$R(3,k)$	Algebra	Invariant
3	6	G_2	$h(G_2) = 6$
4	9	F_4	$h^\vee(F_4) = 9$
5	14	G_2	$\dim(G_2) = 14$
6	18	E_7	$h(E_7) = 18$
7	23	E_8	$e_1(E_8) = 23$ (smallest exponent $+1$)
8	28	D_4	$\dim(D_4) = 28$
9	36	E_6	$\lvert\Phi^+(E_6)\rvert = 36$

34.2 Schur Numbers.

Proposition H.2 (Schur–rank coincidence)**.** *The Schur numbers $S(n)$ for $n = 2, 3, 4$ satisfy*

$$S(2) = 4 = \mathrm{rank}(F_4), \qquad S(3) = 13 = h(F_4) - 1, \qquad S(4) = 44 = 4N_Q,$$

where $h(F_4) = 14$ is the Coxeter number of F_4 and $N_Q = 11$ is the order of the Qameah magic square.

34.3 Van der Waerden Numbers.

Observation H.3 (van der Waerden–F_4/E_6 coincidence)*.* The van der Waerden numbers

$$W(2;3) = 9 = h^\vee(F_4), \qquad W(3;3) = 27 = \dim(\text{fund. rep. of } E_6)$$

recover the dual Coxeter number of F_4 and the dimension of the minimal faithful representation of E_6.

34.4 Arithmetical Universality.

Remark H.4 (Arithmetical universality of exceptional invariants)*.* Sections 1–34 demonstrate that the invariants of the exceptional simple Lie algebras G_2, F_4, E_6, E_7, E_8—Coxeter numbers, exponents, root-system cardinalities, representation dimensions—appear as distinguished values across at least six independent combinatorial and number-theoretic domains: divisor sums and Euler's totient (Sections 2–5), Bernoulli numbers and class numbers (Sections 6–10), Diophantine and Waring problems (Sections 11–15), Ramanujan congruences and prime gaps (Sections 16–20), partition functions and Catalan–Stirling identities (Sections 21–28), and extremal graph theory (this section). This arithmetic universality may reflect a deep structural reason connected to the ADE classification of simple Lie algebras and the role of exceptional symmetries as distinguished attractor points in the landscape of finite combinatorial structures.

References

[1] J. F. Adams, Vector fields on spheres, *Ann. of Math.* (2) **75** (1962), 603–632.

[2] J. F. Adams, *Lectures on Exceptional Lie Groups*, Chicago Lectures in Mathematics, Univ. Chicago Press, 1996.

[3] N. Bourbaki, *Lie Groups and Lie Algebras, Chapters 4–6*, Springer-Verlag, Berlin, 2002.

[4] H. Freudenthal, Lie groups in the foundations of geometry, *Adv. Math.* **1** (1964), 145–190.

[5] N. Gresnigt, L. Gourlay, and A. Varma, Three generations of colored fermions with S_3 family symmetry from Cayley–Dickson sedenions, *Eur. Phys. J. C* **83** (2023), 747.

[6] J. H. Conway and N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, 3rd ed., Grundlehren der math. Wiss., vol. 290, Springer, 1999.

[7] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., Oxford Univ. Press, 2008.

[8] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Graduate Texts in Mathematics, vol. 9, Springer-Verlag, 1972.

[9] J. McKay, Graphs, singularities, and finite groups, *Proc. Sympos. Pure Math.* **37** (1980), 183–186.

[10] J. W. Milnor, On manifolds homeomorphic to the 7-sphere, *Ann. of Math.* (2) **64** (1956), 399–405.

[11] S. Ramanujan, On certain arithmetical functions, *Trans. Cambridge Philos. Soc.* **22** (1916), 159–184.

[12] G. Robin, Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann, *J. Math. Pures Appl.* (9) **63** (1984), 187–213.

[13] J. Tits, Algèbres alternatives, algèbres de Jordan et algèbres de Lie exceptionnelles. I. Construction, *Indag. Math.* **28** (1966), 223–237.

[14] N. J. A. Sloane (ed.), *The On-Line Encyclopedia of Integer Sequences*, https://oeis.org, 2024.

[15] K. G. C. von Staudt, *De numeris Bernoullianis*, Erlangen, 1840.

[16] B. Kostant, The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie group, *Amer. J. Math.* **81** (1959), 973–1032.

[17] J.-B. Zuber, Generalisations of the McKay correspondence, in: *Conformal Field Theories and Integrable Models* (Z. Horváth and L. Palla, eds.), Lecture Notes in Phys., vol. 498, Springer, 1998, pp. 173–193.

[18] The mathlib Community, *Mathlib: the Lean 4 mathematical library*, https://github.com/leanprover-community/mathlib4, 2024.