

# Construction of $\mathrm{Ver}_{p^n}$

Summary of some work from Benson, Etingof, and Ostrik

Andrew Aguilar and Charlie Magland

University of Washington

November 20, 2025

# Overview

---

1. Introduction
2. Tilting Modules
3. Construction of  $\text{Ver}_{p^n}$
4. Tensor Product
5. Properties
6. Spec

# Introduction

---

A symmetric tensor category  $\mathcal{C}$  over an algebraically closed field with objects of finite length is *incompressible* if every tensor functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is an embedding of a tensor subcategory.

By a theorem of Deligne, in characteristic 0, the only such categories (of moderate growth) are **Vec** and **sVec**.

# Introduction

A symmetric tensor category  $\mathcal{C}$  over an algebraically closed field with objects of finite length is *incompressible* if every tensor functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is an embedding of a tensor subcategory.

By a theorem of Deligne, in characteristic 0, the only such categories (of moderate growth) are **Vec** and **sVec**.

## Theorem (Benson, Etingof, and Ostrik)

*Let  $k$  be an algebraically closed field of characteristic  $p$ . Then there are nested sequences of incompressible symmetric tensor categories*

$$\mathrm{Ver}_p \subset \mathrm{Ver}_{p^2} \subset \mathrm{Ver}_{p^3} \subset \dots$$

*and*

$$\mathrm{Ver}_p^+ \subset \mathrm{Ver}_{p^2}^+ \subset \mathrm{Ver}_{p^3}^+ \subset \dots$$

# Introduction

---

Recall:  $\text{Ver}_p$  is the semisimplification of  $\text{Rep}(\mathbb{Z}/p\mathbb{Z})$ .

$\text{Ver}_{p^n}$  is the abelian envelope of  $\mathcal{T}_{n,p}$ , a quotient of the category of tilting modules of  $SL_2$ .

Abelian envelopes, unlike Karoubian envelopes, are difficult to construct. [BEO21] proposes a construction that, under the right conditions, gives the abelian envelope explicitly.

# Tilting Modules

## Definition

A  $G$ -module  $M$  is called *Tilting* if  $M$  and  $M^*$  both have a good filtration. A  $G$ -module  $M$  has a *good filtration* if there is an ascending chain of submodules

$$0 = M_0 \subset M_1 \subset M_2 \subset \dots$$

1.  $M = \bigcup_{i \geq 0} M_i$
2. Each  $M_i/M_{i-1}$  is isomorphic to  $\mathrm{ind}_B^G(k_\lambda)$

Where

- $B = U \rtimes T$  is the Borel subgroup of  $G$
- $k_\lambda$  is  $k$  with the action given by  $\lambda : G \mapsto GL_1 = G_m$

## Tilting Modules for $SL_2$

Let  $V$  be the standard representation for  $SL_2$ .

- Each  $V^{\otimes m}$  has an indecomposable summand,  $T_m$ , not in previous tensor powers.
- The indecomposable tilting modules of highest weight  $m$  are exactly these  $T_m$ ,  $m \geq 0$
- The category of tilting modules, denoted  $\mathcal{T}_p$ , is spanned by the  $T_m$ .
- $\mathcal{T}_p$  is closed under tensoring, so tilting modules form a monoidal category.

## Tilting Modules for $SL_2$

Let  $V$  be the standard representation for  $SL_2$ .

- Each  $V^{\otimes m}$  has an indecomposable summand,  $T_m$ , not in previous tensor powers.
- The indecomposable tilting modules of highest weight  $m$  are exactly these  $T_m$ ,  $m \geq 0$
- The category of tilting modules, denoted  $\mathcal{T}_p$ , is spanned by the  $T_m$ .
- $\mathcal{T}_p$  is closed under tensoring, so tilting modules form a monoidal category.

Special cases:

- The first  $p - 1$  indecomposable tilting modules are all simple.
- The *Steinberg modules*  $T_{p^n-1}$  are simple for every  $n \geq 1$ .



# Tilting Modules

---

- Within  $\mathcal{T}_p$ , we can construct a tensor ideal

$$\mathcal{I}_n = \langle T_m \mid m \geq p^n - 1 \rangle \quad \forall n \geq 1.$$

This is generated by the  $n$ th Steinberg module  $T_{p^n-1}$ .

# Tilting Modules

---

- Within  $\mathcal{T}_p$ , we can construct a tensor ideal

$$\mathcal{I}_n = \langle T_m \mid m \geq p^n - 1 \rangle \quad \forall n \geq 1.$$

This is generated by the  $n$ th Steinberg module  $T_{p^n-1}$ .

- Define the quotient category  $\mathcal{T}_{n,p} = \mathcal{T}_p / \mathcal{I}_n$ , with indecomposables  $T_i$  for  $0 \leq i \leq p^n - 2$ .

# Tilting Modules

---

- Within  $\mathcal{T}_p$ , we can construct a tensor ideal

$$\mathcal{I}_n = \langle T_m \mid m \geq p^n - 1 \rangle \quad \forall n \geq 1.$$

This is generated by the  $n$ th Steinberg module  $T_{p^n-1}$ .

- Define the quotient category  $\mathcal{T}_{n,p} = \mathcal{T}_p / \mathcal{I}_n$ , with indecomposables  $T_i$  for  $0 \leq i \leq p^n - 2$ .
- We are interested in the *new* indecomposables in  $\mathcal{T}_{n,p}$  not contained in  $\mathcal{T}_{n-1,p}$ , highlighted in red below.

$$k \quad T_1 \quad \dots \quad T_{p^{n-1}-2} \quad \textcolor{red}{T_{p^{n-1}-1}} \quad \dots \quad \textcolor{red}{T_{p^n-2}} \quad \textcolor{blue}{T_{p^n-1}} \quad \dots$$

- These generate a tensor ideal in  $\mathcal{T}_{n,p}$ , denoted by  $\mathcal{S}_{n,p} = \mathcal{I}_{n-1} / \mathcal{I}_n$ .

# Construction of $\text{Ver}_{p^n}$

## Definition

Construction of  $\text{Ver}_{p^n}$  Recall  $\mathcal{T}_{n,p}$  with ideal  $\mathcal{S}_{n,p} = \langle T_i \mid p^{n-1} - 1 \leq i < p^n - 1 \rangle$ . Take

$$P = \bigoplus_{i=p^{n-1}-1}^{p^n-2} T_i$$

and let  $A = \text{End}(P)$ .

# Construction of $\text{Ver}_{p^n}$

## Definition

Construction of  $\text{Ver}_{p^n}$  Recall  $\mathcal{T}_{n,p}$  with ideal  $\mathcal{S}_{n,p} = \langle T_i \mid p^{n-1} - 1 \leq i < p^n - 1 \rangle$ . Take

$$P = \bigoplus_{i=p^{n-1}-1}^{p^n-2} T_i$$

and let  $A = \text{End}(P)$ . We define

$$\text{Ver}_{p^n} := \{\text{Finite dimensional } A\text{-modules}\}.$$

# Construction of $\text{Ver}_{p^n}$

## Definition

Construction of  $\text{Ver}_{p^n}$  Recall  $\mathcal{T}_{n,p}$  with ideal  $\mathcal{S}_{n,p} = \langle T_i \mid p^{n-1} - 1 \leq i < p^n - 1 \rangle$ . Take

$$P = \bigoplus_{i=p^{n-1}-1}^{p^n-2} T_i$$

and let  $A = \text{End}(P)$ . We define

$$\text{Ver}_{p^n} := \{\text{Finite dimensional } A\text{-modules}\}.$$

Similarly, let  $P^+ = \bigoplus_{\substack{i=p^{n-1}-1 \\ i \text{ even}}}^{p^n-2} T_i$  and define  $\text{Ver}_{p^n}^+ := \{\text{Finite dimensional } A\text{-modules}\}$

# Construction of $\text{Ver}_{p^n}$

## Projective Objects in $\text{Ver}_{p^n}$

Since  $\text{End}(P) = \bigoplus_{i,j} \text{Hom}(T_i, T_j)$ , we get a map  $\iota : \mathcal{S}_{n,p} \rightarrow \text{Ver}_{p^n}$  by

$$\iota(T_i) = \bigoplus_{j=p^{n-1}-1}^{p^n-2} \text{Hom}(T_j, T_i) = \text{Hom}(P, T_i)$$

which is a summand of  $\text{End}(P)$ .

# Construction of $\text{Ver}_{p^n}$

## Projective Objects in $\text{Ver}_{p^n}$

Since  $\text{End}(P) = \bigoplus_{i,j} \text{Hom}(T_i, T_j)$ , we get a map  $\iota : \mathcal{S}_{n,p} \rightarrow \text{Ver}_{p^n}$  by

$$\iota(T_i) = \bigoplus_{j=p^{n-1}-1}^{p^n-2} \text{Hom}(T_j, T_i) = \text{Hom}(P, T_i)$$

which is a summand of  $\text{End}(P)$ . Thus, the image of each  $T_i$  is projective, and in fact

$$\iota(\mathcal{S}_{n,p}) = \text{Proj}(\text{Ver}_{p^n}).$$



# Construction of $\text{Ver}_{p^n}$

## Projective Objects in $\text{Ver}_{p^n}$

Since  $\text{End}(P) = \bigoplus_{i,j} \text{Hom}(T_i, T_j)$ , we get a map  $\iota : \mathcal{S}_{n,p} \rightarrow \text{Ver}_{p^n}$  by

$$\iota(T_i) = \bigoplus_{j=p^{n-1}-1}^{p^n-2} \text{Hom}(T_j, T_i) = \text{Hom}(P, T_i)$$

which is a summand of  $\text{End}(P)$ . Thus, the image of each  $T_i$  is projective, and in fact

$$\iota(\mathcal{S}_{n,p}) = \text{Proj}(\text{Ver}_{p^n}).$$

Additionally, the indecomposable objects  $\{T_i\}_{p^{n-1}-1 \leq i \leq p^n-2}$  in  $\mathcal{S}_{n,p}$  correspond to the indecomposable projectives in  $\text{Ver}_{p^n}$ .

# Construction of $\text{Ver}_{p^n}$

---

## Example: $\text{Ver}_p$

Let  $n = 1$ .

$\mathcal{S}_{1,p}$  has indecomposables  $k, T_1, \dots, T_{p-2}$ . These are all simple, and are in correspondence with the simple objects of  $\text{Ver}_p$ .

We will construct a tensor product which will agree with the monoidal structure of  $\text{Ver}_p$ .

# Construction of $\text{Ver}_{p^n}$

## Example: $\text{Ver}_4^+$ and $\text{Ver}_4$

Let  $p = 2$ ,  $n = 2$  and  $V = k\{x, y\}$  be the standard module for  $SL_2$ .

- The indecomposables in  $\mathcal{S}_{2,2}$  are  $T_1 = V$  and  $T_2 = V \otimes V$ .
- The tensor product  $T_2$  is uniserial with composition factors  $[k, V^{(1)}, k]$ .
- We have  $\text{Ver}_4^+ = \text{End}(V \otimes V) - \text{mod} \cong k[d]/d^2\text{-mod}$ .
  - The projective module is  $A$  as a right module.
- We have  $\text{Ver}_4 = \text{End}(T_1 \oplus T_2) - \text{mod}$ .
  - $\text{End}(V)$  is a projective simple, while  $\text{End}(V \otimes V)$  is an indecomposable projective the same as in  $\text{Ver}_4^+$ .

# Tensor Product

---

Equivalently,  $\text{Ver}_{p^n}$  can be realized as complexes of projectives in

$$K^-(\text{Pr}(\text{Ver}_{p^n})) \cong K^-(\mathcal{S}_{n,p}).$$

# Tensor Product

---

Equivalently,  $\text{Ver}_{p^n}$  can be realized as complexes of projectives in

$$K^-(\text{Pr}(\text{Ver}_{p^n})) \cong K^-(\mathcal{S}_{n,p}).$$

For two objects  $X, Y \in \text{Ver}_{p^n}$  with resolutions  $P_\bullet \rightarrow X$  and  $Q_\bullet \rightarrow Y$  we can define  $X \otimes Y = H^0(\text{Tot}(P_\bullet \otimes Q_\bullet))$ , or equivalently

$$\text{coker}(P_1 \otimes Q_0 \oplus P_0 \otimes Q_1 \xrightarrow{\varphi} P_0 \otimes Q_0)$$

# Tensor Product

---

Equivalently,  $\text{Ver}_{p^n}$  can be realized as complexes of projectives in

$$K^-(\text{Pr}(\text{Ver}_{p^n})) \cong K^-(\mathcal{S}_{n,p}).$$

For two objects  $X, Y \in \text{Ver}_{p^n}$  with resolutions  $P_\bullet \rightarrow X$  and  $Q_\bullet \rightarrow Y$  we can define  $X \otimes Y = H^0(\text{Tot}(P_\bullet \otimes Q_\bullet))$ , or equivalently

$$\text{coker}(P_1 \otimes Q_0 \oplus P_0 \otimes Q_1 \xrightarrow{\varphi} P_0 \otimes Q_0)$$

The tensor product agrees with the tensor on  $\mathcal{S}_{n,p}$ , and is

- symmetric since  $\mathcal{T}_{n,p}$  is
- right exact in both arguments
- exact if either argument is projective (ie. belongs to  $\mathcal{S}_{n,p}$ )

# Monoidal Unit

Now, we describe the monoidal unit of  $\text{Ver}_{p^n}$ . Since  $P$  is projective we have

$$P \otimes P^* \otimes P \otimes P^* \xrightarrow{\tau} P \otimes P^*$$

where  $\tau := \text{ev} \otimes 1 \otimes 1 - 1 \otimes 1 \otimes \text{ev}$ . Then define  $\text{coker}(\tau) = \mathbb{1}_{\text{Ver}_{p^n}}$ .

## Proposition

Taking  $Q \in \mathcal{S}_{n,p}$  gives isomorphisms

- $\mathbb{1} \otimes Q \cong Q \otimes \mathbb{1} \cong Q$  which are functorial in  $Q$ .
- and induces isomorphisms  $\mathbb{1} \otimes - \cong - \otimes \mathbb{1} \cong \text{Id}_C$ .

So  $\mathbb{1}$  is the monoidal unit in  $\text{Ver}_{p^n}$

# The Functor $F : \mathcal{T}_{n,p} \rightarrow \text{Ver}_{p^n}$

The tensor product gives an action  $\mathcal{T}_{n,p} \curvearrowright \mathcal{S}_{n,p}$  which we can extend to  $\text{Ver}_{p^n}$  via projective resolutions.

Because  $\mathcal{S}_{n,p}$  is a tensor ideal we can tensor the projective resolution of  $\mathbb{1}$  with elements of  $\mathcal{T}_{n,p}$  to get their image in  $\text{Ver}_{p^n}$ .

$$\text{coker}(P \otimes P^* \otimes P \otimes P^* \otimes X \xrightarrow{\tau \otimes X} P \otimes P^* \otimes X) = T_r(X)$$

## Corollary

*There is a monoidal functor  $F : \mathcal{T}_{n,p} \rightarrow \text{Ver}_{p^n}$  defined by  $F(T) = T_r(\mathbb{1})$ . Since  $\mathcal{T}_{n,p}$  is symmetric, so is  $F$ .*

Denote  $\mathbb{T}_i = F(T_i)$  for each indecomposable  $T_i \in \mathcal{T}_{n,p}$



# Fusion Rules

---

Because the tensor is inherited from  $\mathcal{T}_{n,p}$ , we get similar fusion rules as for  $SL_2$ -modules. For  $1 \leq m, r \leq p-1$  we have

$$\mathbb{T}_m \otimes \mathbb{T}_r = \begin{cases} \bigoplus_{\substack{|m-r| \leq k \leq m+r \\ m+r-k \text{ even}}} \mathbb{T}_k & m+r < p \\ \bigoplus_{\substack{|m-r| \leq k \leq 2(p-2)-m-r \\ m+r-k \text{ even}}} \mathbb{T}_k \oplus \bigoplus_{\substack{p-1 \leq k \leq m+r \\ m+r-k \text{ even}}} \mathbb{T}_k & m+r \geq p \end{cases}$$

# $\text{Ver}_{p^n}$ is a symmetric tensor category

---

## Definition

A *Symmetric Tensor Category* is an artinian  $k$ -linear abelian rigid monoidal category with biexact tensor product and  $\text{End}(\mathbb{1}) \cong k$ . It is *finite* if it is equivalent to  $A\text{-mod}$  with  $A$  finite dimensional.

# $\text{Ver}_{p^n}$ is a symmetric tensor category

---

## Definition

A *Symmetric Tensor Category* is an artinian  $k$ -linear abelian rigid monoidal category with biexact tensor product and  $\text{End}(\mathbb{1}) \cong k$ . It is *finite* if it is equivalent to  $A\text{-mod}$  with  $A$  finite dimensional.

Need to show:

- Rigidity
- Biexactness of tensor
- $\text{End}(\mathbb{1}) \cong k$

# $\text{Ver}_{p^n}$ is a symmetric tensor category

---

## Definition

A *Symmetric Tensor Category* is an artinian  $k$ -linear abelian rigid monoidal category with biexact tensor product and  $\text{End}(\mathbb{1}) \cong k$ . It is *finite* if it is equivalent to  $A\text{-mod}$  with  $A$  finite dimensional.

Need to show:

- Rigidity
- Biexactness of tensor
- $\text{End}(\mathbb{1}) \cong k$

## Corollary

*Every projective object in  $\text{Ver}_{p^n}$  is rigid*

# Splitting Ideals

---

## Definition

An ideal  $\mathcal{P} \subseteq \mathcal{T}$  is a *splitting ideal* if for any  $Q_1, Q_2, R \in \mathcal{P}$  and  $f : Q_1 \rightarrow Q_2$ ,  $1_R \otimes f : R \otimes Q_1 \rightarrow R \otimes Q_2$  is split. The objects in  $\mathcal{P}$  are called *splitting objects*.

# Splitting Ideals

## Definition

An ideal  $\mathcal{P} \subseteq \mathcal{T}$  is a *splitting ideal* if for any  $Q_1, Q_2, R \in \mathcal{P}$  and  $f : Q_1 \rightarrow Q_2$ ,  $1_R \otimes f : R \otimes Q_1 \rightarrow R \otimes Q_2$  is split. The objects in  $\mathcal{P}$  are called *splitting objects*.

## Lemma

The Steinberg module  $St_n$  is a splitting object in  $\mathcal{S}_{n,p}$ .

# Splitting Ideals

## Definition

An ideal  $\mathcal{P} \subseteq \mathcal{T}$  is a *splitting ideal* if for any  $Q_1, Q_2, R \in \mathcal{P}$  and  $f : Q_1 \rightarrow Q_2$ ,  $1_R \otimes f : R \otimes Q_1 \rightarrow R \otimes Q_2$  is split. The objects in  $\mathcal{P}$  are called *splitting objects*.

## Lemma

*The Steinberg module  $St_n$  is a splitting object in  $\mathcal{S}_{n,p}$ .*

Taking a resolution of  $Y \in \text{Ver}_{p^n}$  and  $Q \in \mathcal{S}_{n,p}$  the map

$$Q \otimes Q_1 \rightarrow Q \otimes Q_0 \rightarrow Q \otimes Y \rightarrow 0$$

splits, so  $Q \otimes Y$  is a summand.

# Splitting Ideals

## Definition

An ideal  $\mathcal{P} \subseteq \mathcal{T}$  is a *splitting ideal* if for any  $Q_1, Q_2, R \in \mathcal{P}$  and  $f : Q_1 \rightarrow Q_2$ ,  $1_R \otimes f : R \otimes Q_1 \rightarrow R \otimes Q_2$  is split. The objects in  $\mathcal{P}$  are called *splitting objects*.

## Lemma

*The Steinberg module  $St_n$  is a splitting object in  $\mathcal{S}_{n,p}$ .*

Taking a resolution of  $Y \in \text{Ver}_{p^n}$  and  $Q \in \mathcal{S}_{n,p}$  the map

$$Q \otimes Q_1 \rightarrow Q \otimes Q_0 \rightarrow Q \otimes Y \rightarrow 0$$

splits, so  $Q \otimes Y$  is a summand.

## Corollary

*For any  $Y \in \text{Ver}_{p^n}$  and  $Q \in \mathcal{S}_{n,p}$  the tensor products  $Y \otimes Q$  and  $Q \otimes Y$  are projective.*



# Tensor Product II

---

We have the left derived functors  $X \otimes^{L_i} Y = H^{-i}(\text{Tot}(P_{\bullet} \otimes Q_{\bullet})) = H^{-i}(X \otimes Q_{\bullet})$ . Then by previous corollary we get

# Tensor Product II

---

We have the left derived functors  $X \otimes^{L_i} Y = H^{-i}(\text{Tot}(P_{\bullet} \otimes Q_{\bullet})) = H^{-i}(X \otimes Q_{\bullet})$ . Then by previous corollary we get

## Corollary

*The bifunctor  $(X, Y) \rightarrow X \otimes Y$  is exact in both arguments*

# Tensor Product II

We have the left derived functors  $X \otimes^{L_i} Y = H^{-i}(\text{Tot}(P_{\bullet} \otimes Q_{\bullet})) = H^{-i}(X \otimes Q_{\bullet})$ . Then by previous corollary we get

## Corollary

*The bifunctor  $(X, Y) \rightarrow X \otimes Y$  is exact in both arguments*

If  $P_1 \rightarrow P_2 \rightarrow P_3$  is an acyclic complex of projectives then we can show that

$$X \otimes P_1 \rightarrow X \otimes P_2 \rightarrow X \otimes P_3$$

is acyclic when

$$\text{Hom}(R, X \otimes P_1) \rightarrow \text{Hom}(R, X \otimes P_2) \rightarrow \text{Hom}(R, X \otimes P_3)$$

is acyclic for every projective  $R$ .

# Rigidity

---

## Proposition

Let  $X \in \text{Ver}_{p^n}$  which is the cokernel of  $f : Q_1 \rightarrow Q_0$ . Then  $X$  has a dual  $X^*$  which is the kernel of the map  $f^* : Q_0^* \rightarrow Q_1^*$ .

# Rigidity

## Proposition

Let  $X \in \text{Ver}_{p^n}$  which is the cokernel of  $f : Q_1 \rightarrow Q_0$ . Then  $X$  has a dual  $X^*$  which is the kernel of the map  $f^* : Q_0^* \rightarrow Q_1^*$ .

Let  $\iota : X^* \rightarrow Q_0^*$  and  $p : Q_1^* \rightarrow X$ , then the evaluation and coevaluation maps are constructed as

$$\text{ev}_{Q_0} \circ (\iota \circ \text{id}_{Q_0}) : X^* \otimes Q_0 \rightarrow \mathbb{1} \implies \text{ev}_X : X^* \otimes X \rightarrow \mathbb{1}$$

and

$$(p \otimes \text{id}_{Q_0^*}) \circ \text{coev}_{Q_0} : \mathbb{1} \rightarrow X \otimes Q_1^* \implies \text{coev}_X : \mathbb{1} \rightarrow X \otimes X^*$$

$$\mathrm{End}(\mathbb{1}) \cong k$$

---

Let  $\mathcal{S}$  be the set of splitting objects in  $\mathcal{T}$ .

- Define an equivalence relation on  $\mathcal{S}$  by  $Q \sim_I P$  if there exists  $X \in \mathcal{T}$  such that  $Q$  is a direct summand of  $X \otimes R$ .

$$\mathrm{End}(\mathbb{1}) \cong k$$

---

Let  $\mathcal{S}$  be the set of splitting objects in  $\mathcal{T}$ .

- Define an equivalence relation on  $\mathcal{S}$  by  $Q \sim_I P$  if there exists  $X \in \mathcal{T}$  such that  $Q$  is a direct summand of  $X \otimes R$ .
- Then we can take the equivalence classes  $\overline{\mathcal{S}} = \mathcal{S} / \sim_I$ .

# $\text{End}(\mathbb{1}) \cong k$

---

Let  $\mathcal{S}$  be the set of splitting objects in  $\mathcal{T}$ .

- Define an equivalence relation on  $\mathcal{S}$  by  $Q \sim_I P$  if there exists  $X \in \mathcal{T}$  such that  $Q$  is a direct summand of  $X \otimes R$ .
- Then we can take the equivalence classes  $\overline{\mathcal{S}} = \mathcal{S} / \sim_I$ .
- Each  $c \in \overline{\mathcal{S}}$  generates a splitting ideal in  $\mathcal{T}$ .



$$\mathrm{End}(\mathbb{1}) \cong k$$

---

Let  $\mathcal{S}$  be the set of splitting objects in  $\mathcal{T}$ .

- Define an equivalence relation on  $\mathcal{S}$  by  $Q \sim_I P$  if there exists  $X \in \mathcal{T}$  such that  $Q$  is a direct summand of  $X \otimes R$ .
- Then we can take the equivalence classes  $\overline{\mathcal{S}} = \mathcal{S} / \sim_I$ .
- Each  $c \in \overline{\mathcal{S}}$  generates a splitting ideal in  $\mathcal{T}$ .

### Proposition

We have  $\mathrm{Ver}_{p^n} = \bigoplus \mathrm{Ver}_{p^n}(c)$  where  $\mathrm{Ver}_{p^n}(c)$  is the tensor subcategory of objects generated by  $c$ . Then  $\mathbb{1} = \bigoplus_{c \in \overline{\mathcal{S}}} \mathbb{1}_c$ .

$$\text{End}(\mathbb{1}) \cong k$$

---

Let  $\mathcal{S}$  be the set of splitting objects in  $\mathcal{T}$ .

- Define an equivalence relation on  $\mathcal{S}$  by  $Q \sim_I P$  if there exists  $X \in \mathcal{T}$  such that  $Q$  is a direct summand of  $X \otimes R$ .
- Then we can take the equivalence classes  $\overline{\mathcal{S}} = \mathcal{S} / \sim_I$ .
- Each  $c \in \overline{\mathcal{S}}$  generates a splitting ideal in  $\mathcal{T}$ .

### Proposition

We have  $\text{Ver}_{p^n} = \bigoplus \text{Ver}_{p^n}(c)$  where  $\text{Ver}_{p^n}(c)$  is the tensor subcategory of objects generated by  $c$ . Then  $\mathbb{1} = \bigoplus_{c \in \overline{\mathcal{S}}} \mathbb{1}_c$ .

### Proposition

$\mathcal{S}_{n,p}$  is the ideal only splitting ideal in  $\mathcal{T}_{n,p}$

# Abelian Envelope

---

## Theorem

*$\mathcal{T}$  admits a fully faithful monoidal functor  $E : \mathcal{T} \rightarrow \mathcal{D}$  into a multitensor category  $\mathcal{D}$  with enough projectives if and only if  $F : \mathcal{T} \rightarrow \mathcal{C}(\mathcal{T})$  is a fully faithful embedding. In this case  $\mathcal{C}(\mathcal{T})$  is the abelian Envelope of  $\mathcal{T}$ .*

As it turns out,  $F : \mathcal{T}_{n,p} \rightarrow \text{Ver}_{p^n}$  is a fully faithful embedding. Thus,  $\text{Ver}_{p^n}$  is the abelian envelope of  $\mathcal{T}_{n,p}$

# Frobenius Functor

---

Recall  $Fr : \mathcal{C} \rightarrow \mathcal{C}^{(1)} \boxtimes Ver_p$  is the Frobenius functor. Since  $\mathcal{T}_{n,p}$  and  $Ver_{p^n}$  are defined over the prime field we actually have  $Fr : Ver_{p^n} \rightarrow Ver_{p^n} \boxtimes Ver_p$

## Proposition

$Fr(F(\mathcal{T}_{n,p})) \subseteq Ver_{p^n} \boxtimes \mathbb{1} \cong Ver_{p^n}$ , so there is an inclusion  $\mathcal{T}_{n,p} \hookrightarrow Ver_{p^n}$ .

# Frobenius Functor

Recall  $Fr : \mathcal{C} \rightarrow \mathcal{C}^{(1)} \boxtimes Ver_p$  is the Frobenius functor. Since  $\mathcal{T}_{n,p}$  and  $Ver_{p^n}$  are defined over the prime field we actually have  $Fr : Ver_{p^n} \rightarrow Ver_{p^n} \boxtimes Ver_p$

## Proposition

$Fr(F(\mathcal{T}_{n,p})) \subseteq Ver_{p^n} \boxtimes \mathbb{1} \cong Ver_{p^n}$ , so there is an inclusion  $\mathcal{T}_{n,p} \hookrightarrow Ver_{p^n}$ .

$$\begin{array}{ccc}
 \mathcal{T}_{n,p} & \xrightarrow{Fr \circ F_n} & Ver_{p^n} \\
 \pi_{\mathcal{I}_{n-1}} \downarrow & \nearrow \tilde{H} & \uparrow \\
 \mathcal{T}_{n-1,p} & & \\
 F_{n-1} \downarrow & \nearrow H & \\
 Ver_{p^{n-1}} & & 
 \end{array}$$

As it turns out,  $Fr(F(St_{n-2})) \neq 0$  and  $Fr(F(St_{n-1})) = 0$ . Then the map  $\tilde{H}$  in the diagram to the left is induced by  $Fr \circ F$ .

# Frobenius Functor

Recall  $Fr : \mathcal{C} \rightarrow \mathcal{C}^{(1)} \boxtimes Ver_p$  is the Frobenius functor. Since  $\mathcal{T}_{n,p}$  and  $Ver_{p^n}$  are defined over the prime field we actually have  $Fr : Ver_{p^n} \rightarrow Ver_{p^n} \boxtimes Ver_p$

## Proposition

$Fr(F(\mathcal{T}_{n,p})) \subseteq Ver_{p^n} \boxtimes \mathbb{1} \cong Ver_{p^n}$ , so there is an inclusion  $\mathcal{T}_{n,p} \hookrightarrow Ver_{p^n}$ .

$$\begin{array}{ccc}
 \mathcal{T}_{n,p} & \xrightarrow{Fr \circ F_n} & Ver_{p^n} \\
 \pi_{\mathcal{I}_{n-1}} \downarrow & \nearrow \tilde{H} & \\
 \mathcal{T}_{n-1,p} & & \\
 F_{n-1} \downarrow & \nearrow H & \\
 Ver_{p^{n-1}} & & 
 \end{array}$$

As it turns out,  $Fr(F(St_{n-2})) \neq 0$  and  $Fr(F(St_{n-1})) = 0$ . Then the map  $\tilde{H}$  in the diagram to the left is induced by  $Fr \circ F$ .

## Corollary

For  $n \geq 2$  there is a natural inclusion  $Ver_{p^{n-1}} \hookrightarrow Ver_{p^n}$ .

# Properties/Recap

---

The category  $\text{Ver}_{p^n} \dots$

- is a finite symmetric tensor category (ie. artinian  $k$ -linear abelian rigid monoidal category with biexact tensor product and  $\text{End}(\mathbb{1}) \cong k$ )
- Has a fully faithful embedding  $F : \mathcal{T}_{n,p} \rightarrow \text{Ver}_{p^n}$
- is the abelian envelope of  $\mathcal{T}_{n,p}$ .
- has a Frobenius functor  $Fr : \text{Ver}_{p^n} \rightarrow \text{Ver}_{p^n}$
- has an embedding  $H : \text{Ver}_{p^{n-1}} \rightarrow \text{Ver}_{p^n}$
- is Frobenius (ie. projectives coincide with injectives)

# Spec

Since  $\text{Ver}_{p^n}$  is exact and Frobenius, the stable module category  $\text{Stab Ver}_{p^n}$  is triangulated. In fact, it is tensor triangulated.

- The category  $\text{Stab Ver}_{p^n}$  quotients out the ideal of projectives
- The category comes with a shift  $\Omega$ :

$$\begin{array}{ccccccc} P_3 & \longrightarrow & P_1 & \longrightarrow & P_0 & \longrightarrow & 0 \\ & \searrow \text{dashed} & & \nearrow \text{dashed} & & \searrow \text{dashed} & \\ & \Omega^2 X & & \Omega X & & X & \end{array}$$

- and *distinguished triangles*:

$$X \longrightarrow Y \longrightarrow Z \longrightarrow \Omega^{-1}X$$



# Spec

---

We then wish to determine the Balmer Spectrum of our triangulated category  $\text{Stab Ver}_{p^n}$ .

# Spec

We then wish to determine the Balmer Spectrum of our triangulated category  $\text{StabVer}_{p^n}$ .

## Definition

The *Balmer Spectrum* of an essentially small tensor triangulated ( $\Delta$ ) category  $\mathcal{K}$  is the set of primes of  $\mathcal{K}$  denoted  $\text{Spc}(\mathcal{K}) = \{\mathcal{P} | \mathcal{P} \text{ a prime in } \mathcal{K}\}$ .

A *Thick  $\otimes$ -ideal*  $\mathcal{A}$  of  $K$  is a triangulated full subcategory that is closed under  $\otimes$  and  $\oplus$ .

A *prime* of a  $\Delta$ -category is a proper thick  $\otimes$ -ideal  $\mathcal{P} \subsetneq \mathcal{K}$  such that for any  $a, b \in K$  if  $a \otimes b \in \mathcal{P}$ , then  $a \in \mathcal{P}$  or  $b \in \mathcal{P}$

A naive approach to this question would be to determine the fusion rules for non-projective indecomposables.

# Particular Cases

---

Fact ( $n = 1$ )

$\text{Ver}_p$  is semi-simple, so  $\text{Spc}(\text{Ver}_p)$  is trivial.

# Particular Cases

## Fact ( $n = 1$ )

$\text{Ver}_p$  is semi-simple, so  $\text{Spc}(\text{Ver}_p)$  is trivial.

## Fact ( $n = 2$ )

*The indecomposable modules of  $\text{Ver}_{p^2}$  can be read off from a Brauer tree algebra with each block corresponding to a line with  $p$  nodes and no exceptional vertex. It is likely not possible to find explicit fusion rules for non-simple indecomposables.*

$\text{Ver}_{p^2}$  of finite representation type  $\implies \text{Spc}(\text{Ver}_{p^2})$  is finite. Using

- *The support of an indecomposable object must be connected [Bal07]*
- *The tensor product of indecomposables is never 0*

*We see by the Krull-Schmidt property  $\text{Spc}(\text{Ver}_{p^2})$  is a point.*

Conjecture:  $\text{Ver}_8^+$

The category  $\text{Ver}_8^+$  is of tame representation type.

We suspect that the spectrum is  $\mathbb{P}^1$  since  $\text{Proj}(H^*(sl_2^{[p]})) = \mathbb{P}^1$  [BE21]

# Reference

---



Paul Balmer.

The spectrum of prime ideals in tensor triangulated categories, 2004.



Paul Balmer.

Supports and filtrations in algebraic geometry and modular representation theory.  
*Amer. J. Math.*, 129(5):1227–1250, 2007.



Dave Benson and Pavel Etingof.

Symmetric tensor categories in characteristic 2, 2020.



David Benson and Pavel Etingof.

On cohomology in symmetric tensor categories in prime characteristic, 2021.



Dave Benson, Pavel Etingof, and Victor Ostrik.

New incompressible symmetric tensor categories in positive characteristic, 2021.