

Cervical Cancer Prevention and Control in Women Living With Human Immunodeficiency Virus

Philip E. Castle, PhD, MPH ^{1,2}; Mark H. Einstein, MD, MS³; Vikrant V. Sahasrabudhe, MBBS, MPH, DrPH ¹

¹Division of Cancer Prevention, National Cancer Institute, National Institutes of Health, Rockville, Maryland; ²Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Rockville, Maryland; ³Department of Obstetrics, Gynecology, and Reproductive Health, Rutgers New Jersey Medical School, Newark, New Jersey.

Corresponding Author: Philip E. Castle, PhD, MPH, Division of Cancer Prevention, National Cancer Institute, National Institutes of Health, 9609 Medical Center Drive, Room 5E410, Rockville, MD 20850 (philip.castle@nih.gov).

DISCLOSURES: Mark H. Einstein was supported by the The AIDS Malignancy Consortium Grant (UM1CA121947) and the US-Latin American-Caribbean HIV/HPV-Cancer Prevention Clinical Trials Network Partnership Center Grant (U54CA242639) from the National Cancer Institute/National Institutes of Health. Philip E. Castle has received human papillomavirus tests and assays for research at reduced or no cost from Roche, Becton Dickinson, Cepheid, and Arbor Vita Corporation. Mark H. Einstein reports contracts from Johnson & Johnson, Pfizer, Inovio, AstraZeneca, and Merck (payment to institution for costs related to clinical trials); consulting fees from Papivax, Merck, PDS Biotechnologies, and Inovio (payment to institution for Dr. Einstein's time); unpaid committee work for the American Society for Colposcopy and Cervical Pathology and the Society of Gynecologic Oncology; and the National Cancer Institute and the World Health Organization provide payment for Dr. Einstein's time related to cervical cancer prevention. Vikrant V. Sahasrabudhe had no disclosures.

The opinions expressed by the authors are their own, and this material should not be interpreted as representing the official viewpoint of the US Department of Health and Human Services, the National Institutes of Health, or the National Cancer Institute.

doi: 10.3322/caac.21696. Available online at cancerjournal.com

Abstract: Despite being highly preventable, cervical cancer is the fourth most common cancer and cause of cancer death in women globally. In low-income countries, cervical cancer is often the leading cause of cancer-related morbidity and mortality. Women living with human immunodeficiency virus (HIV)/acquired immunodeficiency syndrome are at a particularly high risk of cervical cancer because of an impaired immune response to human papillomavirus, the obligate cause of virtually all cervical cancers. Globally, approximately 1 in 20 cervical cancers is attributable to HIV; in sub-Saharan Africa, approximately 1 in 5 cervical cancers is due to HIV. Here, the authors provide a critical appraisal of the evidence to date on the impact of HIV disease on cervical cancer risk, describe key methodologic issues, and frame the key outstanding research questions, especially as they apply to ongoing global efforts for prevention and control of cervical cancer. Expanded efforts to integrate HIV care with cervical cancer prevention and control, and vice versa, could assist the global effort to eliminate cervical cancer as a public health problem. *CA Cancer J Clin* 2021;71:505–526. © 2021 The Authors. *CA: A Cancer Journal for Clinicians* published by Wiley Periodicals LLC on behalf of American Cancer Society. This article has been contributed to by US Government employees and their work is in the public domain in the USA. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Keywords: Africa, cervical cancer, cytology, human immunodeficiency virus (HIV), human papillomavirus (HPV), low-income and middle-income countries (LMICs), prevention, screening, vaccination, visual inspection after acetic acid (VIA)

Cervical Cancer and HIV/AIDS as Intersecting Epidemics

Invasive cervical cancer is one of the leading causes of cancer-related morbidity and mortality in women globally. It is estimated that approximately 600,000 women are diagnosed with, and more than 300,000 women die from, cervical cancer worldwide annually.¹ A disproportionate amount of its global burden is experienced by women living in low- and middle-income countries (LMICs), where it is often the first or second leading cause of cancer cases and deaths in women.

With the discovery of an obligate causative infectious agent, high-risk human papillomavirus (HPV), elucidation of key steps in its natural history, and availability of highly effective primary and secondary prevention technologies, cervical cancer is an eminently preventable malignancy. Indeed, in countries with routine, effective cervical cancer screening and early detection programs, cervical cancer incidence and mortality rates have declined precipitously over the past one-half century.^{1,2} HPV vaccines will likely lead to even further reductions in decades to come. Buoyed by rising global optimism about the possibility of reducing cervical cancer globally, the World Health Organization (WHO), with endorsement from over 194 countries, including the United States, has recently launched a global initiative to accelerate the elimination of cervical cancer as a public health problem by significant

expansion of efforts to increase HPV vaccination to 90% coverage, screening to 70% coverage in mid-adult women, and treatment to 90% of those in need of it ("90-70-90").³⁻⁵

Over the past 4 decades, the human immunodeficiency virus (HIV)/acquired immunodeficiency syndrome (AIDS) pandemic has emerged and persisted as one of the world's most serious public health, development, and economic challenges. With an estimated 37.7 million (uncertainty bound, 30.2-45.1 million) people living with HIV (PLWH) worldwide in 2020,⁶ 53% of all PLWH were women and girls. HIV/AIDS disproportionately affects people in LMICs, where cervical cancer is also exceedingly common. More than two-thirds of PLWH live in sub-Saharan Africa (SSA).^{7,8} Untreated HIV leads to severe impairment of the immune system, increasing the risk of developing opportunistic infections as well as infection-related cancers and other chronic comorbidities that are otherwise rare in people with normal immune function.

Since the mid-1990s, the availability of combination antiretroviral therapy (ART) for HIV has led to significant improvements in the management of HIV disease, including a partial or near-complete restoration of immunocompetence, with long-term adherence to ART regimens and retention within the HIV care continuum. Over the past 2 decades, catalyzed by generous bilateral and multilateral public sector and philanthropic initiatives (eg, The President's Emergency Plan For AIDS Relief [PEPFAR]),⁹ ART has become affordable and accessible to millions of PLWH, including in some of the most resource-limited settings in SSA. This has led to dramatic increases in the lifespans of PLWH^{10,11} as well as several considerably improved trajectories in the incidence of HIV-associated opportunistic infections,¹² cancers,¹³⁻¹⁵ and comorbidities.¹⁶ In fact, in resource-rich areas of the world, the presence of HIV infection is thought of more like a *chronic* disease compared with pre-ART times in which HIV infection often led to terminal sequelae.

Women living with HIV (WLWH) have long been recognized as having a higher risk for the acquisition, persistence, and progression of high-risk HPV and its downstream consequences, including cervical cancer. In fact, invasive cervical cancer was first included as an AIDS-defining condition (ie, a marker of clinically relevant immunosuppression) by the US Centers for Disease Control and Prevention (CDC) in 1993.^{17,18} Yet, just like the wide heterogeneity seen in the incidence rates of cervical cancer among women in the general population globally, incidence rates in WLWH have also varied substantially between countries. Although incidence rates of cervical cancer in WLWH have been uniformly higher than those in women without HIV (HIV-negative women) within any given setting/geographic region, recent meta-analyses have demonstrated that the attributable fraction of cervical cancers because of HIV is much greater in

LMICs than in high-income countries (HICs).¹⁹ These differences are likely because of a combination of factors, such as a higher percentage women who are WLWH,⁶ differences in the duration²⁰ and adherence²¹ that affect the degree of immune restitution, various social determinants of health (eg, nutrition, housing, and transportation), and better access to cervical cancer screening services in HICs.²² Yet, unlike other AIDS-defining cancers (Kaposi sarcoma and specific types non-Hodgkin lymphoma, ie, Burkitt lymphoma, immunoblastic lymphoma, and primary lymphoma of the brain),²³ which have clearly demonstrated evidence linking dramatic declines with the availability of ART over the past 3 decades,¹⁴ cervical cancer rates have not declined by the same magnitude, and cervical cancer remains a significant comorbidity among WLWH regardless of the setting globally.

Although the relative importance of the 2 factors related to health care delivery, HIV care and cervical cancer screening services, is unclear and difficult to disentangle, their confluence likely contributes to the excessive cervical cancer burden in LMICs.

In this article, we attempt to provide a critical appraisal of the evidence to date on this topic while describing key methodologic issues inherent in studies contributing to this evidence. We also discuss the implications for clinical cancer prevention, especially as they apply to the ongoing global efforts for prevention and control of cervical cancer globally, because both the management of HIV disease and interventions for the prevention and control of cervical cancer (ie, vaccines, screening and treatment of precancerous lesions, and management of invasive cancers) can be conceived as the necessary *dual arms* of a multifocal public health strategy to reduce cervical cancer burden in WLWH.

Quantifying the Burden of Cervical Cancer in WLWH

Over the past 3 decades, there has been significant evolution in quantifying the burden of cervical cancer in WLWH by efforts using a variety of descriptive epidemiology approaches. These have included population registries/records-linkage studies that connect evidence from disparate sources, such as cancer registries, HIV registries/surveillance databases, and national death registers, as well as cohort and case-control studies from pooled databases of clinics and hospitals providing care for WLWH.

Results of 2 large meta-analyses conducted 14 years apart (2007 and 2021)^{19,24} are consistent in their reporting that cervical cancer incidence remains approximately 6 times higher in WLWH compared with the general population or HIV-negative women (pooled risk estimate from the most recent meta-analysis, 6.07; 95% confidence interval [95% CI], 4.40-8.37).¹⁹ In 2018, 5.8% (95% CI,

4.6%-7.3%) of new cervical cancer cases were diagnosed in WLWH, most of which were attributable to HIV infection (4.9%; 95% CI, 3.6%-6.4%).¹⁹ Yet the attributable risk varies greatly across regions and countries because of the very wide variations in HIV prevalence worldwide: <5% in 122 countries and >40% in 9 southern African countries (Eswatini, Lesotho, Botswana, South Africa, Zimbabwe, Namibia, Mozambique, Zambia, and Malawi).¹⁹ These high-burden countries also have some of the highest HIV prevalence (between 11% and 32%) in females aged 15 years or older, and ART has been available only over the past 15 years.²⁵⁻²⁷ Other key findings from the literature on quantification of the burden include:

1. Cervical cancer incidence rates in WLWH in HICs (in contrast to those in LMICs) have declined steadily with access to ART as well as wider availability of cervical cancer screening over the past 3 decades.¹³⁻¹⁵ The age-standardized incidence rate of cervical cancer in US WLWH is predicted to decrease from 60 per 100,000 in 2006 to 10 per 100,000 in 2030.¹⁵
2. Among WLWH, those with higher CD4-positive T-cell levels have a lower risk of cervical precancer and cancer.²⁸⁻³⁰
3. Although cervical precancer among WLWH is common, WLWH who receive regular cervical screenings have low incidence rates of invasive cervical cancer.^{29,31}
4. The magnitude of the effect of precancer risk varies with differences in the use of ART regimens and precancer detection methods.^{28,29}
5. For US WLWH who receive adequate cervical screening and follow-up care for positive screening results, incidence rates of cervical cancer are projected to converge with those of the general population within the next decade.¹⁵
6. WLWH non-White racial and ethnic subgroups in both HICs^{32,33} and LMICs^{34,35} experience significant health disparities because of systemic racism, differential access to health care, and/or other social determinants of health, resulting in higher risks of cervical cancer incidence and mortality compared with their White counterparts. Although these underserved groups are minorities in HICs, they are often the majority population in LMICs.

Therefore, HIV infection and the subsequent disparate care are important contributors to the unequal burden of cervical cancer in some of the lowest resource settings globally.²⁵⁻²⁷ Prevention and control of HIV infection and improved focus on patient-centered care resources to complement efforts for prevention and control of cervical cancer will have a profound impact on reducing the burden of cervical cancer globally.

Impact of HIV and HIV Control on HPV Natural History

There are 4 key stages in the development of cervical cancer: HPV acquisition, HPV persistence, progression to cervical precancer, and development of invasive cancer.³⁶ In addition to factors such as age, parity, smoking status, and hormonal contraceptive use, HIV coinfection, although not causal, has a profound impact on several steps in the natural history of HPV (HPV cofactor) that increases HPV-related cancer risk in WLWH compared with HIV-negative women.

ART modifies these associations by mitigating HIV effects on the immune system. However, the evolution of ART regimens over the past 3 decades has often complicated uniform interpretations of the effect of ART across studies. Furthermore, a one-time measurement of ART status is an imperfect surrogate of adherence to ART and retention in the HIV care cascade over the long-term, which are key factors for HIV viral control and immune restitution. In addition, the development of HIV resistance to ART regimens can reduce their effectiveness for immune reconstitution despite adequate adherence.^{37,38} Variations in methods to detect cervical disease status due to variability in screening and diagnostic tests (especially seen with visual screening approaches and cytologic/histopathologic interpretations) also affect the interpretation of these associations. Finally, in the context of a dynamic immune milieu with swings between relative immunosuppression and immune restoration states in WLWH, cervical HPV detection may represent reactivation of latent/quiescent infections³⁹ as opposed to new infections, as also observed in studies among women undergoing long-term immunosuppressive therapy after solid organ transplants.^{40,41}

Notwithstanding these significant methodologic limitations, meta-analyses of the available literature suggest that WLWH have a higher risk of HPV acquisition, presumably due to shared behavioral risk of sexual transmission but also likely due to immune dysregulation (presumably because more incident HPV infections, even those destined to clear, persist long enough to be detected^{42,43}), as evidenced by the inverse association between HPV acquisition risk and markers of HIV control/immune status (eg, CD4 counts).⁴³⁻⁴⁵ WLWH have an approximately ≥ 2 -fold prevalence of HPV compared with HIV-negative women.⁴⁶⁻⁵² Among WLWH, HPV prevalence was lower among those receiving ART (vs not),^{51,53,54,55,56,57,58,59} on longer (vs shorter) duration of ART,^{20,54} with higher (vs lower) CD4 counts,^{51,54,55,56,57,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75} and with a lower HIV viral load.^{58,70,71}

In multiple HPV type infections, individual HPV type infections likely act independently rather than synergistically and thus contribute additively to the cervical cancer risk. A few studies in the general population have found that multiple HPV infections with non-HPV16 types increase

the risk of cervical precancer and cancer.^{76,77} WLWH are from 2-fold to 3-fold more likely to have multiple HPV infections (among HPV-positive women) compared with HIV-negative women.^{47,48,49,52} Receiving ART⁵⁶ and having a lower HIV viral load⁷⁸ and/or higher CD4 counts^{73,78} are associated with fewer multiple HPV infections.

HIV infection increases the likelihood of cervical HPV infection persisting and, given its persistence, progressing to precancer, and it reduces the likelihood of clearance/regression of low-grade and high-grade cervical abnormalities.⁴⁴ A recent meta-analysis⁵³ reported that ART was associated with reduced incidence of incident high-grade cervical abnormalities (adjusted odds ratio, 0.59), progression from low-grade to high-grade cervical abnormalities (adjusted odds ratio, 0.64), and an increased regression of cervical abnormalities (adjusted hazard ratio [HR], 1.54). Most importantly, data from 2 studies included in that meta-analysis demonstrated that ART was associated with a reduced risk (adjusted HR, 0.50) of invasive cervical cancer.⁵³ Likewise, low CD4 counts were strongly associated with cervical cancer risk,⁷⁹ and low CD4 counts at the time of ART initiation were also associated with an increased risk of cervical cancer.³⁵ Overall, the age at cervical cancer diagnosis in WLWH is approximately 5 to 10 years younger than that in HIV-negative women,^{33,80,81,82,83,84} highlighting the possible role of HIV coinfection as an *accelerant* in cervical carcinogenesis.

It is less clear whether HIV infection and immune status influence the last step of HPV-related carcinogenesis: cervical precancer developing into invasive cancer. It is unethical to observe the progression of cervical precancer into cancer, as was observed tragically in the “The Unfortunate Experiment”: from 1955 to 1976, New Zealand’s National Women’s Hospital withheld treatment from women diagnosed with cervical precancer.⁸⁵ Over the next 30 years, approximately 30% of these women with untreated cervical precancer developed cervical cancer.⁸⁵ Fortunately, no such experiment has been done among WLWH diagnosed with cervical precancer.

There are no high-quality, well powered, population-based, cross-sectional studies comparing the prevalence ratio of cervical intraepithelial neoplasia grade 3 (CIN3)/adenocarcinoma in situ (AIS) and invasive cervical cancer as a proxy for invasive potential in WLWH versus HIV-negative women. Such an approach (vs comparing between separate studies of WLWH or HIV-negative women with different methods, etc), by using statistical methods to control for the effects of differences in screening, variability of pathology diagnosis,⁸⁶ sociodemographic determinants, and behavioral and cultural factors, might provide some indication of whether there are gross differences in the invasive potential of cervical precancer according to HIV status and the degree of immune reconstitution. HPV genotyping data might

provide insights into how these factors influence HPV genotype-specific invasive potential, as has been shown in a meta-analysis of data from the general population.⁸⁷ A few studies report lower CD4 counts in women diagnosed with cervical cancer versus those with precancerous abnormalities,^{71,88} suggesting that immunity may play a role in this final step in cervical carcinogenesis.

Interestingly, of the few studies that have reported on stage distribution of cervical cancer, most did not find a significant difference in cancer stage at diagnosis in WLWH compared with HIV-negative women^{83,89,90,91}; one study reported more stage IIIB cervical cancer in WLWH than in HIV-negative women.⁹² The general lack of a difference in cancer stage distribution between WLWH and HIV-negative women may be a result of the early stage presentation of symptoms, such as postcoital bleeding in both, and/or because WLWH are generally under closer clinical surveillance than HIV-negative women.

There are some data that point to differences in the HPV type-specific natural history in WLWH compared with HIV-negative women. First, the types most commonly found in HIV-negative women, notably HPV16, are the least affected by the immune status in WLWH.⁹³ Second, HPV16 is less commonly detected in CIN3 diagnosed in WLWH than in HIV-negative women.⁹⁴ Finally, there is some evidence to suggest that HPV16 is somewhat less dominant in cervical cancers diagnosed in WLWH than in HIV-negative women.^{95,96} Such type differences might play a role in the time to develop HPV-associated disease, because women with HPV16-related (and HPV18-related) cervical cancers diagnosed in the general population are approximately 5 years younger than women with cervical cancers related to other types.⁹⁷ These differences also could reduce the effectiveness (eg, positive predictive value) of HPV16 and HPV18 genotyping for clinical decision making, as recommended in the general population, in the management of HPV-positive WLWH.⁹⁸

These data highlight the importance of using ART in cervical cancer risk reduction. Counterintuitively, PLWH live longer on ART,^{10,11} giving them more time to develop HPV-related cancers, which may partially offset its cancer risk-reducing benefits.^{99,100} This may explain the time trend of increasing cervical cancer risk in WLWH compared with HIV-negative women observed in South Africa.¹⁰¹ Thus the integration of prevention strategies with the delivery of ART to WLWH will maximize the cervical cancer prevention benefits of ART, as discussed below.

Primary Prevention: Prophylactic HPV Vaccination

The discovery of high-risk HPV infections as the necessary cause of cervical cancer has led to revolutionary advances in cervical cancer prevention, including the development of

prophylactic HPV vaccines for primary prevention. Current prophylactic HPV vaccines are based on the self-assembly of recombinantly expressed L1 protein in cell lines into virus-like particles that resemble native viral capsids but without the viral genome necessary for viral replication.

First-generation HPV vaccines, Gardasil (Merck & Company, Kenilworth, New Jersey)¹⁰² and Cervarix (GlaxoSmithKline, Rixensart, Belgium),¹⁰³ target HPV16 and HPV18 (HPV16/HPV18), which cause approximately 70% of cervical cancers. Gardasil also targets HPV6 and HPV11 (HPV6/HPV11), which are non-high-risk HPV types responsible for 90% of anogenital warts (*Condyloma acuminata*). The next-generation HPV vaccine, Gardasil 9 (Merck & Company),¹⁰⁴ targets HPV31, HPV33, HPV45, HPV52, and HPV58 in addition to HPV6, HPV11, HPV16, and HPV18 and is predicted to prevent approximately 90% of cervical cancers. Notably, the same types of HPV cause cervical cancer in approximately the same proportions in every region of the world, with the exception of Africa, as discussed below.¹⁰⁵

Because HPV vaccination is only prophylactic and not therapeutic, the ideal timing of HPV vaccination is before sexual initiation and exposure to HPV (ie, HPV-naïve women). On a population level, this can be achieved by vaccinating a few years before the population median age of sexual initiation. Because the median age of sexual initiation in many countries is typically from 15 to 17 years, the WHO recommends vaccination programs to target girls aged 9 to 13 years,¹⁰⁶ and the CDC recommends routine HPV vaccination for girls (and boys) aged 11 to 12 years.¹⁰⁷ With older cohorts of women, prophylactic vaccine is equally efficacious but less effective; ie, older age does not reduce vaccine efficacy in HPV-naïve individuals but, because there are fewer HPV-naïve women at older ages, overall vaccine effectiveness and cost-effectiveness of prophylactic vaccination decreases.¹⁰⁸⁻¹¹⁰ On the basis of immunogenicity data, the WHO¹¹¹ and the CDC^{95,112} recommend 2-dose schedules for those younger than 15 years and 3-dose schedules for those aged 15 years and older for all HPV vaccines. There is increasingly strong evidence that a single dose of HPV vaccine is sufficient to confer protection equivalent to multiple doses.^{113,114}

Several countries, eg, Australia,¹¹⁵⁻¹²¹ Scotland,¹²² Denmark,¹²³ and the United States,¹²⁴⁻¹²⁶ were early adopters of HPV vaccination and have documented population reductions in HPV infections and HPV-related diseases and abnormalities. Australia, for example, implemented vaccination with Gardasil in 2007, shortly after its US Food and Drug Administration approval and after a public awareness campaign. Their HPV vaccination program for females achieved >70% coverage in its first year and in subsequent years.¹¹⁵ Within several years after their national HPV vaccination was implemented, there was a significant decrease in anogenital warts among women and heterosexual men (because of herd protection), but not among men who have sex

with men.¹¹⁶ Subsequently, there have been documented decreases in the prevalence of HPV16 and HPV18 in vaccinated women,^{117,118} and now in unvaccinated women (because of herd protection),¹¹⁸ and in the prevalence of high-grade cervical abnormalities, first in younger women^{119,120} and now in slightly older women,¹²¹ as HPV-vaccinated cohorts have aged.¹²⁷ A meta-analysis on the impact of HPV vaccination found reductions in anogenital warts, HPV infections, and CIN grade 2 (CIN2) or more severe (CIN2+) diagnoses among girls and women and in diagnoses of anogenital warts among girls, women, boys, and men.¹²⁸ Recent reports from Finland,¹²⁷ Sweden,¹²⁹ and Denmark¹³⁰ now provide real-world evidence that HPV vaccination prevents cervical cancer.

HPV vaccination in PLWH has been well tolerated and safe and has resulted in good immune responses. One study of Gardasil¹³¹ in children living with HIV aged 7 to 12 years reported high-seroconversion, 4-year persistent, HPV type-specific immunity¹³² and immune memory cells¹³³ and a significant increase in, and persistence of, antibody titers (anamnestic response) after an additional booster (fourth dose).¹³² A study of Gardasil in adult men living with HIV also found high seroconversion in addition to a good safety profile.¹³⁴ Another study reported a seroconversion rate similar to that of Gardasil in WLWH and HIV-negative women aged 13 to 27 years.¹³⁵

Some studies have noted an impact of HIV disease status (CD4 counts and viral suppression) on immune responses to HPV vaccination. One study reported lower HPV seroconversion and antibody titers in young adult WLWH not taking ART compared with those taking ART.¹³⁶ Another study reported lower seroconversion and antibody titers in adult WLWH who had lower (vs higher) CD4 counts.¹³⁷ A third study reported that peak antibody titers after Gardasil vaccination were 2-fold to 3-fold higher in mid-adult WLWH with full HIV viral suppression versus those without viral suppression.¹³⁸

Two studies have compared the immune response to Cervarix versus the response to Gardasil in adult PLWH. Both studies found that the antibody titers after Cervarix immunization were superior to those after Gardasil.^{139,140} However, it remains unclear whether this difference in antibody titers translates into meaningful differences in efficacy or duration of protection.¹⁴¹

Immunogenicity studies in WLWH have been of insufficient sample size to address efficacy/effectiveness in preventing clinical disease. Studies of HPV vaccination in select PLWH at high risk of anal cancer have been limited in their effective sample size because of the high degree of anal HPV exposure (and likely misclassification of exposure) to targeted HPV genotypes before enrollment¹⁴² and the possible reactivation of previously acquired, latent HPV infection, such that few truly incident events can be observed.

Another challenge in conducting placebo-controlled randomized clinical trials to demonstrate efficacy of HPV vaccines

is that projected effect sizes are attenuated by lower-than-expected number of endpoints in placebo-arm participants, who may be practicing lesser risky behaviors with fewer exposures during the course of the study (presumably with adequate counseling and awareness of risk in relatively controlled clinical trial settings). This may result in studies being stopped as per pre-defined rules for futility, as evidenced in a recent trial of Gardasil in older men (aged 27 years and older) to prevent anal HPV infection or anal high-grade squamous intraepithelial lesions.¹⁴³ Yet, this same study suggested a signal of potential benefit for protection against oral HPV that is now under further investigation (ClinicalTrials.gov identifier NCT04255849).

Another study of Gardasil vaccination in perinatally HIV-infected versus HIV-exposed, uninfected youth reported a 5-fold higher incidence of abnormal cervical cytology among the former, suggesting that HPV vaccination is likely less effective in the context of perinatally acquired HIV infection.¹⁴⁴ Not surprisingly, HPV vaccination with Gardasil either before¹⁴⁵ or after¹⁴⁶ excision of high-grade cervical/anal abnormalities in persons with HIV, did not prevent recurrence of lesions,^{145,146} and it seems unlikely that Gardasil 9 will either (ClinicalTrials.gov identifier NCT03284866). However, HPV vaccination should prevent newly acquired (not latent) infection by HPV vaccine-targeted types acquired posttreatment from developing into new (vs recurrence of the same, type-specific) high-grade cervical abnormalities.

A recent study compared the effectiveness of catch-up HPV vaccination (ages 13–26 years) in women with history of immunosuppression (mostly HIV) versus those without a history of immunosuppression.¹⁴⁷ HPV vaccination significantly reduced the risk of CIN2+ by 19% in those without a history of immunosuppression, whereas the risk was not significantly reduced in those with such a history.

There is still a need to assess the efficacy, effectiveness, and duration and determinants of protective immunity of prophylactic HPV vaccines in PLWH. Given the immunodeficiencies related to HIV infection and the lack of defined immune correlates of protection, it is unclear whether immunogenicity, as measured by antibody or neutralizing antibody titers, is a good proxy for the efficacy and long-term effectiveness of HPV vaccines in PLWH. Some key outstanding questions for HPV vaccination in PLWH include: 1) how many doses are needed for protective immunity, and does an adjuvanted vaccine (eg, AS04 adjuvant in Cervarix) improve the protective immunity such that fewer doses are needed to achieve long-lasting, protective immunity compared with unadjuvanted HPV vaccines (eg, Gardasil); 2) does protective immunity wane; and 3) do these vaccines provide sufficiently broad coverage against the HPV types that cause cancer in PLWH?

As noted, there is some evidence to suggest that the distribution of HPV types that cause cervical cancer in WLWH is somewhat different from that in the general population/

HIV-negative women such that HPV16 is somewhat less predominant in cervical precancer and cancer diagnosed in WLWH.^{95,96} HPV35, a type in the same phylogenetic group (α -9) as HPV16, is particularly common in cervical cancers diagnosed among WLWH living in Africa,⁹⁵ precancers diagnosed among WLWH living in SSA and Sweden,¹⁴⁸ and precancers diagnosed among WLWH and HIV-negative women living in South Africa.¹⁴⁹ This may be due in part to viral variants of HPV35 that cause more cancer in women of African descent than in women of other ethnicities.¹⁵⁰ Whether there are any independent effects of HIV coinfection and African descent on the HPV type distribution in cervical cancer is uncertain given the high proportion of WLWH who are of African descent. Regardless, current HPV vaccines might be less effective in preventing cervical cancer in WLWH and/or in women of African descent than in other populations. Of note, there is already some evidence that Cervarix, which targets only HPV16 (vs HPV16, HPV31, HPV33, HPV52, and HPV58 in Gardasil 9) among α -9 types, induces some immunity, cross-protection, and herd immunity against HPV35,^{151–153} including immunity in WLWH.¹⁵⁴

The optimal HPV vaccination strategy for long-term protection against cervical cancer in WLWH, especially those living in LMICs, has yet and needs to be established. The US Advisory Committee on Immunization Practices recommends 3-dose HPV vaccination (at 0, 1 or 2, and 6 months) for females and males aged 9 to 26 years who have primary or secondary immunocompromising conditions.¹⁵⁵ However, as noted by the 2017 American Society of Clinical Oncology (ASCO) resource-stratified recommendations, HPV vaccination of WLWH is a D-level recommendation (ie, *insufficient* evidence quality, *weak* strength of recommendation).¹⁵⁶

The use of more viable end points, such as short-term HPV persistence, which is strongly predictive of high-grade cervical abnormalities^{157,158} and is accepted as a trial end point,¹⁵⁹ will increase the feasibility of studies to answer these questions. However, the use of HPV persistence as an end point must be interpreted with caution because of the possibility of reactivation of latent HPV infections, which may have obfuscated the benefits of HPV vaccination in clinical trials in adult WLWH who were previously exposed to HPV. This will be less of an issue for trials in which HPV vaccines are given before sexual initiation, at a time when the vaccines will provide the greatest benefit regardless of HIV status. Nevertheless, a long-term follow-up of WLWH participants in a trial that used 6-month HPV persistence as its end point will be important to confirm the vaccine effectiveness.

If the next generation of HPV prophylactic vaccines can produce broad-spectrum immunity, either by adding a more active adjuvant than the aluminum-based adjuvant to 9 HPV types included in Gardasil 9 or by expanding the types

included with the AS04 adjuvant used with Cervarix, it will be important to establish their long-term effectiveness and the number of doses needed to achieve that protection in WLWH living in LMICs, perhaps the most vulnerable population. At least one such vaccine is under development.¹⁶⁰

Secondary Prevention: Cervical Screening, Management, and Precancer Treatment

New guidelines¹⁶¹ from the WHO on cervical cancer screening and treatment recommend the use of HPV DNA testing as the primary cervical cancer screening test, although visual inspection after acetic acid (VIA) and Papanicolaou (Pap) tests may continue to be used according to local guidelines in various settings when HPV DNA testing is not *operational*. These guidelines recommend HPV testing-based screening for women aged 30 to 49 years every 5 to 10 years. For WLWH, these guidelines recommend HPV testing-based screening for women aged 25 to 49 years every 3 to 5 years. These are similar to recent resource-stratified guidelines for cervical cancer screening from the ASCO that recommend screening WLWH twice as frequently using HPV testing as what is recommended for HIV-negative women and the general population within the resource strata.¹⁶²

Studies have demonstrated unequivocally that testing for HPV DNA is more effective than Pap testing¹⁶³⁻¹⁶⁶ and VIA¹⁶⁴ in the prevention of cervical cancer. In a cluster randomized trial of approximately 100,000 women aged 30 to 59 years living in India,¹⁶⁴ a single round of HPV DNA testing reduced cervical cancer-related mortality by 50% in 8 years (HR, 0.52; 95% CI, 0.33-0.83), whereas Pap testing (HR, 0.89; 95% CI, 0.62-1.27) and VIA (HR, 0.86; 95% CI, 0.60-1.25) did not reduce cancer-related mortality compared with the community control. Moreover, a negative HPV test provided approximately 4-fold greater reassurance than negative Pap and VIA results against incident cervical cancer (3.7, 15.5, and 16.0 per 100,000 person years, respectively). In a pooled analysis of 4 randomized control trials in Europe that included approximately 175,000 women aged 20 to 64 years who were assigned to either HPV or cytology,¹⁶⁵ HPV testing reduced the incidence of invasive cervical cancer by 40% compared with cytology (rate ratio, 0.60; 95% CI, 0.40-0.89), and a negative HPV test was 70% more reassuring against cancer than cytology (rate ratio, 0.30; 95% CI, 0.15-0.60) over a 6.5-year follow-up period.

As a result of its superior sensitivity for cervical precancer/cancer and reassurance against incident cervical cancer after a negative screen, HPV testing is increasingly being adopted, recommended, and implemented as the primary (standard-of-care) cervical screening method in HICs, including the United States, despite the preexistence of highly successful Pap screening programs.¹⁶⁷⁻¹⁶⁹ The US Preventative Services Task Force¹⁷⁰ and the American Cancer Society¹⁷¹ have

recommended HPV testing as the acceptable and preferred option, respectively, for cervical cancer screening, and it has been endorsed by several professional medical societies, including the Society of Lower Genital Tract Diseases (formerly known as the American Society of Colposcopy and Cervical Pathology),¹⁷² ASCO,¹⁶² The American College of Obstetrics and Gynecology,¹⁷³ and the Society for Gynecologic Oncology.¹⁷⁴ Several European countries have implemented or are implementing HPV testing for cervical cancer screening.¹⁷⁵ Notably, HPV testing for cervical cancer screening is being recommended for LMICs.^{176,177}

One important advantage of HPV testing-based screening over other screening modalities is the possibility of using self-collected cervicovaginal specimens rather than provider-collected specimens. HPV testing of self-collected specimens minimizes the reliance on clinic-based screening and overcomes barriers such as geographic inaccessibility, lack of an adequate health care provider workforce, the need for a pelvic examination, or sociocultural issues that have been historically associated with lack of screening or underscreening. A recent meta-analysis reported that there was virtually no difference in the clinical performance for the detection of cervical precancer and cancer between the 2 specimen types when a polymerase chain reaction-based HPV test (eg, GP5+/6+; cobas [Roche Molecular Systems, Pleasanton, California]; and Onclarity [Becton Dickinson, Franklin, New Jersey]) was used (relative sensitivity [vs provider-collected specimen], 0.99; 95% CI, 0.96-1.02; relative specificity [vs provider-collected specimen], 0.98; 95% CI, 0.97-0.99).¹⁷⁸ By comparison, there was a slight decrement in the clinical performance when a signal amplification-based HPV test (eg, Hybrid Capture 2 [Qiagen, Germantown, Maryland]) was used (relative sensitivity [vs provider-collected specimen], 0.86; 95% CI, 0.76-0.98; relative specificity [vs provider-collected specimen], 0.97; 95% CI, 0.95-0.99).¹⁷⁸ Women were more likely to self-collect when a kit was mailed to their home than to undergo routine screening prompted by invitation/reminder letters (relative participation, 2.33; 95% CI, 1.86-2.91).¹⁷⁸ However, there is a paucity of data on the comparative clinical performance and acceptability of self-collected versus provider-collected specimens for HPV testing in WLWH.

Given its clinical performance and reliability, HPV testing has significant potential for widespread adoption and scalability across LMICs, but only if robust, LMIC-ready HPV tests are made affordable and available through innovative financing and procurement mechanisms. Currently, there are 2 promising polymerase chain reaction-based HPV tests for use in LMICs, the Xpert HPV test (Cepheid, Sunnyvale, California) and the AmpFire Multiplex High Risk HPV Real Time Fluorescent Detection with HPV16/18 Genotyping (Atila BioSystems, Mountain View,

TABLE 1. Simulation of the Impact of HIV Control on Human Papillomavirus Test Performance for the Detection of Cervical Precancer and Cancer (Cervical Intraepithelial Neoplasia Grade 3 or More Severe Diagnoses [CIN3+])^a

WLWH WITHOUT HIV CONTROL	CIN3+ (PREVALENCE: 5%)	<CIN3	TOTAL	PREDICTIVE VALUES
HPV test result				
Positive	450	4550	5000	PPV, 9.0%
Negative	50	4950	5000	NPV, 99.0%
Total	500	9500	10,000	
	Sensitivity, 90.0%	Specificity, 52.1%		
WLWH WITH HIV CONTROL	CIN3+ (PREVALENCE: 2.5%)	<CIN3	TOTAL	PREDICTIVE VALUES
HPV test result				
Positive	225	2275	2500	PPV, 9.0%
Negative	25	7475	7500	NPV, 99.7%
Total	250	9750	10,000	
	Sensitivity, 90.0%	Specificity, 76.7%		
SIMULATION OF US POPULATION FOR REFERENCE	CIN3+ (PREVALENCE: 0.5%)	<CIN3	TOTAL	PREDICTIVE VALUES
HPV test result				
Positive	45	955	1000	PPV, 4.5%
Negative	5	8995	9000	NPV, 99.9%
Total	50	9950	10,000	
	Sensitivity, 90.0%	Specificity, 90.4%		

Abbreviations: <CIN3, a diagnosis of less than grade 3 cervical intraepithelial neoplasia; CIN3+, cervical intraepithelial neoplasia grade 3 or more severe diagnoses; [CIN3+]; HPV, human papillomavirus; NPV, negative predictive value; PPV, positive predictive value; WLWH, women living with HIV.

^aTypically, prevalence of the disease only influences PPV and NPV, not sensitivity and specificity. However because immune status influences both the prevalence of disease (ie, CIN3+) as well as the test biomarker (ie, HPV), the specificity of HPV to detect CIN3+ is indeed influenced by immune status. For this simulation, we assume: 1) the prevalence of HPV in WLWH without HIV control (eg, no antiretroviral therapy) is 50% (top), in WLWH with HIV control is 25% (middle), and the general US population of women (for reference) is 10% (bottom); 2) the prevalence of CIN3+ in WLWH without HIV control (eg, no antiretroviral therapy) is 5% (top), in WLWH with control is 2.5% (middle), and the general US population of women (for reference) is 0.5% (bottom); and 3) the sensitivity of HPV testing for cervical precancer and cancer is 90% and does not differ by immune status or HIV status.

California). The Xpert HPV test is a random-access, rapid (60 minutes), WHO-prequalified¹⁷⁹ HPV test run on the multianalyte GeneXpert clinical laboratory platform; however, a cost of almost \$15 per test is not generally affordable for widespread use in LMICs.¹⁸⁰ AmpFire is an isothermal amplification HPV test but currently lacks sufficient evidence of clinical performance for regulatory approvals and adoption, and its commercial price is unknown.

HPV testing has shown similar sensitivity for high-grade cervical abnormalities among WLWH and HIV-negative women but has lower specificity because of the higher HPV prevalence, which leads to more HPV-positive women in need of follow-up care. Head-to-head comparisons have demonstrated that HPV testing is more sensitive but less specific than VIA and/or Pap testing for the detection of high-grade cervical abnormalities.^{57,181,182,183,184,185,186,187} Importantly, a negative HPV test, even in WLWH, provides excellent reassurance against high-grade cervical¹⁸⁷⁻¹⁸⁹ and anal¹⁹⁰ abnormalities and cancers.

HPV test specificity is influenced by HPV prevalence, which, in turn, is influenced by HIV prevalence and control, as previously discussed¹⁹¹ and illustrated in Table 1.

As noted above, among WLWH, HPV positivity decreases and HPV test specificity increases with a higher CD4 count,^{51,54,55,56,57,60,61,62,63,64,65,66,67,68,69,70,71} ART use and duration,^{51,53,54,55,56,57,58,59,192} and lower HIV viral load.^{58,70,71} One study reported that national implementation of ART for PLWH in Rwanda led to a time trend of higher CD4 counts and lower HPV prevalence.⁶⁷

Likewise, VIA and Pap test specificity is influenced by HIV infection and its control, probably as a result of their influence on HPV prevalence, as shown in one study that controlled for the presence of HPV.¹⁹³ VIA positivity is higher (and its specificity is lower) for WLWH (vs HIV-negative women)^{182,194,195,196,197,198,199,200} and, among WLWH, for those with lower CD4 counts^{183,201} and lack of ART use.⁵⁷ Pap test positivity is higher (and its specificity is lower) for WLWH (vs HIV-negative women)^{71,202} with higher CD4 counts^{71,183,193,203,204,205,206,207,208} and lower HIV viral loads.^{71,205}

Therefore, although there is not consistency across all studies about which types of biomarkers^{44,88} (eg, absolute CD4 count, CD4 percentage of total T-cell count, and HIV viral load) or clinical markers (eg, ART status, duration of ART use, and ART adherence measures) of HIV disease

status/immune competency are most strongly associated with the accuracy of cervical cancer screening tests, and not all studies have reported or controlled for all such markers, there is consistent evidence that HIV control in general improves cervical cancer screening test performance by improving its specificity. The implication of these data is that, with good HIV control, fewer WLWH will screen positive, which is especially important in LMICs with limited capacity and resources to manage them. Conversely, with uncontrolled HIV, the benefits versus harms of HPV testing are less certain not only because the high HPV positivity results in poor specificity, leading to overtreatment, but because of the increased burden on limited, financially strapped health care systems to manage patients who have positive results.

A secondary triage test of HPV-positive WLWH can be included to increase specificity but at the price of reducing sensitivity for high-grade cervical abnormalities as well as increasing cost, complexity, and burden on the health care system. Studies have evaluated visual triage (VIA or visual inspection with Lugol iodine),^{181,184,187,209,210} cytology,^{187,210,211} p16/Ki-67 dual-stain immunocytochemistry,²⁰⁹ HPV genotyping/type restriction,^{181,187,210,211,212,213} changing the positive cutoff point/viral load,²¹³ and host^{214,215} and viral methylation²¹⁶ biomarkers measured from the cervical specimen. There is no consensus approach to choosing which triage test to use for HPV-positive WLWH, and the choice may depend on local capacities and which HPV test is being used for primary screening. A large study of screening and triage strategies in 5000 WLWH living in Rwanda may provide important data on triage strategies for HPV-positive WLWH.²¹⁷

Although no studies are published on the impact of HIV control on the performance of colposcopy, it is reasonable to expect that greater HIV control will lead to fewer and smaller visible acetowhite abnormalities, like what is observed for VIA, because of fewer and better controlled prevalent HPV infections. Whether better HIV control might result in poorer colposcopic performance will likely depend on the skill of the colposcopist and their willingness to take multiple biopsies, which increases the sensitivity of colposcopic-directed diagnoses.^{218,219}

The treatment of cervical cancer precursors also is affected by HIV. Treatment by any method is more likely to fail in WLWH than in HIV-negative women.²²⁰ In HICs, the primary treatment modality is excision (loop electrosurgical excision procedure, large loop excision of the transformation zone, or cone excision) of histologically confirmed CIN2, CIN3, or AIS. Margins with disease (*positive margins*) after excision are a strong predictor of recurrent (incomplete treatment) disease in the general population²²¹ and also in WLWH.^{220,222,223} As a result of having larger areas of abnormality, WLWH are more likely to have positive margins^{224,225} and recurrence^{220,224,226,227,228,229,230} than

HIV-negative women. Among WLWH, a higher CD4 count,^{224,231,232,233,234,235} ART before treatment,²³⁶ and a lower HIV viral load²²² are associated with a lower risk of recurrence.

Where the capacity for excising cervical abnormalities is lacking, primarily in low-resource settings, tissue ablation by cryotherapy or thermal ablation is the primary method of treating cervical abnormalities. Tissue ablation, which potentially does less destructive damage to the cervix, may have a role in the treatment of smaller lesions or persistent HPV infection in WLWH,²³⁷ especially within single-visit *screen-and-treat* programs that seek to improve operational efficiencies by reducing attrition. Those cervical abnormalities that cover >75% of the cervical transformation zone, that are suspicious of cancer, and/or that cannot be fully visualized are deemed ineligible for ablative treatment and are referred for excisional treatment or cancer treatment, as needed.²³⁸

WLWH are more likely than HIV-negative women to have an ablation-ineligible cervical abnormality,^{239,240} again suggesting that WLWH have larger, harder-to-treat cervical abnormalities than HIV-negative women. Low CD4 counts^{241–243} and high HIV viral loads²²² in WLWH are associated with the recurrence of cervical abnormalities after treatment. Loop electrosurgical excision procedure/large loop excision of the transformation zone has shown more efficacy in treating CIN2+ than cryotherapy in WLWH, especially in those with CD4 counts <250/mm³, perhaps because of the size of the abnormality,²⁴⁴ indirectly supporting WHO recommendations to refer larger cervical abnormalities for excisional rather than ablative therapy.²³⁸ Although excisional treatments may be more efficacious than ablative treatments in WLWH, especially for those with uncontrolled HIV infections, the cancer prevention benefits of each must be weighed carefully against their potential harms, such as preterm birth and procedural complications, especially in relation to a woman's desire to have children in the future and the ability of the health care system to manage those complications.^{245,246}

Thermal ablation (thermocoagulation) is now recommended as an alternative to cryotherapy²⁴⁷ and should be applied using the same WHO guidelines as cryotherapy.^{238,247} However, there remains a dearth of data on the short-term and long-term effectiveness of any tissue ablative methods to treat high-grade cervical abnormalities and reduce cancer risk in WLWH.^{247,248} A new clinical trial may help address these knowledge gaps.²⁴⁹

Cancer Management

The survival of WLWH with locally advanced cervical cancer (LACC) after treatment in the era of ART differs from patient outcomes that were observed at the onset of the AIDS epidemic. Cervical cancer screening is often part of the basic services provided in HIV screening and treatment

settings, resulting in the earlier detection of cancers. In addition, most patients are on ART at the time of their cervical cancer presentation and have high CD4 counts.²⁵⁰ Many have been viremically HIV-suppressed on ART for several years before cancer diagnosis compared with older cohorts before increased HIV care capacity existed.²⁵¹ Also, for those who are not on ART at the time of their cervical cancer diagnosis, as part of treatment, they are all started on ART, which has led to historical differences in treatment completion, tolerability, and survival compared with historical cohorts. In prospectively followed patients with HIV-associated LACC who were treated on protocol in Zimbabwe and South Africa, concomitant chemoradiotherapy with cisplatin, at the same doses used for HIV-negative women with LACC, was well tolerated with excellent survival.²⁵² Furthermore, those WLWH who had LACC with the best HIV viremic control tolerated treatment better and had the best survival outcomes.^{253,254}

Radiation therapy is the primary treatment for LACC, regardless of HIV status. However, because of the lack of radiation equipment, trained personnel, as well as cost differences or delays because of limited services in public-sector hospitals in LMICs, more aggressive surgical approaches have been considered. This, however, requires experienced gynecologic surgeons with expertise in radical hysterectomies and the complications that arise from these complex surgeries—personnel who are in limited supply and high demand in LMICs. Mostly retrospective investigations of the use of neoadjuvant chemotherapy to shrink the tumor to make the completion of surgery possible have been done and have shown mixed results, with no studies exclusively focusing on WLWH.²⁵⁵⁻²⁵⁸ A 2003 meta-analysis found that studies with chemotherapy cycle lengths ≤ 14 days or cisplatin dose intensities ≥ 25 mg/m² per week showed a slight improvement in survival with neoadjuvant chemotherapy and radiation versus radiation alone, whereas longer cycle lengths or lower cisplatin dose intensities reduced survival.²⁵⁷ A randomized controlled trial in India found poorer disease-free survival (HR, 1.38) but similar overall survival (HR, 1.03) for neoadjuvant chemotherapy plus surgery compared with concomitant chemoradiation.²⁵⁸ A retrospective review of data from 3 northern Italian hospitals found better disease-free survival (HR, 3.95) and overall survival (HR, 5.33) for neoadjuvant chemotherapy plus surgery compared with concomitant chemoradiation.²⁵⁵ None of those studies focused on WLWH or reported on the effects on WLWH in a subgroup analysis. Ethically, such an approach might present challenges in some regions of the world because additional high-risk features (eg, lymph node positivity, positive margins) might still require the need for radiation therapy.

Immunotherapies and targeted therapies have been shown to improve survival in the recurrent setting.²⁵⁹⁻²⁶²

However, there is limited information on the role of these therapies in the setting of HIV and no information on a survival benefit in HIV. Early phase clinical trials to assess safety are underway.²⁶³ In addition, many immunotherapies and targeted therapies might be cost prohibitive in the regions of the world that need it the most, which calls into question the ethics of studying such agents in women with HIV-associated LACC in LMICs unless some provision is made to make them available and affordable.

A recent meta-analysis of retrospective or prospective cohort studies, most of which were conducted in SSA and published between 2012 and 2018, on the chemoradiation treatment of LACC in WLWH found that: 1) 8 of 13 studies identified reported no significant differences in treatment outcomes, 2) 6 of 8 studies identified reported no significant difference in survival, 3) all 4 studies identified reported no significant differences in treatment response, and 4) all 6 studies identified reported no significant differences in toxicity between WLWH and HIV-negative women.²⁶⁴ Of note, one of the 13 studies included in the meta-analysis found that there was a difference in overall survival if the WLWH group included those not on ART.^{264,265} Another of the studies included in the meta-analysis commented that, with regard to the effect of HIV status on the survival of women with invasive cervical cancer, 27.5% of patients tested positive for HIV and had poorer survival compared with those who were HIV-positive, although the difference was not statistically significant.²⁶⁶ A recent AIDS Malignancy Consortium feasibility study found that a high proportion of WLWH with LACC will complete concomitant chemoradiotherapy with the same cisplatin dose as, and with tolerability comparable to, that in HIV-negative women.²⁵² Two studies have reported that lower CD4 counts^{90,253} among WLWH were associated with poorer survival from cervical cancer. Importantly, although some studies have reported worse outcomes for WLWH compared with HIV-negative women, possibly related to being immunocompromised, none of the studies have reported better outcomes for WLWH versus HIV-negative women.

Taken together, immune status, or at least HIV status, might play a role in survival from LACC. Additional investigations are needed to clarify the role of HIV status in cervical cancer management outcomes and to optimize care.

Barriers to Implementation

Although not specific to cervical cancer prevention in WLWH, there are several barriers to implementing a global cervical cancer prevention and control program in LMICs where most WLWH live. In relation to primary and secondary prevention strategies, because the burden of cervical cancer disproportionately affects the poorest countries and populations, a global procurement strategy with innovative

financing mechanisms is needed to improve the availability and affordability of HPV vaccines as well as HPV tests for cervical cancer screening. In addition, without addressing patient-specific social determinants of health care, even with effective access to primary and secondary prevention like what we see in HICs, disparities will persist. Models for reaching the lowest income countries through entities such as GAVI, the Vaccine Alliance should also spur efforts to reach GAVI-support ineligible LMICs that are also largely unable to afford these interventions because of competing priorities within overstretched health budgets.

Underlying the challenges of implementing and scaling up cervical cancer prevention, treatment, and care services for women with HIV in LMICs, especially in some of the highest burden countries in SSA, are some of the most persistent challenges that have historically affected similar vertical, disease-focused programs. In particular, the lack of availability of a specialized and trained workforce and suboptimal health care infrastructures often limit widespread, equitable, and affordable access to high-quality services such as surgery, oncology, radiation, and pathology.^{267–273} Although there are ongoing efforts to address these gaps,^{274–276} it is likely to be decades before these services are of sufficient quality and quantity to meet the demands of cervical cancer prevention and control in these regions as well as many other health care demands. In the meantime, cervical cancer prevention and control programs will need work within those limitations to make progress in averting cervical cancer-related morbidity and mortality.

Financing for cancer-specific services and infrastructures, especially in the public sector and with other competing disease priorities, is often a challenge for already severely constrained national health care budgets. External financing efforts, including through multilateral or bilateral donor initiatives^{277,278} and philanthropic/nonprofit sector investments,²⁷⁹ have been critical to augment and expand nascent efforts by in-country agencies, although ensuring the sustainability of such approaches over the long term remains an ongoing challenge.²⁸⁰

Rwanda is an illustrative example of both the successes and challenges of cervical cancer prevention and control in LMICs. In 2010, Rwanda entered public-private partnerships with Merck and Qiagen to launch a national cervical cancer prevention program of prophylactic HPV vaccination in preadolescent females and HPV testing-based screening in mid-adult women, respectively. From 2011 to 2013, Merck donated Gardasil to the Rwanda government, which launched arguably the most successful national HPV vaccination program globally, achieving a laudable >90% coverage of 12-year-old girls in its first year.²⁸¹ In contrast, the establishment of a complementary national cervical cancer screening program for mid-adult women remained

unrealized, largely because of costs and the lack of health care infrastructure to provide the necessary follow-up care of screen-positive women.

Summary

HIV is an important risk factor for cervical cancer. HIV exerts its effects by acting as an HPV cofactor to increase the likelihood of viral persistence and perhaps progression and invasion.

Because cervical cancer is only one and, relatively speaking, an infrequent comorbidity of HIV infection, its prevention in itself is not a reason for HIV control. Yet, in high-burden countries, where there are the least resources and capacities to directly address the cervical cancer burden and the prevalence of HIV is often high, good HIV care might be the easiest and most cost-effective first step in cervical cancer prevention. Better HIV care through increased access to ART will reduce cervical cancer risk overall, although there might be a short-term increase in cervical cancer incidence as the result of reducing HIV-related deaths as a competing risk.²⁸²

Therefore, achieving the WHO goal for HIV control of “90-90-90” (ie, 90% diagnosis, 90% treated, and 90% virally suppressed) for HIV care²⁸³ will contribute significantly to the WHO’s goal of eliminating cervical cancer as a public health problem globally.³ This is especially so in SSA, where there is a *perfect storm* of high HIV prevalence and a profound lack of preventive services, a consequence of which is that one-fifth of all cervical cancers are attributable to HIV. That is, approximately 2-fold more HIV-related cervical cancers occur in SSA than of all the cervical cancers that occur in the United States annually.¹⁹ Given the general lack of surgical, oncology, and radiation services,^{268–271} as well as incomplete reporting of cancer cases,²⁸⁴ it is likely that the numbers and fractions of HIV-attributable cervical cancer-related deaths are higher still in SSA.

However, benefits of HIV care go beyond cervical cancer risk reduction and impact every step in the cervical cancer control continuum: 1) improved immune responses to prophylactic HPV vaccination, which might be important for long-term vaccine effectiveness in immunocompromised WLWH populations; 2) more effective secondary prevention through screening, diagnosis, and treatment of cervical precancer and early cancer; and 3) perhaps even tertiary prevention through better outcomes and longer survival.

Most notably, comprehensive HIV care will likely improve the clinical and cost effectiveness of secondary cervical cancer prevention by screening by: 1) reducing the number of screen-positive women who will need follow-up care because of fewer HPV infections, 2) reducing the likelihood of

HPV persistence and progression to high-grade cervical abnormalities and increasing the likelihood of regression of the latter, and 3) reducing the size of cervical abnormalities that need to be treated, making them more amenable to be treated effectively, even in low-resource settings, and less likely to recur. As noted above, positive margins after treatment are a strong predictor of recurrent disease, and larger abnormalities, which are more common in WLWH, especially in those with poorly controlled HIV infection, than in HIV-negative women, are more likely to have positive margins.

Given the call for cervical cancer elimination,³ programs implementing cervical cancer care in areas where HIV is highly endemic should consider screening for HIV and, if positive, linkage to HIV care, which will reduce the risk of cervical cancer. If possible, cervical cancer prevention through prophylactic HPV vaccination of younger WLWH and screening of mid-adult WLWH should be integrated with their HIV care, as has been supported widely in high-burden countries in SSA through initiatives such as PEPFAR.^{277,285} The introduction of cervical cancer screening with VIA into PEPFAR has proven to be effective in increasing screening rates in WLWH.²⁸⁶ Efforts to mitigate HIV-related stigma^{267,287} and better approaches for the prevention of attrition²⁸⁸ may be necessary to optimize integration and better care for both epidemics. A meta-analysis reported widely variable low levels of knowledge, attitude, and practice of cervical cancer screening among WLWH living in SSA,²⁸⁹ which will need to be overcome to implement screening in this population.

Not only does better HIV care influence cervical cancer risk, but the converse may also be true. Several studies have demonstrated that prevalent HPV infection increases the risk of HIV acquisition by approximately 2.0-fold to 2.5-fold,^{290,291} perhaps as a result of the recruitment of T cells, the target cell for HIV infection, in the immune response to HPV infection. Therefore, in addition to the protection against both infections offered by barrier methods such as condoms,²⁹²⁻²⁹⁴ and perhaps one day by topical microbicides, reducing the endemicity of HPV by primary prevention through vaccination and secondary prevention through screening and treatment may even help reduce the spread of HIV.

Several key questions and issues need to be addressed in the prevention and control of cervical cancer in WLWH. First, are prophylactic HPV vaccines efficacious in WLWH, and how do the number of doses and the type of adjuvant vaccine influence the effectiveness and duration of protection against even viral end points (eg, the prevention of incident persistent infections, recommended as an acceptable surrogate end point for vaccine trials instead of incident precancerous lesions¹⁵⁹)? Understanding the protective effect of HPV vaccines in WLWH will help to optimize their use in this highly vulnerable, highest risk population.

Second, what is the best screening protocol, including the optimal management, of HPV positive results for WLWH? Because the majority of WLWH live in LMICs, and specifically SSA, these screening protocols need to work in/be applicable to these settings, which have limited resources and health care infrastructure and services. Relying on pathology for cervical diagnosis to guide clinical management is not practical in many of these LMIC settings. A robust, broadly available, and affordable molecular triage method using the residual cervical specimen or, better yet, the self-collected cervicovaginal specimen has the distinct advantage of not burdening women or the health care system with an extra clinical visit and its associated costs. Artificial intelligence-based image analysis of cervical images is a promising method for the triage of HPV positive results²⁹⁵⁻²⁹⁷ and treatment decisions²⁹⁸ because it could remove the interrater variability inherent to visual methods due to their subjectivity. Such a method, if validated in WLWH, might be combined with self-collection and/or rapid, affordable, point-of-care HPV testing (when it becomes available) to greatly reduce the number of clinical visits and associated costs for a cervical cancer screening algorithm for WLWH.

Finally, given the scale-up of cervical cancer screening activities, which will lead to large increases in screen-detected cervical cancers that will need care, are the current cancer treatment protocols optimized for WLWH, especially those living in LMICs? Currently, there are limited data on the comparative effectiveness of cervical cancer treatment modalities, including the management of LACC, in WLWH compared with HIV-negative women, with some, albeit inconsistent, evidence that WLWH may have poorer outcomes. Moreover, in LMICs where HIV prevalence is highest, there is a limited number of trained cancer surgeons²⁶⁸ to effectively perform the radical surgery in localized tumors. Thus cancer management protocols may need to be adapted to improve outcomes. One possible adaptation is the use of neoadjuvant chemotherapy before surgery to shrink the tumor and allow for better completion of surgery. This is being used widely in many SSA settings (P. Castle, personal observation), especially in areas where radiation therapy is not routinely available, but its effectiveness and best practices need to be established to inform international recommendations.

In conclusion, a significant number of cervical cancers and related deaths (as well as other cancers and related deaths) can be averted through better, more comprehensive HIV care alone. Further reductions in cervical cancer incidence and mortality will be achieved through primary, secondary, and tertiary cervical cancer prevention services, all of which are likely to be more effective because of that better HIV care. That is, “90-90-90” for HIV care will help achieve a more *effective* “90-70-90” for cervical cancer control. ■

References

- Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2021;71:209–249.
- Vaccarella S, Lortet-Tieulent J, Plummer M, Franceschi S, Bray F. Worldwide trends in cervical cancer incidence: impact of screening against changes in disease risk factors. *Eur J Cancer*. 2013;49:3262–3273.
- Das M. WHO launches strategy to accelerate elimination of cervical cancer. *Lancet Oncol*. 2021;22:20–21.
- Simelela PN. WHO global strategy to eliminate cervical cancer as a public health problem: an opportunity to make it a disease of the past. *Int J Gynaecol Obstet*. 2021;152:1–3.
- Kuehn BM. WHO launches global push to eliminate cervical cancer. *JAMA*. 2021;325:213.
- The Joint United Nations Programme on HIV/AIDS (UNAIDS). Global HIV & AIDS Statistics—Fact Sheet. Accessed August 5, 2021. [unaids.org/en/resources/fact-sheet](https://www.unaids.org/en/resources/fact-sheet)
- The Joint United Nations Programme on HIV/AIDS (UNAIDS). Global Fact Sheet. Accessed June 27, 2021. [unaids.org/en/media/unaids/contentassets/documents/epidemiology/2012/gr2012/201212120_FactSheet_Global_en.pdf](https://www.unaids.org/en/media/unaids/contentassets/documents/epidemiology/2012/gr2012/201212120_FactSheet_Global_en.pdf)
- Kharsany AB, Karim QA. HIV infection and AIDS in sub-Saharan Africa: current status, challenges and opportunities. *Open AIDS J*. 2016;10:34–48.
- US Department of State. The United States President's Emergency Plan for AIDS Relief (PEPFAR). Accessed June 27, 2021. [state.gov/pepfar/](https://www.state.gov/pepfar/)
- Lohse N, Obel N. Update of survival for persons with HIV infection in Denmark. *Ann Intern Med*. 2016;165:749–750.
- Nsanzimana S, Remera E, Kanters S, et al. Life expectancy among HIV-positive patients in Rwanda: a retrospective observational cohort study. *Lancet Glob Health*. 2015;3:e169–e177.
- Brooks JT, Kaplan JE, Holmes KK, Benson C, Pau A, Masur H. HIV-associated opportunistic infections—going, going, but not gone: the continued need for prevention and treatment guidelines. *Clin Infect Dis*. 2009;48:609–611.
- Horner MJ, Shiels MS, Pfeiffer RM, Engels EA. Deaths attributable to cancer in the US human immunodeficiency virus population during 2001–2015. *Clin Infect Dis*. 2021;72:e224–e231.
- Shiels MS, Engels EA. Evolving epidemiology of HIV-associated malignancies. *Curr Opin HIV AIDS*. 2017;12:6–11.
- Shiels MS, Islam JY, Rosenberg PS, Hall HI, Jacobson E, Engels EA. Projected cancer incidence rates and burden of incident cancer cases in HIV-infected adults in the United States through 2030. *Ann Intern Med*. 2018;168:866–873.
- Marcus JL, Leyden WA, Alexeeff SE, et al. Comparison of overall and comorbidity-free life expectancy between insured adults with and without HIV infection, 2000–2016. *JAMA Netw Open*. 2020;3:e207954.
- From the Centers for Disease Control and Prevention. 1993 revised classification system for HIV infection and expanded surveillance case definition for AIDS among adolescents and adults. *JAMA*. 1993;269:729–730.
- 1993 revised classification system for HIV infection and expanded surveillance case definition for AIDS among adolescents and adults. *MMWR Recomm Rep*. 1992;41(RR-17):1–19.
- Stelzle D, Tanaka LF, Lee KK, et al. Estimates of the global burden of cervical cancer associated with HIV. *Lancet Glob Health*. 2021;9:e161–e169.
- Kelly HA, Sawadogo B, Chikandiwa A, et al. Epidemiology of high-risk human papillomavirus and cervical lesions in African women living with HIV/AIDS: effect of anti-retroviral therapy. *AIDS*. 2017;31:273–285.
- Minkoff H, Zhong Y, Burk RD, et al. Influence of adherent and effective antiretroviral therapy use on human papillomavirus infection and squamous intraepithelial lesions in human immunodeficiency virus-positive women. *J Infect Dis*. 2010;201:681–690.
- Massad LS, Hessol NA, Darragh TM, et al. Cervical cancer incidence after up to 20 years of observation among women with HIV. *Int J Cancer*. 2017;141:1561–1565.
- Centers for Disease Control and Prevention. Appendix A: AIDS-Defining Conditions. Accessed June 27, 2021. [cdc.gov/mmwr/preview/mmwrhtml/rr5710a2.htm](https://www.cdc.gov/mmwr/preview/mmwrhtml/rr5710a2.htm)
- Grulich AE, van Leeuwen MT, Falster MO, Vajdic CM. Incidence of cancers in people with HIV/AIDS compared with immunosuppressed transplant recipients: a meta-analysis. *Lancet*. 2007;370:59–67.
- World Health Organization (WHO). Estimated number of people living with HIV, 2016 By WHO region. Accessed June 27, 2021. [gamapserver.who.int/](https://gamapserver.who.int/mapLibrary/Files/Maps/HIV_all_2016.png)
- World Health Organization (WHO). HIV Country Intelligence: HIV Country Profiles, 2018. WHO; 2018.
- Kaiser Family Foundation. The Global HIV/AIDS Epidemic. Kaiser Family Foundation; 2019.
- Silverberg MJ, Leyden WA, Chi A, et al. Human immunodeficiency virus (HIV)- and non-HIV-associated immunosuppression and risk of cervical neoplasia. *Obstet Gynecol*. 2018;131:47–55.
- Massad LS, Xie X, D'Souza G, et al. Incidence of cervical precancers among HIV-seropositive women. *Am J Obstet Gynecol*. 2015;212:606.e1–e8.
- Abraham AG, D'Souza G, Jing Y, et al. Invasive cervical cancer risk among HIV-infected women: a North American multicohort collaboration prospective study. *J Acquir Immune Defic Syndr*. 2013;62:405–413.
- Lee JS, Cole SR, Achenbach CJ, et al. Cancer risk in HIV patients with incomplete viral suppression after initiation of antiretroviral therapy. *PLoS One*. 2018;13:e0197665.
- Ortiz AP, Engels EA, Nogueras-Gonzalez GM, et al. Disparities in human papillomavirus-related cancer incidence and survival among human immunodeficiency virus-infected Hispanics living in the United States. *Cancer*. 2018;124:4520–4528.
- Rositch AF, Levinson K, Suneja G, et al. Epidemiology of cervical adenocarcinoma and squamous cell carcinoma among women living with HIV compared to the general population in the United States. *Clin Infect Dis*. Published online June 18, 2021. doi:10.1093/cid/ciab561
- Phaswana-Mafuya N, Peltzer K. Breast and cervical cancer screening prevalence and associated factors among women in the South African general population. *Asian Pac J Cancer Prev*. 2018;19:1465–1470.
- Rohner E, Butikofer L, Schmidlin K, et al. Cervical cancer risk in women living with HIV across four continents: a multicohort study. *Int J Cancer*. 2020;146:601–609.
- Schiffman M, Castle PE, Jeronimo J, Rodriguez AC, Wacholder S. Human papillomavirus and cervical cancer. *Lancet*. 2007;370:890–907.
- Schultze A, Paredes R, Sabin C, et al. The association between detected drug resistance mutations and CD4(+) T-cell decline in HIV-positive individuals

- maintained on a failing treatment regimen. *Antivir Ther.* 2018;23:105-116.
38. Schultze A, Torti C, Cozzi-Lepri A, et al. The effect of primary drug resistance on CD4+ cell decline and the viral load set-point in HIV-positive individuals before the start of antiretroviral therapy. *AIDS.* 2019;33:315-326.
 39. Gravitt PE. Evidence and impact of human papillomavirus latency. *Open Virol J.* 2012;6:198-203.
 40. Hinten F, Hilbrands LB, Meeuwis KAP, et al. Reactivation of latent HPV infections after renal transplantation. *Am J Transplant.* 2017;17:1563-1573.
 41. Meeuwis KAP, Hilbrands LB, Int'Hout J, et al. Cervicovaginal HPV infection in female renal transplant recipients: an observational, self-sampling based, cohort study. *Am J Transplant.* 2015;15:723-733.
 42. McHome B, Linde DS, Manongi R, et al. Incident detection of human papillomavirus—a prospective follow-up study among Tanzanian women with focus on HIV status. *Int J Infect Dis.* 2021;110:165-170.
 43. Ermel A, Tong Y, Tonui P, et al. Longer duration of anti-retroviral therapy is associated with decreased risk of human papillomaviruses detection in Kenyan women living with HIV. *Int J STD AIDS.* Published online July 7, 2021. doi:10.1177/09564624211030766
 44. Liu G, Sharma M, Tan N, Barnabas RV. HIV-positive women have higher risk of human papilloma virus infection, pre-cancerous lesions, and cervical cancer. *AIDS.* 2018;32:795-808.
 45. Strickler HD, Burk RD, Fazzari M, et al. Natural history and possible reactivation of human papillomavirus in human immunodeficiency virus-positive women. *J Natl Cancer Inst.* 2005;97:577-586.
 46. Ma X, Wang Q, Ong JJ, et al. Prevalence of human papillomavirus by geographical regions, sexual orientation and HIV status in China: a systematic review and meta-analysis. *Sex Transm Infect.* 2018;94:434-442.
 47. Camargo M, Del Rio-Ospina L, Soto-De Leon SC, et al. Association of HIV status with infection by multiple HPV types. *Trop Med Int Health.* 2018;23:1259-1268.
 48. Okoye JO, Ofordile CA, Adeleke OK, Okechi O. Prevalence of high-risk HPV genotypes in sub-Saharan Africa based on human immunodeficiency virus status: a 20-year systematic review. *Epidemiol Health.* Published online May 25, 2021. doi:10.4178/epih.e2021039
 49. Adebamowo SN, Olawande O, Famooto A, Dareng EO, Offiong R, Adebamowo CA. Persistent low-risk and high-risk human papillomavirus infections of the uterine cervix in HIV-negative and HIV-positive women. *Front Public Health.* 2017;5:178.
 50. Obiri-Yeboah D, Akakpo PK, Mutocheluh M, et al. Epidemiology of cervical human papillomavirus (HPV) infection and squamous intraepithelial lesions (SIL) among a cohort of HIV-infected and uninfected Ghanaian women. *BMC Cancer.* 2017;17:688.
 51. Hanisch RA, Sow PS, Toure M, et al. Influence of HIV-1 and/or HIV-2 infection and CD4 count on cervical HPV DNA detection in women from Senegal, West Africa. *J Clin Virol.* 2013;58:696-702.
 52. Ermel A, Tonui P, Titus M, et al. A cross-sectional analysis of factors associated with detection of oncogenic human papillomavirus in human immunodeficiency virus-infected and uninfected Kenyan women. *BMC Infect Dis.* 2019;19:352.
 53. Kelly H, Weiss HA, Benavente Y, et al. Association of antiretroviral therapy with high-risk human papillomavirus, cervical intraepithelial neoplasia, and invasive cervical cancer in women living with HIV: a systematic review and meta-analysis. *Lancet HIV.* 2018;5:e45-e58.
 54. Menon S, Rossi R, Zdravetska N, et al. Associations between highly active antiretroviral therapy and the presence of HPV, premalignant and malignant cervical lesions in sub-Saharan Africa, a systematic review: current evidence and directions for future research. *BMJ Open.* 2017;7:e015123.
 55. Thorsteinsson K, Storgaard M, Katzenstein TL, et al. Prevalence and distribution of cervical high-risk human papillomavirus and cytological abnormalities in women living with HIV in Denmark—the SHADE. *BMC Cancer.* 2016;16:866.
 56. De Vuyst H, Mugo NR, Chung MH, et al. Prevalence and determinants of human papillomavirus infection and cervical lesions in HIV-positive women in Kenya. *Br J Cancer.* 2012;107:1624-1630.
 57. Chung MH, McKenzie KP, De Vuyst H, et al. Comparing Papanicolaou smear, visual inspection with acetic acid and human papillomavirus cervical cancer screening methods among HIV-positive women by immune status and antiretroviral therapy. *AIDS.* 2013;27:2909-2919.
 58. Abel S, Najioullah F, Volumenie JL, et al. High prevalence of human papillomavirus infection in HIV-infected women living in French Antilles and French Guiana. *PLoS One.* 2019;14:e0221334.
 59. Zeier MD, Botha MH, Engelbrecht S, et al. Combination antiretroviral therapy reduces the detection risk of cervical human papilloma virus infection in women living with HIV. *AIDS.* 2015;29:59-66.
 60. Delory T, Ngo-Giang-Huong N, Rangdaeng S, et al. Human papillomavirus infection and cervical lesions in HIV infected women on antiretroviral treatment in Thailand. *J Infect.* 2017;74:501-511.
 61. Ortiz AP, Tamayo V, Scorsone A, et al. Prevalence and correlates of cervical HPV infection in a clinic-based sample of HIV-positive Hispanic women. *Papillomavirus Res.* 2017;4:39-44.
 62. Adedimeji A, Ajeh R, Dzudie A, et al. Cervical human papillomavirus DNA detection in women living with HIV and HIV-uninfected women living in Limbe, Cameroon. *J Clin Virol.* 2020;128:104445.
 63. McHome BL, Kjaer SK, Manongi R, et al. HPV types, cervical high-grade lesions and risk factors for oncogenic human papillomavirus infection among 3416 Tanzanian women. *Sex Transm Infect.* 2021;97:56-62.
 64. Teixeira MF, Sabido M, Leturiondo AL, de Oliveira Ferreira C, Torres KL, Benzaken AS. High risk human papillomavirus prevalence and genotype distribution among women infected with HIV in Manaus, Amazonas. *Virol J.* 2018;15:36.
 65. Silva L, Miranda A, Batalha R, Ferreira L, Santos M, Talhari S. High-risk human papillomavirus and cervical lesions among women living with HIV/AIDS in Brazilian Amazon, Brazil. *Braz J Infect Dis.* 2015;19:557-562.
 66. Dames DN, Blackman E, Butler R, et al. High-risk cervical human papillomavirus infections among human immunodeficiency virus-positive women in the Bahamas. *PLoS One.* 2014;9:e85429.
 67. Murenzi G, Kanyabwisha F, Murangwa A, et al. Twelve-year trend in the prevalence of high-risk human papillomavirus infection among Rwandan women living with HIV. *J Infect Dis.* 2020;222:74-81.
 68. Konopnicki D, Manigart Y, Gilles C, et al. High-risk human papillomavirus infection in HIV-positive African women living in Europe. *J Int AIDS Soc.* 2013;16:18023.
 69. Teixeira NC, Araujo AC, Correa CM, et al. Prevalence and risk factors for cervical intraepithelial neoplasia among

- HIV-infected women. *Braz J Infect Dis*. 2012;16:164–169.
70. Denny L, Boa R, Williamson AL, et al. Human papillomavirus infection and cervical disease in human immunodeficiency virus-1-infected women. *Obstet Gynecol*. 2008;111:1380–1387.
 71. Hawes SE, Critchlow CW, Faye Niang MA, et al. Increased risk of high-grade cervical squamous intraepithelial lesions and invasive cervical cancer among African women with human immunodeficiency virus type 1 and 2 infections. *J Infect Dis*. 2003;188:555–563.
 72. Rocha-Brischiliari SC, Gimenes F, de Abreu AL, et al. Risk factors for cervical HPV infection and genotypes distribution in HIV-infected South Brazilian women. *Infect Agent Cancer*. 2014;9:6.
 73. Mane A, Nirmalkar A, Risbud AR, Vermund SH, Mehendale SM, Sahasrabudhe VV. HPV genotype distribution in cervical intraepithelial neoplasia among HIV-infected women in Pune, India. *PLoS One*. 2012;7:e38731.
 74. Garbuglia AR, Piselli P, Lapa D, et al. Frequency and multiplicity of human papillomavirus infection in HIV-1 positive women in Italy. *J Clin Virol*. 2012;54:141–146.
 75. Meyrelles ARI, Siqueira JD, Hofer CB, et al. HIV/HPV co-infection during pregnancy in southeastern Brazil: prevalence, HPV types, cytological abnormalities and risk factors. *Gynecol Oncol*. 2013;128:107–112.
 76. Herrero R, Castle PE, Schiffman M, et al. Epidemiologic profile of type-specific human papillomavirus infection and cervical neoplasia in Guanacaste, Costa Rica. *J Infect Dis*. 2005;191:1796–1807.
 77. Trottier H, Mahmud S, Costa MC, et al. Human papillomavirus infections with multiple types and risk of cervical neoplasia. *Cancer Epidemiol Biomarkers Prev*. 2006;15:1274–1280.
 78. Massad L, Keller M, Xie X, et al. Multitype infections with human papillomavirus: impact of human immunodeficiency virus coinfection. *Sex Transm Dis*. 2016;43:637–641.
 79. Clifford GM, Franceschi S, Keiser O, et al. Immunodeficiency and the risk of cervical intraepithelial neoplasia 2/3 and cervical cancer: a nested case-control study in the Swiss HIV Cohort Study. *Int J Cancer*. 2016;138:1732–1740.
 80. Musa J, Achenbach CJ, Evans CT, et al. HIV status, age at cervical cancer screening and cervical cytology outcomes in an opportunistic screening setting in Nigeria: a 10-year cross-sectional data analysis. *Infect Agent Cancer*. 2019;14:43.
 81. Kapambwe S, Sahasrabudhe VV, Blevins M, et al. Implementation and operational research: age distribution and determinants of invasive cervical cancer in a “screen-and-treat” program integrated with HIV/AIDS care in Zambia. *J Acquir Immune Defic Syndr*. 2015;70:e20–e26.
 82. van Bogaert LJ. Age at diagnosis of pre-invasive and invasive cervical neoplasia in South Africa: HIV-positive versus HIV-negative women. *Int J Gynecol Cancer*. 2011;21:363–366.
 83. Wu ES, Urban RR, Krantz EM, et al. The association between HIV infection and cervical cancer presentation and survival in Uganda. *Gynecol Oncol Rep*. 2020;31:100516.
 84. Tawe L, MacDuffie E, Narasimhamurthy M, et al. Human papillomavirus genotypes in women with invasive cervical cancer with and without human immunodeficiency virus infection in Botswana. *Int J Cancer*. 2020;146:1667–1673.
 85. McCredie MR, Paul C, Sharples KJ, et al. Consequences in women of participating in a study of the natural history of cervical intraepithelial neoplasia 3. *Aust N Z J Obstet Gynaecol*. 2010;50:363–370.
 86. Stoler MH, Schiffman M. Interobserver reproducibility of cervical cytologic and histologic interpretations: realistic estimates from the ASCUS-LSIL Triage Study. *JAMA*. 2001;285:1500–1505.
 87. Guan P, Howell-Jones R, Li N, et al. Human papillomavirus types in 115,789 HPV-positive women: a meta-analysis from cervical infection to cancer. *Int J Cancer*. 2012;131:2349–2359.
 88. Chambuso R, Ramesar R, Kaambo E, et al. Age, absolute CD4 count, and CD4 percentage in relation to HPV infection and the stage of cervical disease in HIV-1-positive women. *Medicine (Baltimore)*. 2020;99:e19273.
 89. Menon MP, Coghill A, Mutyaba IO, et al. Association between HIV infection and cancer stage at presentation at the Uganda Cancer Institute. *J Glob Oncol*. 2018;4:1–9.
 90. Dryden-Peterson S, Bvochora-Nsingo M, Suneja G, et al. HIV infection and survival among women with cervical cancer. *J Clin Oncol*. 2016;34:3749–3757.
 91. Ntekim A, Campbell O, Rothenbacher D. Optimal management of cervical cancer in HIV-positive patients: a systematic review. *Cancer Med*. 2015;4:1381–1393.
 92. Simonds HM, Wright JD, du Toit N, Neugut AI, Jacobson JS. Completion of and early response to chemoradiation among human immunodeficiency virus (HIV)-positive and HIV-negative patients with locally advanced cervical carcinoma in South Africa. *Cancer*. 2012;118:2971–2979.
 93. Castle PE, Burk RD, Massad LS, et al. Epidemiological evidence that common HPV types may be common because of their ability to evade immune surveillance: results from the Women's Interagency HIV Study. *Int J Cancer*. 2020;146:3320–3328.
 94. Massad LS, Xie X, Burk RD, et al. Association of cervical precancer with human papillomavirus types other than 16 among HIV co-infected women. *Am J Obstet Gynecol*. 2016;214:354.e1–e6.
 95. Clifford GM, de Vuyst H, Tenet V, Plummer M, Tully S, Franceschi S. Effect of HIV infection on human papillomavirus types causing invasive cervical cancer in Africa. *J Acquir Immune Defic Syndr*. 2016;73:332–339.
 96. Mpunga T, Chantal Umulisa M, Tenet V, et al. Human papillomavirus genotypes in cervical and other HPV-related anogenital cancer in Rwanda, according to HIV status. *Int J Cancer*. 2020;146:1514–1522.
 97. de Sanjose S, Wheeler CM, Quint WGV, et al. Age-specific occurrence of HPV16- and HPV18-related cervical cancer. *Cancer Epidemiol Biomarkers Prev*. 2013;22:1313–1318.
 98. Perkins RB, Guido RS, Castle PE, et al. 2019 ASCCP risk-based management consensus guidelines for abnormal cervical cancer screening tests and cancer precursors. *J Low Genit Tract Dis*. 2020;24:102–131.
 99. Ghebrey RG, Grover S, Xu MJ, Chuang LT, Simonds H. Cervical cancer control in HIV-infected women: past, present and future. *Gynecol Oncol Rep*. 2017;21:101–108.
 100. Palefsky J. Human papillomavirus-related disease in people with HIV. *Curr Opin HIV AIDS*. 2009;4:52–56.
 101. Dhokotera T, Bohlius J, Spoerri A, et al. The burden of cancers associated with HIV in the South African public health sector, 2004–2014: a record linkage study. *Infect Agent Cancer*. 2019;14:12.
 102. Munoz N, Kjaer SK, Sigurdsson K, et al. Impact of human papillomavirus (HPV)-6/11/16/18 vaccine on all HPV-associated genital diseases in young women. *J Natl Cancer Inst*. 2010;102:325–339.
 103. Lehtinen M, Paavonen J, Wheeler CM, et al. Overall efficacy of HPV-16/18 AS04-adjuvanted vaccine against grade

- 3 or greater cervical intraepithelial neoplasia: 4-year end-of-study analysis of the randomised, double-blind PATRICIA trial. *Lancet Oncol.* 2012;13:89-99.
104. Joura EA, Giuliano AR, Iversen OE, et al. A 9-valent HPV vaccine against infection and intraepithelial neoplasia in women. *N Engl J Med.* 2015;372:711-723.
105. de Sanjose S, Quint WG, Alemany L, et al. Human papillomavirus genotype attribution in invasive cervical cancer: a retrospective cross-sectional worldwide study. *Lancet Oncol.* 2010;11:1048-1056.
106. Human papillomavirus vaccines: WHO position paper, October 2014. *Wkly Epidemiol Rec.* 2014;89:465-491.
107. Centers for Disease Control and Prevention (CDC). FDA licensure of quadrivalent human papillomavirus vaccine (HPV4, Gardasil®) for use in males and guidance from the Advisory Committee on Immunization Practices (ACIP). *MMWR Morb Mortal Wkly Rep.* 2010;59:630-632.
108. Brisson M, Laprise JF. Cost-effectiveness of extending HPV vaccination above age 26 years in the US. Advisory Committee on Immunization Practices, Centers for Disease Control and Prevention; 2019.
109. Kim JJ, Goldie SJ. Health and economic implications of HPV vaccination in the United States. *N Engl J Med.* 2008;359:821-832.
110. Kjaer SK, Sigurdsson K, Iversen OE, et al. A pooled analysis of continued prophylactic efficacy of quadrivalent human papillomavirus (types 6/11/16/18) vaccine against high-grade cervical and external genital lesions. *Cancer Prev Res (Phila).* 2009;2:868-878.
111. World Health Organization (WHO). Immunization, Vaccines and Biologicals: Human Papillomavirus (HPV). Accessed March 27, 2018. who.int/immunization/diseases/hpv/en/
112. Centers for Disease Control and Prevention (CDC). HPV Vaccine. Accessed March 27, 2018. cdc.gov/hpv/parents/vaccine.html
113. Whitworth HS, Gallagher KE, Howard N, et al. Efficacy and immunogenicity of a single dose of human papillomavirus vaccine compared to no vaccination or standard three and two-dose vaccination regimens: a systematic review of evidence from clinical trials. *Vaccine.* 2020;38:1302-1314.
114. Kreimer AR, Sampson JN, Porras C, et al. Evaluation of durability of a single dose of the bivalent HPV vaccine: the CVT Trial. *J Natl Cancer Inst.* 2020;112:1038-1046.
115. Historical data from the National HPV Vaccination Program Register. Accessed August 26, 2021. <https://www.health.gov.au/resources/collections/historical-data-from-the-national-hpv-vaccination-program-register>
116. Donovan B, Franklin N, Guy R, et al. Quadrivalent human papillomavirus vaccination and trends in genital warts in Australia: analysis of national sentinel surveillance data. *Lancet Infect Dis.* 2011;11:39-44.
117. Tabrizi SN, Brotherton JM, Kaldor JM, et al. Assessment of herd immunity and cross-protection after a human papillomavirus vaccination programme in Australia: a repeat cross-sectional study. *Lancet Infect Dis.* 2014;14:958-966.
118. Machalek DA, Garland SM, Brotherton JML, et al. Very low prevalence of vaccine human papillomavirus types among 18- to 35-year old Australian women, nine years following implementation of vaccination. *J Infect Dis.* 2018;217:1590-1600.
119. Gertig DM, Brotherton JM, Budd AC, Drennan K, Chappell G, Saville AM. Impact of a population-based HPV vaccination program on cervical abnormalities: a data linkage study. *BMC Med.* 2013;11:227.
120. Crowe E, Pandeya N, Brotherton JM, et al. Effectiveness of quadrivalent human papillomavirus vaccine for the prevention of cervical abnormalities: case-control study nested within a population based screening programme in Australia. *BMJ.* 2014;348:g1458.
121. Brotherton JM, Gertig DM, May C, Chappell G, Saville M. HPV vaccine impact in Australian women: ready for an HPV-based screening program. *Med J Aust.* 2016;204:184-184.e1.
122. Kavanagh K, Pollock KG, Cuschieri K, et al. Changes in the prevalence of human papillomavirus following a national bivalent human papillomavirus vaccination programme in Scotland: a 7-year cross-sectional study. *Lancet Infect Dis.* 2017;17:1293-1302.
123. Baldur-Felskov B, Dehlendorff C, Munk C, Kjaer SK. Early impact of human papillomavirus vaccination on cervical neoplasia—nationwide follow-up of young Danish women. *J Natl Cancer Inst.* 2014;106:djt460.
124. Markowitz LE, Hariri S, Lin C, et al. Reduction in human papillomavirus (HPV) prevalence among young women following HPV vaccine introduction in the United States, National Health and Nutrition Examination Surveys, 2003-2010. *J Infect Dis.* 2013;208:385-393.
125. Markowitz LE, Liu G, Hariri S, Steinau M, Dunne EF, Unger ER. Prevalence of HPV after introduction of the vaccination program in the United States. *Pediatrics.* 2016;137:e20151968.
126. Oliver SE, Unger ER, Lewis R, et al. Prevalence of human papillomavirus among females after vaccine introduction—National Health and Nutrition Examination Survey, United States, 2003-2014. *J Infect Dis.* 2017;216:594-603.
127. Luostarinen T, Apter D, Dillner J, et al. Vaccination protects against invasive HPV-associated cancers. *Int J Cancer.* 2018;142:2186-2187.
128. Drolet M, Benard E, Perez N, Brisson M, HPV Vaccination Impact Study Group. Population-level impact and herd effects following the introduction of human papillomavirus vaccination programmes: updated systematic review and meta-analysis. *Lancet.* 2019;394:497-509.
129. Lei J, Ploner A, Elfstrom KM, et al. HPV vaccination and the risk of invasive cervical cancer. *N Engl J Med.* 2020;383:1340-1348.
130. Kjaer SK, Dehlendorff C, Belmonte F, Baandrup L. Real-world effectiveness of human papillomavirus vaccination against cervical cancer. *J Natl Cancer Inst.* Published online April 20 2021. doi:10.1093/jnci/djab080
131. Levin MJ, Moscicki AB, Song LY, et al. Safety and immunogenicity of a quadrivalent human papillomavirus (types 6, 11, 16, and 18) vaccine in HIV-infected children 7 to 12 years old. *J Acquir Immune Defic Syndr.* 2010;55:197-204.
132. Levin MJ, Huang S, Moscicki AB, et al. Four-year persistence of type-specific immunity after quadrivalent human papillomavirus vaccination in HIV-infected children: effect of a fourth dose of vaccine. *Vaccine.* 2017;35:1712-1720.
133. Weinberg A, Huang S, Moscicki AB, Saah A, Levin MJ. Persistence of memory B-cell and T-cell responses to the quadrivalent HPV vaccine in HIV-infected children. *AIDS.* 2018;32:851-860.
134. Wilkin T, Lee JY, Lensing SY, et al. Safety and immunogenicity of the quadrivalent human papillomavirus vaccine in HIV-1-infected men. *J Infect Dis.* 2010;202:1246-1253.
135. Giacomet V, Penagini F, Trabattini D, et al. Safety and immunogenicity of a quadrivalent human papillomavirus vaccine in HIV-infected and HIV-negative

- adolescents and young adults. *Vaccine*. 2014;32:5657–5661.
136. Kahn JA, Xu J, Kapogiannis BG, et al. Immunogenicity and safety of the human papillomavirus 6, 11, 16, 18 vaccine in HIV-infected young women. *Clin Infect Dis*. 2013;57:735–744.
137. Kojic EM, Kang M, Cespedes MS, et al. Immunogenicity and safety of the quadrivalent human papillomavirus vaccine in HIV-1-infected women. *Clin Infect Dis*. 2014;59:127–135.
138. Money DM, Moses E, Blitz S, et al. HIV viral suppression results in higher antibody responses in HIV-positive women vaccinated with the quadrivalent human papillomavirus vaccine. *Vaccine*. 2016;34:4799–4806.
139. Folschweiller N, Teixeira J, Joshi S, et al. Immunogenicity and safety of the AS04-HPV-16/18 and HPV-6/11/16/18 human papillomavirus vaccines in asymptomatic young women living with HIV aged 15–25 years: a phase IV randomized comparative study. *EClinicalMedicine*. 2020;23:100353.
140. Toft L, Storgaard M, Muller M, et al. Comparison of the immunogenicity and reactogenicity of Cervarix™ and Gardasil® human papillomavirus vaccines in HIV-infected adults: a randomized, double-blind clinical trial. *J Infect Dis*. 2014;209:1165–1173.
141. Einstein MH, Schiller JT, Viscidi RP, et al. Clinician's guide to human papillomavirus immunology: knowns and unknowns. *Lancet Infect Dis*. 2009;9:347–356.
142. Cranston RD, Cespedes MS, Paczuski P, et al. High baseline anal human papillomavirus and abnormal anal cytology in a phase 3 trial of the quadrivalent human papillomavirus vaccine in human immunodeficiency virus-infected individuals older than 26 years: ACTG 5298. *Sex Transm Dis*. 2018;45:266–271.
143. Wilkin TJ, Chen H, Cespedes MS, et al. A randomized, placebo-controlled trial of the quadrivalent human papillomavirus vaccine in human immunodeficiency virus-infected adults aged 27 years or older: AIDS Clinical Trials Group protocol A5298. *Clin Infect Dis*. 2018;67:1339–1346.
144. Moscicki AB, Karalius B, Tassiopoulos K, et al. Human papillomavirus antibody levels and quadrivalent vaccine clinical effectiveness in perinatally human immunodeficiency virus-infected and exposed, uninfected youth. *Clin Infect Dis*. 2019;69:1183–1191.
145. Firnhaber C, Swarts A, Jezile V, et al. HPV vaccination prior to loop electro-excision procedure does not prevent recurrent cervical high grade squamous intraepithelial lesions in women living with HIV: a randomized, double-blind, placebo-controlled trial. *Clin Infect Dis*. Published online September 25, 2020. doi:10.1093/cid/ciaa1456
146. Gosens KCM, van der Zee RP, van Heukelom MLS, et al. HPV vaccination to prevent recurrence of anal intraepithelial neoplasia in HIV+ MSM. *AIDS*. 2021;35(11):1753–1764. doi:10.1097/qad.0000000000002928.
147. Silverberg MJ, Leyden WA, Lam JO, et al. Effectiveness of 'catch-up' human papillomavirus vaccination to prevent cervical neoplasia in immunosuppressed and non-immunosuppressed women. *Vaccine*. 2020;38:4520–4523.
148. Carlander C, Lagheden C, Eklund C, et al. HPV types in cervical precancer by HIV status and birth region: a population-based register study. *Cancer Epidemiol Biomarkers Prev*. 2020;29:2662–2668.
149. Taku O, Mbulawa ZZA, Phohlo K, Garcia-Jardon M, Businge CB, Williamson AL. Distribution of human papillomavirus (HPV) genotypes in HIV-negative and HIV-positive women with cervical intraepithelial lesions in the Eastern Cape Province, South Africa. *Viruses*. 2021;13:280.
150. Pinheiro M, Gage JC, Clifford GM, et al. Association of HPV35 with cervical carcinogenesis among women of African ancestry: evidence of viral-host interaction with implications for disease intervention. *Int J Cancer*. 2020;147:2677–2686.
151. Tsang SH, Sampson JN, Schussler J, et al. Durability of cross-protection by different schedules of the bivalent HPV vaccine: the CVT Trial. *J Natl Cancer Inst*. 2020;112:1030–1037.
152. Woestenbergh PJ, King AJ, van Benthem BHB, et al. Bivalent vaccine effectiveness against type-specific HPV positivity: evidence for cross-protection against oncogenic types among Dutch STI clinic visitors. *J Infect Dis*. 2018;217:213–222.
153. Lehtinen M, Luostarinen T, Vanska S, et al. Gender-neutral vaccination provides improved control of human papillomavirus types 18/31/33/35 through herd immunity: results of a community randomized trial (III). *Int J Cancer*. 2018;143:2299–2310.
154. Faust H, Toft L, Sehr P, et al. Human papillomavirus neutralizing and cross-reactive antibodies induced in HIV-positive subjects after vaccination with quadrivalent and bivalent HPV vaccines. *Vaccine*. 2016;34:1559–1565.
155. Meites E, Kempe A, Markowitz LE. Use of a 2-dose schedule for human papillomavirus vaccination—updated recommendations of the Advisory Committee on Immunization Practices. *MMWR Morb Mortal Wkly Rep*. 2016;65:1405–1408.
156. Arrossi S, Temin S, Garland S, et al. Primary prevention of cervical cancer: American Society of Clinical Oncology resource-stratified guideline. *J Glob Oncol*. 2017;3:611–634.
157. Castle PE, Rodriguez AC, Burk RD, et al. Short term persistence of human papillomavirus and risk of cervical precancer and cancer: population based cohort study. *BMJ*. 2009;339:b2569.
158. Kjaer SK, Frederiksen K, Munk C, Iftner T. Long-term absolute risk of cervical intraepithelial neoplasia grade 3 or worse following human papillomavirus infection: role of persistence. *J Natl Cancer Inst*. 2010;102:1478–1488.
159. Lowy DR, Herrero R, Hildesheim A. Primary endpoints for future prophylactic human papillomavirus vaccine trials: towards infection and immunobridging. *Lancet Oncol*. 2015;16:e226–233.
160. XINHUANET.com. GSK, INNOVAX and Xiamen University to jointly develop next-generation cervical cancer vaccine: Xinhua, 2019. Accessed June 27, 2021. xinhuanet.com/english/2019--09/07/c_138371938.htm
161. World Health Organization (WHO). WHO Guideline for Screening and Treatment of Cervical Pre-Cancer Lesions for Cervical Cancer Prevention. 2nd ed. WHO; 2021. Accessed July 1, 2021. int/publications/i/item/9789240030824
162. Castle PE, Jeronimo J, Temin S, Shastri SS. Screening to prevent invasive cervical cancer: ASCO resource-stratified clinical practice guideline. *J Clin Oncol*. 2017;35:1250–1252.
163. Castle PE, Stoler MH, Wright TC Jr, Sharma A, Wright TL, Behrens CM. Performance of carcinogenic human papillomavirus (HPV) testing and HPV16 or HPV18 genotyping for cervical cancer screening of women aged 25 years and older: a subanalysis of the ATHENA study. *Lancet Oncol*. 2011;12:880–890.
164. Sankaranarayanan R, Nene BM, Shastri SS, et al. HPV screening for cervical cancer in rural India. *N Engl J Med*. 2009;360:1385–1394.
165. Ronco G, Dillner J, Elfstrom KM, et al. Efficacy of HPV-based screening for prevention of invasive cervical

- cancer: follow-up of four European randomised controlled trials. *Lancet*. 2014;383:524-532.
166. Katki HA, Kinney WK, Fetterman B, et al. Cervical cancer risk for women undergoing concurrent testing for human papillomavirus and cervical cytology: a population-based study in routine clinical practice. *Lancet Oncol*. 2011;12:663-672.
 167. Habbema D, De Kok IMCM, Brown ML. Cervical cancer screening in the United States and the Netherlands: a tale of two countries. *Milbank Q*. 2012;90:5-37.
 168. Kitchener HC, Castle PE, Cox JT. Chapter 7: Achievements and limitations of cervical cytology screening. *Vaccine*. 2006;24(suppl 3):S3/63-70.
 169. International Agency for Research on Cancer (IARC), World Health Organization. Cervix Cancer Screening. IARC Handbooks of Cancer Prevention. Vol 10. IARC Press; 2005.
 170. Curry SJ, Krist AH, Owens DK, et al. Screening for cervical cancer: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;320:674-686.
 171. Fontham ETH, Wolf AMD, Church TR, et al. Cervical cancer screening for individuals at average risk: 2020 guideline update from the American Cancer Society. *CA Cancer J Clin*. 2020;70:321-346.
 172. Marcus JZ, Cason P, Downs LS Jr, Einstein MH, Flowers L. The ASCCP Cervical Cancer Screening Task Force endorsement and opinion on the American Cancer Society updated cervical cancer screening guidelines. *J Low Genit Tract Dis*. 2021;25:187-191.
 173. The American College of Obstetrics and Gynecology (ACOG). Updated Cervical Cancer Screening Guidelines. Practice Advisory. ACOG; 2021. Accessed July 1, 2021. [acog.org/clinical/clinical-guidance/practice-advisory/articles/2021/04/updated-cervical-cancer-screening-guidelines](https://www.acog.org/clinical/clinical-guidance/practice-advisory/articles/2021/04/updated-cervical-cancer-screening-guidelines)
 174. Society for Gynecologic Oncology (SGO). SGO endorses ACOG new practice advisory of USPSTF cervical cancer screening recommendations. SGO; 2021. Accessed July 1, 2021. [sgo.org/news/sgo-endorses-acog-new-practice-advisory-of-uspstf-cervical-cancer-screening-recommendations/](https://www.sgo.org/news/sgo-endorses-acog-new-practice-advisory-of-uspstf-cervical-cancer-screening-recommendations/)
 175. Maver PJ, Poljak M. Primary HPV-based cervical cancer screening in Europe: implementation status, challenges, and future plans. *Clin Microbiol Infect*. 2020;26:579-583.
 176. World Health Organization (WHO). WHO Guidelines for Screening and Treatment of Precancerous Lesions for Cervical Cancer Prevention. WHO; 2013.
 177. Jeronimo J, Castle PE, Temin S, et al. Secondary prevention of cervical cancer: ASCO resource-stratified clinical practice guideline. *J Glob Oncol*. 2017;3: 635-657.
 178. Arbyn M, Smith SB, Temin S, Sultana F, Castle P. Detecting cervical precancer and reaching underscreened women by using HPV testing on self samples: updated meta-analyses. *BMJ*. 2018;363:k4823.
 179. World Health Organization (WHO). WHO Prequalification of In Vitro Diagnostics. PUBLIC REPORT. Product: Xpert® HPV. WHO reference number: PQDx 0268-070-00. Accessed August 6, 2021. [who.int/diagnostics_laboratory/evaluations/pq-list/hiv-vrl/171221_final_pq_report_pqdx_0268_070_00.pdf?ua=1](https://www.who.int/diagnostics_laboratory/evaluations/pq-list/hiv-vrl/171221_final_pq_report_pqdx_0268_070_00.pdf?ua=1)
 180. FIND. GeneXpert® negotiated prices. Accessed August 6, 2021. finddx.org/pricing/genexpert/
 181. Castle PE, Ajeh R, Dzudie A, et al. A comparison of screening tests for detection of high-grade cervical abnormalities in women living with HIV from Cameroon. *Infect Agent Cancer*. 2020;15:45.
 182. Dartell MA, Rasch V, Iftner T, et al. Performance of visual inspection with acetic acid and human papillomavirus testing for detection of high-grade cervical lesions in HIV positive and HIV negative Tanzanian women. *Int J Cancer*. 2014;135:896-904.
 183. Firnhaber C, Mayisela N, Mao L, et al. Validation of cervical cancer screening methods in HIV positive women from Johannesburg South Africa. *PLoS One*. 2013;8:e53494.
 184. Joshi S, Sankaranarayanan R, Muwonge R, Kulkarni V, Somanathan T, Divate U. Screening of cervical neoplasia in HIV-infected women in India. *AIDS*. 2013;27:607-615.
 185. Bansil P, Lim J, Byamugisha J, Kumakech E, Nakisige C, Jeronimo JA. Performance of cervical cancer screening techniques in HIV-infected women in Uganda. *J Low Genit Tract Dis*. 2015;19:215-219.
 186. Chibwesha CJ, Frett B, Katundu K, et al. Clinical performance validation of 4 point-of-care cervical cancer screening tests in HIV-infected women in Zambia. *J Low Genit Tract Dis*. 2016;20: 218-223.
 187. Kelly HA, Chikandiwa A, Sawadogo B, et al. Diagnostic accuracy of cervical cancer screening and screening-triage strategies among women living with HIV-1 in Burkina Faso and South Africa: a cohort study. *PLoS Med*. 2021;18:e1003528.
 188. Castle PE, Fetterman B, Poitras N, Lorey T, Kinney W. Safety against cervical precancer and cancer following negative human papillomavirus and Papanicolaou test results in human immunodeficiency virus-infected women. *Arch Intern Med*. 2012;172:1041-1043.
 189. Keller MJ, Burk RD, Xie X, et al. Risk of cervical precancer and cancer among HIV-infected women with normal cervical cytology and no evidence of oncogenic HPV infection. *JAMA*. 2012;308:362-369.
 190. Clarke MA, Cheung LC, Lorey T, et al. 5-Year prospective evaluation of cytology, human papillomavirus testing, and biomarkers for detection of anal precancer in human immunodeficiency virus-positive men who have sex with men. *Clin Infect Dis*. 2019;69:631-638.
 191. Giorgi-Rossi P, Franceschi S, Ronco G. HPV prevalence and accuracy of HPV testing to detect high-grade cervical intraepithelial neoplasia. *Int J Cancer*. 2012;130:1387-1394.
 192. Duan R, Zhao X, Zhang H, et al. Performance of cervical cancer screening and triage strategies among women living with HIV in China. *Cancer Med*. Published online August 2, 2021. doi:10.1002/cam4.4152
 193. Parham GP, Sahasrabudhe VV, Mwanahamuntu MH, et al. Prevalence and predictors of squamous intraepithelial lesions of the cervix in HIV-infected women in Lusaka, Zambia. *Gynecol Oncol*. 2006;103:1017-1022.
 194. Namale G, Mayanja Y, Kamacooko O, et al. Visual inspection with acetic acid (VIA) positivity among female sex workers: a cross-sectional study highlighting one-year experiences in early detection of pre-cancerous and cancerous cervical lesions in Kampala, Uganda. *Infect Agent Cancer*. 2021;16:31.
 195. Henke A, Kluge U, Borde T, McHome B, Serventi F, Henke O. Tanzanian women's knowledge about cervical cancer and HPV and their prevalence of positive VIA cervical screening results. Data from a prevention and awareness campaign in Northern Tanzania, 2017-2019. *Glob Health Action*. 2021;14:1852780.
 196. Thay S, Goldstein A, Goldstein LS, Govind V, Lim K, Seang C. Prospective cohort study examining cervical cancer screening methods in HIV-positive and HIV-negative Cambodian women: a comparison of human papillomavirus testing, visualization with acetic acid and digital colposcopy. *BMJ Open*. 2019;9:e026887.

197. Baldur-Felskov B, Mwaiselage J, Faber MT, et al. Factors associated with a cervical high-grade lesion on cytology or a positive visual inspection with acetic acid among more than 3300 Tanzanian women. *Trop Med Int Health*. 2019;24:229-237.
198. Jolly PE, Mthethwa-Hleta S, Padilla LA, et al. Screening, prevalence, and risk factors for cervical lesions among HIV positive and HIV negative women in Swaziland. *BMC Public Health*. 2017;17:218.
199. Anderson J, Wysong M, Estep D, et al. Evaluation of cervical cancer screening programs in Cote d'Ivoire, Guyana, and Tanzania: effect of HIV status. *PLoS One*. 2015;10:e0139242.
200. Mwanahamuntu MH, Sahasrabudde VV, Blevins M, et al. Utilization of cervical cancer screening services and trends in screening positivity rates in a 'screen-and-treat' program integrated with HIV/AIDS care in Zambia. *PLoS One*. 2013;8:e74607.
201. Huchko MJ, Sneden J, Zakaras JM, et al. A randomized trial comparing the diagnostic accuracy of visual inspection with acetic acid to visual inspection with Lugol's iodine for cervical cancer screening in HIV-infected women. *PLoS One*. 2015;10:e0118568.
202. Chirenje ZM, Loeb L, Mwale M, Nyamapfeni P, Kamba M, Padian N. Association of cervical SIL and HIV-1 infection among Zimbabwean women in an HIV/STI prevention study. *Int J STD AIDS*. 2002;13:765-768.
203. Sansone M, Saccone G, Migliucci A, et al. Screening for cervical carcinoma in HIV-infected women: analysis of main risk factors for cervical cytologic abnormalities. *J Obstet Gynaecol Res*. 2017;43:352-357.
204. Aho I, Kivela P, Haukka J, Sutinen J, Heikinheimo O. Declining prevalence of cytological squamous intraepithelial lesions of the cervix among women living with well controlled HIV—most women living with HIV do not need annual Pap smear screening. *Acta Obstet Gynecol Scand*. 2017;96:1330-1337.
205. Massad LS, Riester KA, Anastos KM, et al. Prevalence and predictors of squamous cell abnormalities in Papanicolaou smears from women infected with HIV-1. Women's Interagency HIV Study Group. *J Acquir Immune Defic Syndr*. 1999;21:33-41.
206. Delmas MC, Larsen C, van Benthem B, et al. Cervical squamous intraepithelial lesions in HIV-infected women: prevalence, incidence and regression. European Study Group on Natural History of HIV Infection in Women. *AIDS*. 2000;14:1775-1784.
207. Duerr A, Kieke B, Warren D, et al. Human papillomavirus-associated cervical cytologic abnormalities among women with or at risk of infection with human immunodeficiency virus. *Am J Obstet Gynecol*. 2001;184:584-590.
208. Cardillo M, Hagan R, Abadi J, Abadi MA. CD4 T-cell count, viral load, and squamous intraepithelial lesions in women infected with the human immunodeficiency virus. *Cancer*. 2001;93:111-114.
209. Orang'o EO, Were E, Rode O, et al. Novel concepts in cervical cancer screening: a comparison of VIA, HPV DNA test and p16(INK4a)/Ki-67 dual stain cytology in Western Kenya. *Infect Agent Cancer*. 2020;15:57.
210. Luckett R, Mogowa N, Li HJ, et al. Performance of two-stage cervical cancer screening with primary high-risk human papillomavirus testing in women living with human immunodeficiency virus. *Obstet Gynecol*. 2019;134:840-849.
211. Strickler HD, Keller MJ, Hessol NA, et al. Primary HPV and molecular cervical cancer screening in US women living with human immunodeficiency virus. *Clin Infect Dis*. 2021;72:1529-1537.
212. Johnson LG, Saidu R, Mbulawa Z, et al. Selecting human papillomavirus genotypes to optimize the performance of screening tests among South African women. *Cancer Med*. 2020;9:6813-6824.
213. Kuhn L, Saidu R, Boa R, et al. Clinical evaluation of modifications to a human papillomavirus assay to optimise its utility for cervical cancer screening in low-resource settings: a diagnostic accuracy study. *Lancet Glob Health*. 2020;8:e296-e304.
214. De Vuyst H, Franceschi S, Plummer M, et al. Methylation levels of CADM1, MAL, and MIR124-2 in cervical scrapes for triage of HIV-infected, high-risk HPV-positive women in Kenya. *J Acquir Immune Defic Syndr*. 2015;70:311-318.
215. Kremer WW, van Zummeren M, Breytenbach E, et al. The use of molecular markers for cervical screening of women living with HIV in South Africa. *AIDS*. 2019;33:2035-2042.
216. Gradissimo A, Lam J, Attonito JD, et al. Methylation of high-risk human papillomavirus genomes are associated with cervical precancer in HIV-positive women. *Cancer Epidemiol Biomarkers Prev*. 2018;27:1407-1415.
217. Murenzi G, Dusingize JC, Rurangwa T, et al. Protocol for the study of cervical cancer screening technologies in HIV-infected women living in Rwanda. *BMJ Open*. 2018;8:e020432.
218. Wentzensen N, Walker JL, Gold MA, et al. Multiple biopsies and detection of cervical cancer precursors at colposcopy. *J Clin Oncol*. 2015;33:83-89.
219. Pretorius RG, Zhang WH, Belinson JL, et al. Colposcopically directed biopsy, random cervical biopsy, and endocervical curettage in the diagnosis of cervical intraepithelial neoplasia II or worse. *Am J Obstet Gynecol*. 2004;191:430-434.
220. Debeaudrap P, Sobngwi J, Tebeu PM, Clifford GM. Residual or recurrent precancerous lesions after treatment of cervical lesions in human immunodeficiency virus-infected women: a systematic review and meta-analysis of treatment failure. *Clin Infect Dis*. 2019;69:1555-1565.
221. Arbyn M, Redman CWE, Verdoodt F, et al. Incomplete excision of cervical precancer as a predictor of treatment failure: a systematic review and meta-analysis. *Lancet Oncol*. 2017;18:1665-1679.
222. Reimers LL, Sotardi S, Daniel D, et al. Outcomes after an excisional procedure for cervical intraepithelial neoplasia in HIV-infected women. *Gynecol Oncol*. 2010;119:92-97.
223. Lima MI, Tafuri A, Araujo AC, de Miranda Lima L, Melo VH. Cervical intraepithelial neoplasia recurrence after conization in HIV-positive and HIV-negative women. *Int J Gynaecol Obstet*. 2009;104:100-104.
224. Cejtin HE, Malapati R, Chaparala S. A comparison of loop electrosurgical excision procedures between human immunodeficiency virus-seropositive and -seronegative women. *J Low Genit Tract Dis*. 2011;15:37-41.
225. van Bogaert LJ. Histometry of high-grade cervical intraepithelial neoplasia and LEEP excision margin in HIV-infected and uninfected women. *Anal Quant Cytopathol Histopathol*. 2014;36:183-188.
226. Cejtin HE, Zimmerman L, Mathews M, Patel A. Predictors of persistent or recurrent disease after loop electrosurgical excision procedure. *J Low Genit Tract Dis*. 2017;21:59-63.
227. Teixeira da Costa Lodi C, Michelin MA, Miranda Lima MI, et al. Predicting cervical intraepithelial neoplasia recurrence in HIV-infected and -noninfected women by detecting aberrant promoter methylation in the CDH1, TIMP3, and

- MGMT genes. *Arch Gynecol Obstet*. 2018;298:971-979.
228. Russomano F, Paz BR, Camargo MJ, et al. Recurrence of cervical intraepithelial neoplasia in human immunodeficiency virus-infected women treated by means of electrosurgical excision of the transformation zone (LLETZ) in Rio de Janeiro, Brazil. *Sao Paulo Med J*. 2013;131:405-410.
 229. Malapati R, Chaparala S, Cejtin HE. Factors influencing persistence or recurrence of cervical intraepithelial neoplasia after loop electrosurgical excision procedure. *J Low Genit Tract Dis*. 2011;15:177-179.
 230. Lodi CT, Michelin MA, Lima MI, et al. Factors associated with recurrence of cervical intraepithelial neoplasia after conization in HIV-infected and non-infected women. *Arch Gynecol Obstet*. 2011;284:191-197.
 231. Huchko MJ, Leslie H, Maloba M, Zakaras J, Bukusi E, Cohen CR. Outcomes up to 12 months after treatment with loop electrosurgical excision procedure for cervical intraepithelial neoplasia among HIV-infected women. *J Acquir Immune Defic Syndr*. 2015;69:200-205.
 232. Babkina N, Heller DS, Goldsmith LT, Houck KL. Cervical conization for cervical intraepithelial neoplasia (CIN) 2 and 3 in HIV-positive women: a case-control study. *J Low Genit Tract Dis*. 2015;19:110-114.
 233. Berrebi A, Badiou W, Duclusaud A. [Frequency, persistence and recurrence of HPV lesions of the uterine cervix in HIV-seropositive women]. *Gynecol Obstet Fertil*. 2008;36:521-524.
 234. Carlander C, Wagner P, van Beirs A, et al. Suppressive antiretroviral therapy associates with effective treatment of high-grade cervical intraepithelial neoplasia. *AIDS*. 2018;32:1475-1484.
 235. Huchko MJ, Leslie H, Maloba M, Bukusi EA, Cohen CR. Factors associated with recurrence of cervical intraepithelial neoplasia 2+ after treatment among HIV-infected women in Western Kenya. *J Acquir Immune Defic Syndr*. 2014;66:188-192.
 236. Zeier MD, Nachega JB, Van Der Merwe FH, et al. Impact of timing of antiretroviral therapy initiation on survival of cervical squamous intraepithelial lesions: a cohort analysis from South Africa. *Int J STD AIDS*. 2012;23:890-896.
 237. Firnhaber C, Swarts A, Goeieman B, et al. Cryotherapy reduces progression of cervical intraepithelial neoplasia grade 1 in South African HIV-infected women: a randomized, controlled trial. *J Acquir Immune Defic Syndr*. 2017;76:532-538.
 238. Santesso N, Mustafa RA, Schunemann HJ, et al. World Health Organization guidelines for treatment of cervical intraepithelial neoplasia 2-3 and screen-and-treat strategies to prevent cervical cancer. *Int J Gynaecol Obstet*. 2016;132:252-258.
 239. Pfaendler KS, Mwanahamuntu MH, Sahasrabudhe VV, Mudenda V, Stringer JS, Parham GP. Management of cryotherapy-ineligible women in a "screen-and-treat" cervical cancer prevention program targeting HIV-infected women in Zambia: lessons from the field. *Gynecol Oncol*. 2008;110:402-407.
 240. Salcedo MP, Oliveira C, Andrade V, et al. The Capulana study: a prospective evaluation of cervical cancer screening using human papillomavirus testing in Mozambique. *Int J Gynecol Cancer*. 2020;30:1292-1297.
 241. Fruchter RG, Maiman M, Sedlis A, Bartley L, Camilien L, Arrastia CD. Multiple recurrences of cervical intraepithelial neoplasia in women with the human immunodeficiency virus. *Obstet Gynecol*. 1996;87:338-344.
 242. Maiman M, Fruchter RG, Serur E, Levine PA, Arrastia CD, Sedlis A. Recurrent cervical intraepithelial neoplasia in human immunodeficiency virus-seropositive women. *Obstet Gynecol*. 1993;82:170-174.
 243. Tate DR, Anderson RJ. Recrudescence of cervical dysplasia among women who are infected with the human immunodeficiency virus: a case-control analysis. *Am J Obstet Gynecol*. 2002;186:880-882.
 244. Greene SA, De Vuyst H, John-Stewart GC, et al. Effect of cryotherapy vs loop electrosurgical excision procedure on cervical disease recurrence among women with HIV and high-grade cervical lesions in Kenya: a randomized clinical trial. *JAMA*. 2019;322:1570-1579.
 245. Kyrgiou M, Athanasiou A, Paraskeva M, et al. Adverse obstetric outcomes after local treatment for cervical preinvasive and early invasive disease according to cone depth: systematic review and meta-analysis. *BMJ*. 2016;354:i3633.
 246. Santesso N, Mustafa RA, Wiercioch W, et al. Systematic reviews and meta-analyses of benefits and harms of cryotherapy, LEEP, and cold knife conization to treat cervical intraepithelial neoplasia. *Int J Gynaecol Obstet*. 2016;132:266-271.
 247. World Health Organization (WHO). WHO Guidelines for the Use of Thermal Ablation for Cervical Pre-Cancer Lesions. World Health Organization; 2019.
 248. Forhan SE, Godfrey CC, Watts DH, Langley CL. A systematic review of the effects of visual inspection with acetic acid, cryotherapy, and loop electrosurgical excision procedures for cervical dysplasia in HIV-infected women in low- and middle-income countries. *J Acquir Immune Defic Syndr*. 2015;68(suppl 3):S350-S356.
 249. National Institutes of Health, National Cancer Institute. A Randomized Clinical Trial to Assess the Effectiveness of Ablative Treatments for Cervical-Cancer Risk Reduction in HIV+ Women Living in Mozambique. Accessed April 20, 2021. reporter.nih.gov/search/L2-n6mYuhEi_2uh17sfkgb/project-details/9927140
 250. Grover S, Bvochora-Nsingo M, Yeager A, et al. Impact of human immunodeficiency virus infection on survival and acute toxicities from chemoradiation therapy for cervical cancer patients in a limited-resource setting. *Int J Radiat Oncol Biol Phys*. 2018;101:201-210.
 251. Simonds HM, Neugut AI, Jacobson JS. HIV status and acute hematologic toxicity among patients with cervix cancer undergoing radical chemoradiation. *Int J Gynecol Cancer*. 2015;25:884-890.
 252. Einstein MH, Ndlovu N, Lee J, et al. Cisplatin and radiation therapy in HIV-positive women with locally advanced cervical cancer in sub-Saharan Africa: a phase II study of the AIDS malignancy consortium. *Gynecol Oncol*. 2019;153:20-25.
 253. Grover S, Mehta P, Wang Q, et al. Association between CD4 count and chemoradiation therapy outcomes among cervical cancer patients with HIV. *J Acquir Immune Defic Syndr*. 2020;85:201-208.
 254. Grover S, MacDuffie EC, Wang Q, et al. HIV infection is not associated with the initiation of curative treatment in women with cervical cancer in Botswana. *Cancer*. 2019;125:1645-1653.
 255. Sala P, Bogliolo S, Barra F, et al. Neoadjuvant chemotherapy followed by radical surgery versus concurrent chemoradiotherapy in the treatment of locally advanced cervical cancer: a multicenter retrospective analysis. *J Invest Surg*. Published online December 8, 2020. doi:10.1080/08941939.2020.1856239
 256. Plante M. Evolution in fertility-preserving options for early-stage cervical cancer: radical trachelectomy, simple trachelectomy, neoadjuvant chemotherapy. *Int J Gynecol Cancer*. 2013;23:982-989.
 257. Neoadjuvant Chemotherapy for Locally Advanced Cervical Cancer Meta-Analysis

- Collaboration. Neoadjuvant chemotherapy for locally advanced cervical cancer: a systematic review and meta-analysis of individual patient data from 21 randomised trials. *Eur J Cancer*. 2003;39:2470-2486.
258. Gupta S, Maheshwari A, Parab P, et al. Neoadjuvant chemotherapy followed by radical surgery versus concomitant chemotherapy and radiotherapy in patients with stage IB2, IIA, or IIB squamous cervical cancer: a randomized controlled trial. *J Clin Oncol*. 2018;36:1548-1555.
 259. Chung HC, Ros W, Delord JP, et al. Efficacy and safety of pembrolizumab in previously treated advanced cervical cancer: results from the phase II KEYNOTE-158 Study. *J Clin Oncol*. 2019;37:1470-1478.
 260. Coleman RL, Lorusso D, Gennigens C, et al. Efficacy and safety of tisotumab vedotin in previously treated recurrent or metastatic cervical cancer (innovaTV 204/GOG-3023/ENGOT-cx6): a multicentre, open-label, single-arm, phase 2 study. *Lancet Oncol*. 2021;22:609-619.
 261. Friedman CF, Snyder Charen A, Zhou Q, et al. Phase II study of atezolizumab in combination with bevacizumab in patients with advanced cervical cancer. *J Immunother Cancer*. 2020;8:3001126.
 262. Mayadev J, Nunes AT, Li M, Marcovitz M, Lanasa MC, Monk BJ. CALLA: efficacy and safety of concurrent and adjuvant durvalumab with chemoradiotherapy versus chemoradiotherapy alone in women with locally advanced cervical cancer: a phase III, randomized, double-blind, multicenter study. *Int J Gynecol Cancer*. 2020;30:1065-1070.
 263. Rajdev L, Cioa EY, Lensing S, et al. AMC 095 (AIDS Malignancy Consortium): a phase I study of ipilimumab (IPI) and nivolumab (NIVO) in advanced HIV associated solid tumors (ST) with expansion cohorts in HIV associated solid tumors and classical Hodgkin lymphoma (cHL) [abstract]. *J Clin Oncol*. 2018;36(15 suppl):TPS2597.
 264. Shah S, Xu M, Mehta P, Zetola NM, Grover S. Differences in outcomes of chemoradiation in women with invasive cervical cancer by human immunodeficiency virus status: a systematic review. *Pract Radiat Oncol*. 2021;11:53-65.
 265. Simonds HM, Botha MH, Neugut AI, Van Der Merwe FH, Jacobson JS. Five-year overall survival following chemoradiation among HIV-positive and HIV-negative patients with locally advanced cervical carcinoma in a South African cohort. *Gynecol Oncol*. 2018;151:215-220.
 266. Maranga IO, Hampson L, Oliver AW, et al. Analysis of factors contributing to the low survival of cervical cancer patients undergoing radiotherapy in Kenya. *PLoS One*. 2013;8:e78411.
 267. Adedimeji A, Ajeh R, Pierz A, et al. Challenges and opportunities associated with cervical cancer screening programs in a low income, high HIV prevalence context. *BMC Womens Health*. 2021;21:74.
 268. Sullivan R, Alatisse OI, Anderson BO, et al. Global cancer surgery: delivering safe, affordable, and timely cancer surgery. *Lancet Oncol*. 2015;16:1193-1224.
 269. Stefan DC. Cancer care in Africa: an overview of resources. *J Glob Oncol*. 2015;1:30-36.
 270. Balogun O, Rodin D, Ngwa W, Grover S, Longo J. Challenges and prospects for providing radiation oncology services in Africa. *Semin Radiat Oncol*. 2017;27:184-188.
 271. Kingham TP, Alatisse OI, Vanderpuye V, et al. Treatment of cancer in sub-Saharan Africa. *Lancet Oncol*. 2013;14:e158-e167.
 272. Adesina A, Chumba D, Nelson AM, et al. Improvement of pathology in sub-Saharan Africa. *Lancet Oncol*. 2013;14:e152-e157.
 273. Nelson AM, Milner DA, Rebeck TR, Iliyasu Y. Oncologic care and pathology resources in Africa: survey and recommendations. *J Clin Oncol*. 2016;34:20-26.
 274. Arneson WL. Essential upgrades: the ASCP lends a hand in sub-Saharan Africa. *MLO Med Lab Obs*. 2015;47:28.
 275. Moretti-Marques R, Salcedo MP, Callegaro Filho D, et al. Telementoring in gynecologic oncology training: changing lives in Mozambique. *Int J Gynecol Cancer*. 2020;30:150-151.
 276. Randall T, Chuang L, Ng SYJ, Schmeler KM, Quinn MA. The IGCS global curriculum mentorship and training program: building human capacity for gynecologic cancer treatment and research where the need is greatest. *Int J Gynecol Cancer*. Published online March 16, 2021. doi:10.1136/ijgc-2021-002480
 277. Godfrey C, Prainito A, Lapidus-Salaiz I, Barnhart M, Watts DH. Reducing cervical cancer deaths in women living with HIV: PEPFAR and the Go Further Partnership. *Prev Med*. 2021;144:106295.
 278. Wagner CM, Antillon F, Uwinkindi F, et al. Establishing cancer treatment programs in resource-limited settings: lessons learned from Guatemala, Rwanda, and Vietnam. *J Glob Oncol*. 2018;4:1-14.
 279. Jailobaeva K, Falconer J, Loffreda G, Arakelyan S, Witter S, Ager A. An analysis of policy and funding priorities of global actors regarding noncommunicable disease in low- and middle-income countries. *Global Health*. 2021;17:68.
 280. Johnson SE, Tasker R, Mired D, D'Cruz A, Adams C. Expanding access to essential quality services for cancer patients as part of universal health coverage: reflections from the World Cancer Leaders' Summit. *JCO Glob Oncol*. 2020;6:500-503.
 281. Binagwaho A, Wagner CM, Gatera M, Karema C, Nutt CT, Ngabo F. Achieving high coverage in Rwanda's national human papillomavirus vaccination programme. *Bull World Health Organ*. 2012;90:623-628.
 282. Hall MT, Smith MA, Simms KT, Barnabas RV, Canfell K, Murray JM. The past, present and future impact of HIV prevention and control on HPV and cervical disease in Tanzania: a modelling study. *PLoS One*. 2020;15:e0231388.
 283. The Joint United Nations Programme on HIV/AIDS (UNAIDS). UNAIDS 90-90-90: An Ambitious Treatment Target to Help End the AIDS Epidemic. UNAIDS; 2014.
 284. Parkin DM, Ferlay J, Jemal A, et al, eds. Cancer in Sub-Saharan Africa. International Agency for Research on Cancer (IARC) Scientific Publications No. 167. IARC; 2018.
 285. Parham GP, Mwanahamuntu MH, Kapambwe S, et al. Population-level scale-up of cervical cancer prevention services in a low-resource setting: development, implementation, and evaluation of the cervical cancer prevention program in Zambia. *PLoS One*. 2015;10:e0122169.
 286. George W. Bush Presidential Center. Go Further Partnership Reaches More Than One Million Women Living With HIV With First-Time Cervical Cancer Screenings. George W. Bush Presidential Center; 2020.
 287. Nyblade L, Mingkwan P, Stockton MA. Stigma reduction: an essential ingredient to ending AIDS by 2030. *Lancet HIV*. 2021;8:e106-e113.
 288. Rohner E, Mulongo M, Pasipamire T, et al. Mapping the cervical cancer screening cascade among women living with HIV in Johannesburg, South Africa. *Int J Gynaecol Obstet*. 2021;152:53-59.
 289. Bogale AL, Teklehaymanot T, Haidar Ali J, Kassie GM. Knowledge, attitude and practice of cervical cancer screening among women infected with HIV in Africa: systematic review and meta-analysis. *PLoS One*. 2021;16:e0249960.

290. Houlihan CF, Larke NL, Watson-Jones D, et al. Human papillomavirus infection and increased risk of HIV acquisition. A systematic review and meta-analysis. *AIDS*. 2012;26:2211-2222.
291. Liu G, Mugo NR, Brown ER, et al. Prevalent HPV infection increases the risk of HIV acquisition in African women: advancing the argument for HPV immunization. *AIDS*. Published online June 24, 2021. doi:10.1097/QAD.0000000000003004
292. Winer RL, Hughes JP, Feng Q, et al. Condom use and the risk of genital human papillomavirus infection in young women. *N Engl J Med*. 2006;354:2645-2654.
293. Giannou FK, Tsiara CG, Nikolopoulos GK, et al. Condom effectiveness in reducing heterosexual HIV transmission: a systematic review and meta-analysis of studies on HIV serodiscordant couples. *Expert Rev Pharmacoecon Outcomes Res*. 2016;16:489-499.
294. Pinkerton SD, Abramson PR. Effectiveness of condoms in preventing HIV transmission. *Soc Sci Med*. 1997;44:1303-1312.
295. Desai KT, Ajenifuja KO, Banjo A, et al. Design and feasibility of a novel program of cervical screening in Nigeria: self-sampled HPV testing paired with visual triage. *Infect Agent Cancer*. 2020;15:60.
296. Xue Z, Novetsky AP, Einstein MH, et al. A demonstration of automated visual evaluation of cervical images taken with a smartphone camera. *Int J Cancer*. 2020;147:2416-2423.
297. Hu L, Bell D, Antani S, et al. An observational study of deep learning and automated evaluation of cervical images for cancer screening. *J Natl Cancer Inst*. 2019;111:923-932.
298. Guo P, Xue Z, Jeronimo J, et al. Network visualization and pyramidal feature comparison for ablative treatability classification using digitized cervix images. *J Clin Med*. 2021;10:953.