

On conformational changes of proteins using collective motions in torsion angles and L_1 regularization

Ryabinina Raisa, Emtsev Daniil

Moscow Institute of Physics and Technology




ryabinina.rb@phystech.edu, daniil.emcev.ru@yandex.ru

December 10, 2018

- 1 Conformational changes and normal modes in torsion angle space
- 2 Overview of regularization methods of obtaining conformational changes
- 3 Results

Conformational changes and normal modes in torsion angle space

Figure: Protein backbone and 1MQ4-10L6 conformational change



Problem

We address the question how well these conformational changes are described using only torsion angles, and how can the change in torsion angles be determined from the observed changes of Cartesian coordinates.

$$\Delta r = J \Delta \phi$$

$$\Delta \phi - ???$$

$$\frac{1}{2n} \|\Delta r - J \Delta \phi\|_2^2 + \alpha \|\Delta \phi\|_1$$

Overview of methods

Ridge regression

$$\min_{\Delta\phi} (\Delta\phi, J^T M J \Delta\phi) - 2(\Delta\phi, J^T M \Delta r) + \lambda(\Delta\phi, \Delta\phi)$$


Least absolute shrinkage and selection operator

$$\min_{\Delta\phi} (\Delta\phi, J^T M J \Delta\phi) - 2(\Delta\phi, J^T M \Delta r) + \lambda \sum_{j=1}^p |\Delta\phi_j|$$

Elastic net regularization

$$\min_{\Delta\phi} (\Delta\phi, J^T M J \Delta\phi) - 2(\Delta\phi, J^T M \Delta r) + \alpha(\Delta\phi, \Delta\phi) + (1 - \alpha) \sum_{j=1}^p |\Delta\phi_j|$$

Figure: Correlation of RMSD in Grid search Lasso and Ridge regression



Results




Figure: Comparison of correlations of RMSD in all methods(using linear approximation)

Thank you for your attention!