

# Chapter 7

## Inference for numerical data<sup>1</sup>

Department of Mathematics & Statistics  
North Carolina A&T State University


---

<sup>1</sup>These notes use content from OpenIntro Statistics Slides by Mine Cetinkaya-Rundel.

One-sample means with the  $t$  distribution

# Heights

- ▶ According to the CDC, the mean height of U.S. adults ages 20 and older is about 66.5 inches (69.3 inches for males, and 63.8 inches for females).
- ▶ In our sample data, we have a sample of 430 college students from a single college.



## Summary statistics

n	$\bar{x}$	s	minimum	maximum
430	67.09	4.86	53.78	83.21

**Objective:** We would like to investigate if the mean height of students at this college is significantly different than 66.5 inches.

## Conditions


- ▶ **Independence:** We are told to assume that cases (rows) are independent.

## Conditions

- ▶ **Independence:** We are told to assume that cases (rows) are independent.
- ▶ **Sample size / skew:**


# Conditions

- ▶ **Independence:** We are told to assume that cases (rows) are independent.
- ▶ **Sample size / skew:**
- ▶ The sample distribution does not appear to be extremely skewed, but it's very difficult to assess with such a small sample size. We might want to think about whether we would expect the population distribution to be skewed or not — probably not, it should be equally likely to have days with lower than average traffic and higher than average traffic.
- ▶ We do not know  $\sigma$  and  $n$  is too small to assume  $s$  is a reliable estimate for  $\sigma$ .



# Conditions

- ▶ **Independence:** We are told to assume that cases (rows) are independent.
- ▶ **Sample size / skew:**
- ▶ The sample distribution does not appear to be extremely skewed, but it's very difficult to assess with such a small sample size. We might want to think about whether we would expect the population distribution to be skewed or not — probably not, it should be equally likely to have days with lower than average traffic and higher than average traffic.
- ▶ We do not know  $\sigma$  and  $n$  is too small to assume  $s$  is a reliable estimate for  $\sigma$ .



So what do we do when the sample size is small?



## Review: what purpose does a large sample serve?

As long as observations are independent, and the population distribution is not extremely skewed, a large sample would ensure that...

- ▶ The sampling distribution of the mean is nearly normal.
- ▶ The estimate of the standard error, as  $\frac{s}{\sqrt{n}}$ , is reliable.

## The normality condition

- ▶ The CLT, which states that sampling distributions will be nearly normal, holds true for **any** sample size as long as the population distribution is nearly normal.

## The normality condition

- ▶ The CLT, which states that sampling distributions will be nearly normal, holds true for **any** sample size as long as the population distribution is nearly normal.
- ▶ While this is helpful special case, it's inherently difficult to verify normality in small data sets.

# The normality condition

- ▶ The CLT, which states that sampling distributions will be nearly normal, holds true for **any** sample size as long as the population distribution is nearly normal.
- ▶ While this is helpful special case, it's inherently difficult to verify normality in small data sets.
- ▶ We should exercise caution when verifying the normality condition for small samples. It is important to not only examine the data but also think about where the data come from.
  - ▶ For example, ask: would I expect this distribution to be symmetric, and am I confident that outliers are rare?

## The $t$ distribution

- ▶ When the population standard deviation is unknown (almost always), the uncertainty of the standard error estimate is addressed by using a new distribution: the  **$t$  distribution**.

## The $t$ distribution


- ▶ When the population standard deviation is unknown (almost always), the uncertainty of the standard error estimate is addressed by using a new distribution: the  **$t$  distribution**.
- ▶ This distribution also has a bell shape, but its tails are **thicker** than the normal model's.

## The $t$ distribution

- ▶ When the population standard deviation is unknown (almost always), the uncertainty of the standard error estimate is addressed by using a new distribution: the  **$t$  distribution**.
- ▶ This distribution also has a bell shape, but its tails are **thicker** than the normal model's.
- ▶ Therefore observations are more likely to fall beyond two SDs from the mean than under the normal distribution.

## The $t$ distribution


- ▶ When the population standard deviation is unknown (almost always), the uncertainty of the standard error estimate is addressed by using a new distribution: the  **$t$  distribution**.
- ▶ This distribution also has a bell shape, but its tails are **thicker** than the normal model's.
- ▶ Therefore observations are more likely to fall beyond two SDs from the mean than under the normal distribution.
- ▶ Extra thick tails are helpful for resolving our problem with a less reliable estimate the standard error (since  $n$  is small).






# The $t$ distribution

- ▶ Always centered at zero, like the standard normal ( $z$ ) distribution.
- ▶ Has a single parameter: **degrees of freedom (df)**.



## The $t$ distribution


- ▶ Always centered at zero, like the standard normal ( $z$ ) distribution.
- ▶ Has a single parameter: **degrees of freedom** ( $df$ ).



What happens to shape of the  $t$  distribution as  $df$  increases?

## The $t$ distribution

- ▶ Always centered at zero, like the standard normal ( $z$ ) distribution.
- ▶ Has a single parameter: **degrees of freedom** ( $df$ ).



What happens to shape of the  $t$  distribution as  $df$  increases?

Approaches normal.

## Back to the student heights survey

n	$\bar{x}$	s	minimum	maximum
430	67.09	4.86	53.78	83.21

**Objective:** We would like to investigate if the mean height of students at this college is significantly different than 66.5 inches.

# Hypotheses

What are the hypotheses for testing for the mean of college student heights being different from 67 inches?

A)  $H_0 : \mu = 66.5$

$H_A : \mu \neq 66.5$

B)  $H_0 : \mu = 66.5$

$H_A : \mu > 66.5$

C)  $H_0 : \mu = 66.5$

$H_A : \mu < 66.5$

D)  $H_0 : \mu \neq 66.5$

$H_A : \mu > 66.5$

# Hypotheses

What are the hypotheses for testing for the mean of college student heights being different from 66.5 inches?

A)  $H_0 : \mu = 66.5$

$H_A : \mu \neq 66.5$

B)  $H_0 : \mu = 66.5$

$H_A : \mu > 66.5$

C)  $H_0 : \mu = 66.5$

$H_A : \mu < 66.5$

D)  $H_0 : \mu \neq 66.5$

$H_A : \mu > 66.5$

## Finding the test statistic

The test statistic for inference on sample mean is the  $T$  statistic with  $df = n - 1$ .

$$T_{df} = \frac{\text{point estimate} - \text{null value}}{SE}$$

## Finding the test statistic

The test statistic for inference on sample mean is the  $T$  statistic with  $df = n - 1$ .

$$T_{df} = \frac{\text{point estimate} - \text{null value}}{SE}$$

in context...

$$\text{point estimate} = \bar{x} = 67.09$$



## Finding the test statistic

The test statistic for inference on sample mean is the  $T$  statistic with  $df = n - 1$ .

$$T_{df} = \frac{\text{point estimate} - \text{null value}}{SE}$$

in context...

$$\text{point estimate} = \bar{x} = 67.09$$

$$SE = \frac{s}{\sqrt{n}} = \frac{4.86}{\sqrt{430}} = 0.234$$

## Finding the test statistic

The test statistic for inference on sample mean is the  $T$  statistic with  $df = n - 1$ .

$$T_{df} = \frac{\text{point estimate} - \text{null value}}{SE}$$

in context...

$$\text{point estimate} = \bar{x} = 67.09$$

$$SE = \frac{s}{\sqrt{n}} = \frac{4.86}{\sqrt{430}} = 0.234$$

$$T = \frac{67.09 - 66.5}{0.234} = 2.52$$

## Finding the test statistic

The test statistic for inference on sample mean is the  $T$  statistic with  $df = n - 1$ .

$$T_{df} = \frac{\text{point estimate} - \text{null value}}{SE}$$

in context...

$$\text{point estimate} = \bar{x} = 67.09$$

$$SE = \frac{s}{\sqrt{n}} = \frac{4.86}{\sqrt{430}} = 0.234$$

$$T = \frac{67.09 - 66.5}{0.234} = 2.52$$

$$df = 430 - 1 = 429$$

---

**Note:** Null value is 66.5 because in the null hypothesis we set  $\mu = 66.5$ .

## Finding the p-value

- ▶ The p-value is, once again, calculated as the area tail area under the  $t$  distribution.

## Finding the p-value

- ▶ The p-value is, once again, calculated as the area tail area under the  $t$  distribution.
- ▶ Using a web app:  
[https://gallery.shinyapps.io/dist\\_calc/](https://gallery.shinyapps.io/dist_calc/)

## Finding the p-value

- ▶ The p-value is, once again, calculated as the area tail area under the  $t$  distribution.
- ▶ Using a web app:  
[https://gallery.shinyapps.io/dist\\_calc/](https://gallery.shinyapps.io/dist_calc/)
- ▶ Or when these aren't available, we can use a  $t$ -table.

## Conclusion of the test

What is the conclusion of this hypothesis test?

## Conclusion of the test

What is the conclusion of this hypothesis test?

We saw that the p-value was extremely low. Thus, we reject the null hypothesis. Based on the p-value, we conclude that the survey provide strong evidence that the mean of the college students height is different from the mean height of U.S. adults over 20.



## What is the difference?

- ▶ We concluded that there is a difference in the mean heights of the college students compared to the mean height of U.S. adults

# What is the difference?

- ▶ We concluded that there is a difference in the mean heights of the college students compared to the mean height of U.S. adults
- ▶ But it would be more interesting to find out what exactly this difference is.

## What is the difference?

- ▶ We concluded that there is a difference in the mean heights of the college students compared to the mean height of U.S. adults
- ▶ But it would be more interesting to find out what exactly this difference is.
- ▶ We can use a confidence interval to estimate this difference.

## Confidence interval for a sample mean

- Confidence intervals are always of the form

$$\text{point estimate} \pm ME$$

## Confidence interval for a sample mean

- ▶ Confidence intervals are always of the form

$$\text{point estimate} \pm ME$$

- ▶ ME is always calculated as the product of a critical value and SE.

## Confidence interval for a sample mean

- ▶ Confidence intervals are always of the form

$$\text{point estimate} \pm ME$$

- ▶ ME is always calculated as the product of a critical value and SE.

- ▶  $ME = t^* \times SE$

$$\text{point estimate} \pm t^* \times SE$$

## Finding the critical $t(t^*)$

- ▶ We want to find the 95% confidence interval.

## Constructing a CI for a small sample mean

Which of the following is the correct calculation of a 95% confidence interval for the heights of the college students?

$$\bar{x} = 67.09 \quad s = 4.86 \quad n = 430 \quad SE = 0.234$$

- A)  $66.5 \pm 1.96 \times 0.234$
- B)  $67.09 \pm 1.97 \times 0.234$
- C)  $67.09 \pm -2.26 \times 0.234$
- D)  $66.5 \pm 2.26 \times 4.86$



## Constructing a CI for a small sample mean

Which of the following is the correct calculation of a 95% confidence interval for the heights of the college students?

$$\bar{x} = 67.09 \quad s = 4.86 \quad n = 430 \quad SE = 0.234$$

- A)  $66.5 \pm 1.96 \times 0.234$
- B)  $67.09 \pm 1.97 \times 0.234 \rightarrow (66.63, 67.55)$
- C)  $67.09 \pm -2.26 \times 0.234$
- D)  $66.5 \pm 2.26 \times 4.86$

# Synthesis

Does the conclusion from the hypothesis test agree with the findings of the confidence interval?

# Synthesis

Does the conclusion from the hypothesis test agree with the findings of the confidence interval?

Yes, the hypothesis test found a significant difference, and the CI does not contain the null value of 66.5.

## Recap: Inference using the $t$ -distribution

- ▶ If  $\sigma$  is unknown, use  $t$ -distribution with  $SE = \frac{s}{\sqrt{n}}$ .

## Recap: Inference using the $t$ -distribution

- ▶ If  $\sigma$  is unknown, use  $t$ -distribution with  $SE = \frac{s}{\sqrt{n}}$ .
- ▶ Conditions:
  - ▶ Independence of observations (often verified by random sample, and if sampling w/o replacement,  $n < 10\%$  of population).
  - ▶ No extreme skew.

## Recap: Inference using the $t$ -distribution

- ▶ If  $\sigma$  is unknown, use  $t$ -distribution with  $SE = \frac{s}{\sqrt{n}}$ .
- ▶ Conditions:
  - ▶ Independence of observations (often verified by random sample, and if sampling w/o replacement,  $n < 10\%$  of population).
  - ▶ No extreme skew.
- ▶ Hypothesis testing:

$$T_{df} = \frac{\text{point estimate} - \text{null value}}{SE}, \text{ where } df = n - 1$$

## Recap: Inference using the $t$ -distribution

- ▶ If  $\sigma$  is unknown, use  $t$ -distribution with  $SE = \frac{s}{\sqrt{n}}$ .
- ▶ Conditions:
  - ▶ Independence of observations (often verified by random sample, and if sampling w/o replacement,  $n < 10\%$  of population).
  - ▶ No extreme skew.
- ▶ Hypothesis testing:

$$T_{df} = \frac{\text{point estimate} - \text{null value}}{SE}, \text{ where } df = n - 1$$

- ▶ Confidence interval:  $\text{point estimate} \pm t_{df}^* \times SE$