

NATIONAL UNIVERSITY OF SINGAPORE  
Department of Mathematics

MA1522 Linear Algebra for Computing

Tutorial 9

1. A father wishes to distribute an amount of money among his three sons Jack, Jim, and John. He wish to distribute such that the following conditions are all satisfied.
  - (i) The amount Jack receives plus twice the amount Jim receives is \$300.
  - (ii) The amount Jim receives plus the amount John receives is \$300.
  - (iii) Jack receives \$300 more than twice of what John receives.

- (a) Is it possible for the following conditions to all be satisfied?
  - (b) If it is not possible, find a least square solution. (Make sure that your least square solution is feasible. For example, one cannot give a negative amount of money to anybody.)

2. (a) Suppose  $\mathbf{A}$  is a  $m \times n$  matrix where  $m > n$ . Let  $\mathbf{A} = \mathbf{Q}\mathbf{R}$  be a QR factorization of  $\mathbf{A}$ . Explain how you might use this to write

$$\mathbf{A} = \mathbf{Q}'\mathbf{R}',$$

where  $\mathbf{Q}'$  is an  $m \times m$  orthogonal matrix, and  $\mathbf{R}'$  a  $m \times n$  matrix with  $m - n$  zero rows at the bottom. This is known as the *full* QR factorization of  $\mathbf{A}$ .

- (b) In MATLAB, enter the following.

```
>> A=sym([1 1 0;1 1 0;1 1 1;0 1 1])
```

```
>> [Q R]=qr(A)
```

What is  $\mathbf{Q}$  and  $\mathbf{R}$ ? Compare this with the answer in tutorial 8 question 5(a).

- (c) Explain how you might use the command `qr` in MATLAB to find a QR factorization of a  $m \times n$  matrix  $\mathbf{A}$ ?
3. (Cayley-Hamilton theorem)  
Consider

$$p(\mathbf{X}) = \mathbf{X}^3 - 4\mathbf{X}^2 - \mathbf{X} + 4\mathbf{I}.$$

- (a) Compute  $p(\mathbf{X})$  for  $\mathbf{X} = \begin{pmatrix} 1 & 1 & 2 \\ 1 & 2 & 1 \\ 2 & 1 & 1 \end{pmatrix}$ .

- (b) Find the characteristic polynomial of  $\mathbf{X}$ .

- (c) Show that  $\mathbf{X}$  invertible. Express the inverse of  $\mathbf{X}$  as a function of  $\mathbf{X}$ .

This question demonstrated the *Cayley-Hamilton theorem*, which states that if  $p(x)$  is the characteristic polynomial of  $\mathbf{X}$ , then  $p(\mathbf{X}) = \mathbf{0}$ . This also show that if 0 is not an eigenvalue of  $\mathbf{X}$ , then the constant term of the characteristic polynomial  $p(x)$  is nonzero, and we can use that to compute the inverse of  $\mathbf{X}$ .

4. For each of the following matrices  $\mathbf{A}$ , determine if  $\mathbf{A}$  is diagonalizable. If  $\mathbf{A}$  is diagonalizable, find an invertible  $\mathbf{P}$  that diagonalizes  $\mathbf{A}$  and determine  $\mathbf{P}^{-1}\mathbf{A}\mathbf{P}$ .

(a)  $\mathbf{A} = \begin{pmatrix} 1 & -3 & 3 \\ 3 & -5 & 3 \\ 6 & -6 & 4 \end{pmatrix}.$

(b)  $\mathbf{A} = \begin{pmatrix} 9 & 8 & 6 & 3 \\ 0 & -1 & 3 & -4 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix}$

(c)  $\mathbf{A} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}.$

(d)  $\mathbf{A} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$

(e)  $\mathbf{A} = \begin{pmatrix} -1 & 1 & 1 \\ 1 & 1 & -1 \\ -4 & 2 & 3 \end{pmatrix}.$

5. (a) Show that  $\lambda$  is an eigenvalue of  $\mathbf{A}$  if and only if it is an eigenvalue of  $\mathbf{A}^T$ .  
 (b) Suppose  $\mathbf{A}$  is diagonalizable. Is  $\mathbf{A}^T$  diagonalizable? Justify your answer.  
 (c) Suppose  $\mathbf{v}$  is an eigenvector of  $\mathbf{A}$  associated to eigenvalue  $\lambda$ . Show that  $\mathbf{v}$  is an eigenvector of  $\mathbf{A}^k$  associated to eigenvalue  $\lambda^k$  for any positive integer  $k$ .  
 (d) If  $\mathbf{A}$  is invertible, show that  $\mathbf{v}$  is an eigenvector of  $\mathbf{A}^k$  associated to eigenvalue  $\lambda^k$  for any negative integer  $k$ .  
 (e) A square matrix is said to be *nilpotent* if there is a positive integer  $k$  such that  $\mathbf{A}^k = \mathbf{0}$ . Show that if  $\mathbf{A}$  is nilpotent, then 0 is the only eigenvalue.  
 (f) Let  $\mathbf{A}$  be a  $n \times n$  matrix with one eigenvalue  $\lambda$  with algebraic multiplicity  $n$ . Show that  $\mathbf{A}$  is diagonalizable if and only if  $\mathbf{A}$  is a scalar matrix,  $\mathbf{A} = \lambda\mathbf{I}$ .  
 (g) Show that the only diagonalizable nilpotent matrix is the zero matrix.

## Extra problems

1. Let  $\mathbf{A}$  be an orthogonal matrix of order  $n$  and let  $\mathbf{u}, \mathbf{v}$  be any two vectors in  $\mathbb{R}^n$ . Show that
  - (a)  $\|\mathbf{u}\| = \|\mathbf{A}\mathbf{u}\|$ ;
  - (b)  $d(\mathbf{u}, \mathbf{v}) = d(\mathbf{A}\mathbf{u}, \mathbf{A}\mathbf{v})$ ; and
  - (c) the angle between  $\mathbf{u}$  and  $\mathbf{v}$  is equal to the angle between  $\mathbf{A}\mathbf{u}$  and  $\mathbf{A}\mathbf{v}$ .
2. Let  $\mathbf{A}$  be an orthogonal matrix of order  $n$ . Let  $S = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$  and define  $T = \{\mathbf{A}\mathbf{u}_1, \mathbf{A}\mathbf{u}_2, \dots, \mathbf{A}\mathbf{u}_k\}$ .
  - (a) If  $S$  is orthogonal, show that  $T$  is orthogonal.
  - (b) If  $S$  is orthonormal, is  $T$  orthonormal?
3.
  - (a) Suppose  $\mathbf{A}$  and  $\mathbf{B}$  are similar matrices, that is,  $\mathbf{A} = \mathbf{P}\mathbf{B}\mathbf{P}^{-1}$  for some invertible matrix  $\mathbf{P}$ . Then the characteristic polynomial of  $\mathbf{A}$  and  $\mathbf{B}$  are equal.
  - (b) Suppose the characteristic polynomial of  $\mathbf{A}$  and  $\mathbf{B}$  are equal. Can we conclude that  $\mathbf{A}$  and  $\mathbf{B}$  are similar? Justify your answer.
  - (c) Suppose  $\mathbf{A}$  and  $\mathbf{B}$  are  $n \times n$  matrices with the same determinant. Is it true that their characteristic polynomials are equal? Justify your answer.
4. (a) Let  $\mathbf{A}$  be a  $2 \times 2$  matrix. Prove that the characteristic polynomial of  $\mathbf{A}$  is

$$x^2 - \text{tr}(\mathbf{A})x + \det(\mathbf{A}),$$

where  $\text{tr}(\mathbf{A})$  is the sum of the diagonal entries of  $\mathbf{A}$ .

- (b) Let  $\mathbf{A}$  be a  $n \times n$  matrix and  $p(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$ . Prove that  $a_0 = (-1)^n \det(\mathbf{A})$ .