

Recursive Rows in Rome

AH & JGM

ACM Reference Format:

AH & JGM. 2023. Recursive Rows in Rome. 1, 1 (December 2023), 6 pages. <https://doi.org/10.1145/nnnnnnnn>.
nnnnnnnn

1 IX: THE INDEX CALCULUS

1.1 Syntax

Sorts	$\sigma ::= \star \mid \mathcal{U}$
Terms	$M, N, T ::= \star \mid x \mid$ $\mathbb{N} \mid Z \mid S M \mid$ $\text{case}_{\mathbb{N}} M N T \mid$ $\text{Ix } M \mid \text{I}_0 \mid \text{I}_S M \mid$ $\text{case}_{\text{Ix}} M N \mid \text{case}_{\text{Ix}} M N T \mid$ $\{\!\{M_1, \dots, M_n\}\!\} \mid \lambda() \mid$ $\tau \mid \text{tt} \mid$ $\forall \alpha : T. N \mid \lambda x : T. N \mid M N \mid$ $\exists \alpha : T. M \mid \langle\langle \alpha : T, M \rangle\rangle \mid \text{case}_{\exists} M N \mid$ $M + N \mid \text{left } M \mid \text{right } M \mid$ $\text{case}_+ M N T \mid$ $M \equiv N \mid \text{refl } T M N \mid$
Environments	$\Gamma ::= \varepsilon \mid \Gamma, \alpha : T$

Fig. 1. Syntax

1.1.1 Meta-syntax & syntactic sugar. Let

- (1) $\tau \rightarrow v$ denote the non-dependent universal quantification $\forall(_ : \tau).v$;
- (2) $\tau \times v$ denote the non-dependent existential quantification $\exists(_ : \tau).v$;
- (3) $0, 1, 2, \dots$ denote object-level natural numbers in the intuitive fashion;
- (4) i_n denote the index obtained by n applications of I_S to I_0 ; and

Author's address: AH & JGM.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

© 2023 Association for Computing Machinery.

XXXX-XXXX/2023/12-ART \$15.00

<https://doi.org/10.1145/nnnnnnnn.nnnnnnnn>

(5) the syntax

$$\{\!\{M_1, \dots, M_n\}\!\}$$

denote the large elimination of a known, finite quantity of indices to types M_1, \dots, M_n , elaborated by the equations:

$$\begin{aligned}\{\!\{M_1\}\!\} &:= \lambda(i : \text{Ix } 1).\text{case}_{\text{Ix}} i M_1 \\ \{\!\{M_1, \dots, M_n\}\!\} &:= \lambda(i : \text{Ix } n).\text{case}_{\text{Ix}} i M_1 \{\!\{M_2, \dots, M_n\}\!\}\end{aligned}$$

1.2 Typing

Many of these are fucked or in need of repair; refer to the translation as the SSOT.

$$\begin{array}{c} \boxed{\vdash \Gamma} \\[10pt] (\text{EMP}) \frac{}{\vdash \varepsilon} \quad (\text{VAR}) \frac{\vdash \Gamma \quad \Gamma \vdash M : \sigma}{\vdash \Gamma, x : M} \\[10pt] \boxed{\Gamma \vdash M : \sigma} \\[10pt] \begin{array}{llll} (\star) \frac{\vdash \Gamma}{\Gamma \vdash \star : \mathcal{U}} & (\top) \frac{\vdash \Gamma}{\Gamma \vdash \top : \sigma} & (\text{NAT}) \frac{\vdash \Gamma}{\Gamma \vdash \mathbb{N} : \star} & (\text{IX}) \frac{\Gamma \vdash n : \mathbb{N}}{\Gamma \vdash \text{Ix } n : \star} \\[10pt] (\forall) \frac{\Gamma \vdash M : \sigma_1 \quad \Gamma, \alpha : M \vdash N : \sigma_2}{\Gamma \vdash \forall \alpha : M.N : \sigma_2} & (\exists) \frac{\Gamma \vdash M : \sigma_1 \quad \Gamma, \alpha : M \vdash N : \sigma_2}{\Gamma \vdash \exists \alpha : M.N : \sigma_2} & & \\[10pt] (+) \frac{\Gamma \vdash M : \sigma \quad \Gamma \vdash N : \sigma}{\Gamma \vdash M + N : \sigma} & (\equiv) \frac{\Gamma \vdash M : \sigma \quad \Gamma \vdash N : \sigma}{\Gamma \vdash M \equiv N : \sigma} & & \end{array} \end{array}$$

Fig. 2. Context and type well-formedness

$$\boxed{\Gamma \vdash M : N}$$

$$\begin{array}{c}
(\text{VAR}) \frac{\vdash \Gamma \quad x : M \in \Gamma}{\Gamma \vdash x : M} \quad (\text{tt}) \frac{\vdash \Gamma}{\Gamma \vdash \text{tt} : \top} \\
(Z) \frac{\vdash \Gamma}{\Gamma \vdash Z : \mathbb{N}} \quad (S) \frac{\Gamma \vdash n : \mathbb{N}}{\Gamma \vdash S n : \mathbb{N}} \quad (\mathbb{N}E) \frac{\Gamma \vdash A : \sigma \quad \Gamma \vdash M : \mathbb{N} \quad \Gamma \vdash N : A \quad \Gamma \vdash P : \mathbb{N} \rightarrow A}{\Gamma \vdash \text{case}_{\mathbb{N}} M N P : A} \\
(l_0) \frac{\Gamma \vdash n : \mathbb{N}}{\Gamma \vdash l_0 : \text{Ix}(S n)} \quad (l_S) \frac{\Gamma \vdash n : \mathbb{N} \quad \Gamma \vdash i : \text{Ix } n}{\Gamma \vdash l_S i : \text{Ix}(S n)} \quad (\lambda()) \frac{\Gamma \vdash A : \sigma}{\Gamma \vdash \lambda() : \text{Ix } 0 \rightarrow A} \\
(l_S E) \frac{\Gamma \vdash A : \sigma \quad \Gamma \vdash M : \text{Ix}(S n) \quad \Gamma \vdash N : A \quad \Gamma \vdash P : \text{Ix } n \rightarrow A}{\Gamma \vdash \text{case}_{\text{Ix}} M N P : A} \\
(\forall I) \frac{\Gamma \vdash T : \sigma \quad \Gamma, x : T \vdash M : N}{\Gamma \vdash \lambda x : T. M : \forall(x : T). N} \quad (\forall E) \frac{\Gamma \vdash M : \forall(x : T_1). T_2 \quad \Gamma \vdash N : T_1}{\Gamma \vdash M N : T_2[N/x]} \\
(\exists I) \frac{\Gamma \vdash T_1 : \sigma \quad \Gamma \vdash M : T_1 \quad \Gamma \vdash N : T_2[M/x]}{\Gamma \vdash \langle\langle M : T_1, N \rangle\rangle : \exists(x : T_1). T_2} \\
(\exists E) \frac{\Gamma \vdash A : \sigma \quad \Gamma \vdash M : \Sigma(x : T_1). T_2 \quad \Gamma \vdash N : \forall(x : T_1). T_2 \rightarrow A}{\Gamma \vdash \text{case}_{\exists} M N : A} \\
(+_1 I) \frac{\Gamma \vdash M : A}{\Gamma \vdash \text{left } M : A + B} \quad (+_2 I) \frac{\Gamma \vdash N : B}{\Gamma \vdash \text{right } N : A + B} \\
(+E) \frac{\Gamma \vdash C : \sigma \quad \Gamma \vdash M : A + B \quad \Gamma \vdash N : A \rightarrow C \quad \Gamma \vdash P : B \rightarrow C}{\Gamma \vdash \text{case}_+ M N P : C} \\
(\equiv I) \frac{\Gamma \vdash M : \sigma}{\Gamma \vdash \text{refl} : M \equiv M} \quad (\text{CONV}) \frac{\Gamma \vdash M : T_1 \quad \Gamma \vdash T_1 = T_2 : \sigma}{\Gamma \vdash M : T_2} \\
\begin{array}{c}
\Gamma \vdash P : T_1 \equiv T_2 \\
\Gamma \vdash M : T_1 \\
\Gamma \vdash N : T_1 \\
\Gamma, x : T_1, y : T_1, p : x \equiv y \vdash T : \star \\
\Gamma, z : T_1 \vdash H : T[z/x, z/y, \text{refl}/p] \\
(\equiv E) \frac{}{\Gamma \vdash \mathcal{J} H M N P : T[M/x, N/y, P/p]}
\end{array}
\end{array}$$

Fig. 3. Typing lx terms

$$\boxed{\Gamma \vdash M = N : \sigma}$$

$$\begin{array}{c}
(\text{E-REFL}) \frac{\Gamma \vdash M : \sigma}{\Gamma \vdash M = M : \sigma} \quad (\text{E-SYM}) \frac{\Gamma \vdash N = M : \sigma}{\Gamma \vdash M = N : \sigma} \quad (\text{E-TRANS}) \frac{\Gamma \vdash M = P : \sigma \quad \Gamma \vdash P = N : \sigma}{\Gamma \vdash M = N : \sigma} \\
\boxed{\Gamma \vdash M = N : T} \\
(\text{C-REFL}) \frac{\Gamma \vdash M : T}{\Gamma \vdash M = M : T} \quad (\text{C-SYM}) \frac{\Gamma \vdash N = M : T}{\Gamma \vdash M = N : T} \quad (\text{C-TRANS}) \frac{\Gamma \vdash M = P : T \quad \Gamma \vdash P = N : T}{\Gamma \vdash M = N : T}
\end{array}$$

Fig. 4. Definitional equality & computational laws

1.3 Properties

THEOREM 1 (WELL-SORTEDNESS). *if $\Gamma \vdash M : N$ then $\vdash \Gamma$ and there exists σ such that $\Gamma \vdash N : \sigma$.*

2 TRANSLATION FROM $R\omega$

2.1 Untyped Translation

We follow the approach of [Morris and McKinna 2019] and give both typed and untyped translations of $R\omega$ types. Figure 5 describe the untyped translation, which is used to show translational soundness of the typed translation (Figure 5).

$$\boxed{\llbracket \kappa \rrbracket}$$

$$\llbracket \star \rrbracket = \star$$

$$\llbracket \mathbf{L} \rrbracket = \top$$

$$\llbracket \kappa_1 \rightarrow \kappa_2 \rrbracket = \llbracket \kappa_1 \rrbracket \rightarrow \llbracket \kappa_2 \rrbracket$$

$$\llbracket \mathbf{R}^\kappa \rrbracket = \exists (n : \mathbb{N}). \text{Ix } n \rightarrow \llbracket \kappa \rrbracket$$

$$\boxed{\llbracket \Gamma \vdash \tau : \kappa \rrbracket}$$

$$\llbracket \Gamma \vdash \alpha : \kappa \rrbracket = \alpha$$

$$\llbracket \Gamma \vdash \tau_1 \rightarrow \tau_2 : \star \rrbracket = \llbracket \tau_1 \rrbracket \rightarrow \llbracket \tau_2 \rrbracket$$

$$\llbracket \Gamma \vdash \forall \alpha : \kappa. \tau : \star \rrbracket = \forall (\alpha : \llbracket \kappa \rrbracket). \llbracket \tau \rrbracket$$

$$\llbracket \Gamma \vdash \lambda \alpha : \kappa. \tau : \kappa \rightarrow \kappa' \rrbracket = \forall (\alpha : \llbracket \kappa \rrbracket). \llbracket \tau \rrbracket$$

$$\llbracket \Gamma \vdash \pi \Rightarrow \tau : \kappa \rrbracket = \forall (\alpha : \llbracket \pi \rrbracket). \llbracket \tau \rrbracket$$

$$\llbracket \Gamma \vdash \tau v : \kappa \rrbracket = \llbracket \tau \rrbracket \llbracket v \rrbracket$$

$$\llbracket \Gamma \vdash \ell : \mathbf{L} \rrbracket = \top$$

$$\llbracket \Gamma \vdash \lfloor \xi \rfloor : \star \rrbracket = \top$$

$$\llbracket \Gamma \vdash (\xi \triangleright \tau) : \kappa \rrbracket = \llbracket \tau \rrbracket$$

$$\llbracket \Gamma \vdash \Pi \rho : \star \rrbracket = \text{case}_{\exists} \llbracket \rho \rrbracket (\lambda n : \mathbb{N}. \lambda P : \text{Ix } n \rightarrow \star. \forall (i : \text{Ix } n). P i)$$

$$\llbracket \Gamma \vdash \Sigma \rho : \star \rrbracket = \text{case}_{\exists} \llbracket \rho \rrbracket (\lambda n : \mathbb{N}. \lambda P : \text{Ix } n \rightarrow \star. \exists (i : \text{Ix } n). P i)$$

$$\llbracket \Gamma \vdash \epsilon : \mathbf{R}^\kappa \rrbracket = \langle \langle 0 : \mathbb{N}, \lambda () \rangle \rangle$$

$$\llbracket \Gamma \vdash \rho [v] : \mathbf{R}^{\kappa_2} \rrbracket = \text{case}_{\exists} \llbracket \rho \rrbracket (\lambda n : \mathbb{N}. \lambda (P : \text{Ix } n \rightarrow \llbracket \kappa_1 \rrbracket \rightarrow \llbracket \kappa_2 \rrbracket). \langle \langle n : \mathbb{N}, \lambda (j : \text{Ix } n). (P j) \llbracket \tau \rrbracket \rangle \rangle)$$

$$\llbracket \Gamma \vdash [\tau] \rho : \mathbf{R}^{\kappa_2} \rrbracket = \text{case}_{\exists} \llbracket \rho \rrbracket (\lambda n : \mathbb{N}. \lambda (P : \text{Ix } n \rightarrow \llbracket \kappa_1 \rrbracket). \langle \langle n : \mathbb{N}, \lambda (j : \text{Ix } n). \llbracket \tau \rrbracket (P j) \rangle \rangle)$$

$$\llbracket \Gamma \vdash (\xi \triangleright_{\mathbf{R}} \tau) : \mathbf{R}^\kappa \rrbracket = \langle \langle 1 : \mathbb{N}, \llbracket \tau \rrbracket \rangle \rangle$$

$$\boxed{\llbracket \Gamma \vdash \pi : \kappa \rrbracket \text{ (Predicate well-formedness)}}$$

$$\text{case } \llbracket \rho_1 \rrbracket (\lambda n : \mathbb{N}. \lambda (P : \text{Ix } n \rightarrow \llbracket \kappa \rrbracket).$$

$$\llbracket \Gamma \vdash \rho_1 \lesssim \rho_2 : \kappa \rrbracket = \text{case } \llbracket \rho_2 \rrbracket (\lambda m : \mathbb{N}. \lambda (Q : \text{Ix } m \rightarrow \llbracket \kappa \rrbracket). \\ \forall (i : \text{Ix } n). \exists (j : \text{Ix } m). P i \equiv Q j))$$

$$\text{case } \llbracket \rho_1 \rrbracket (\lambda n : \mathbb{N}. \lambda (P : \text{Ix } n \rightarrow \llbracket \kappa \rrbracket).$$

$$\text{case } \llbracket \rho_2 \rrbracket (\lambda m : \mathbb{N}. \lambda (Q : \text{Ix } m \rightarrow \llbracket \kappa \rrbracket).$$

$$\text{case } \llbracket \rho_3 \rrbracket (\lambda l : \mathbb{N}. \lambda (R : \text{Ix } l \rightarrow \llbracket \kappa \rrbracket).$$

$$(\forall (i : \text{Ix } n). \exists (k : \text{Ix } l). P i \equiv R k)$$

$$\llbracket \Gamma \vdash \rho_1 \odot \rho_2 \sim \rho_3 : \kappa \rrbracket = \times (\forall (j : \text{Ix } m). \exists (k : \text{Ix } l). Q j \equiv R k)$$

$$\times (\forall (k : \text{Ix } l).$$

$$(\exists (i : \text{Ix } n). P i \equiv R k)$$

$$+ (\exists (j : \text{Ix } m). Q j \equiv R k))))$$

Fig. 5. A compositional translation of typed $\mathbf{R}\omega$ kinds and predicates to untyped Ix terms

2.2 Typed translation

$$\begin{array}{c}
 \boxed{\Gamma \vdash \tau \rightsquigarrow v : \kappa} \\
 \\
 \text{(C-FOO)} \frac{A}{B} \\
 \boxed{\Gamma \vdash M \rightsquigarrow N : \tau} \\
 \\
 \text{(C-FOO)} \frac{A}{B} \\
 \boxed{\Gamma \Vdash \pi \rightsquigarrow N} \\
 \\
 \text{(C-FOO)} \frac{A}{B} \\
 \boxed{\tau \equiv v \rightsquigarrow P} \\
 \\
 \text{(C-FOO)} \frac{A}{B}
 \end{array}$$

Fig. 6. Translation of $R\omega$ derivations to lx derivations

2.3 Properties of Translation

Presume an $R\omega$ instantiation of the simple row theory. A lot of this is likely bullshit.

THEOREM 2 (TRANSLATIONAL SOUNDNESS (TYPES)). *if $\Gamma \vdash \tau : \kappa$ such that $\Gamma \vdash \tau \rightsquigarrow v : \kappa$ then $\llbracket \Gamma \rrbracket \vdash v : \llbracket \kappa \rrbracket$.*

THEOREM 3 (TRANSLATIONAL SOUNDNESS (TYPE EQUIVALENCE)). *if*

- (1) $\Gamma \vdash \tau_1 \rightsquigarrow v_1 : \kappa_1$;
- (2) $\Gamma \vdash \tau_2 \rightsquigarrow v_2 : \kappa_2$; and
- (3) $\tau_1 \equiv \tau_2 \rightsquigarrow P$,

then $\llbracket \Gamma \rrbracket \vdash P : v_1 \equiv v_2$.

THEOREM 4 (TRANSLATIONAL SOUNDNESS (PREDICATES)). *if $\Gamma \Vdash \pi$ such that $\Gamma \Vdash \pi \rightsquigarrow N$ then $\llbracket \Gamma \rrbracket \vdash N : \llbracket \pi \rrbracket$.*

Finally,

THEOREM 5 (TRANSLATIONAL SOUNDNESS). *if $\Gamma \vdash M : \tau$ such that $\Gamma \vdash M \rightsquigarrow N : \tau$ then $\llbracket \Gamma \rrbracket \vdash N : \llbracket \tau \rrbracket$.*

3 OPERATIONAL SEMANTICS

REFERENCES

- J. Garrett Morris and James McKinna. 2019. Abstracting extensible data types: or, rows by any other name. *Proc. ACM Program. Lang.* 3, POPL (2019), 12:1–12:28. <https://doi.org/10.1145/3290325>