


School of Physics and Materials Science
Thapar Institute of Engineering & Technology, Patiala
APPLIED PHYSICS (UPH 004)

TUTORIAL SHEET # 1: OSCILLATIONS

1. A mass of 3 kg is attached to the free end of a spring as shown in the figure. For this spring, a force of 6 N causes a displacement of 0.03 m. If the mass is pulled aside by a distance of 0.06 m and released, then



- (a) Compute the force constant of the spring.
 - (b) Compute the period and frequency of vibration.
 - (c) Compute the maximum velocity attained by the vibrating mass.
 - (d) Compute the maximum acceleration.
 - (e) Compute the velocity and acceleration when the mass has moved halfway toward the centre from its initial position.
 - (f) How much time is required for the mass to move halfway to the centre from its initial position?
2. A 50 g mass hangs at the end of a spring. When 20 g more are added to the end of the spring, it stretches 5 cm more.
- (a) Find the spring constant.
 - (b) If the 20 g mass is now removed, what will be the period of the motion?
3. A block of mass 4 kg hangs from a spring of Force constant $k = 400 \text{ N/m}$. The block is pulled down 15 cm below equilibrium and released. Find
- (a) Amplitude, frequency and period of motion.
 - (b) Kinetic energy when the block is 10 cm above equilibrium.
4. Compute the oscillation frequency of LCR circuit having $L = 0.1 \text{ mH}$, $C = 20 \mu\text{F}$, $R = 0.1 \text{ ohm}$. What will happen if R is made zero?
5. A simple pendulum executes 90 oscillations per minute. It takes 15 oscillations to reduce its amplitude, which is 15° , by 50%. Find relaxation time of the pendulum.
6. A galvanometer coil oscillates with a periodic time of 5 second and successive maximum displacements are observed at 76.0, 34.2, 15.5, and 6.9 scale divisions. Compute
- (a) Logarithmic decrement.
 - (b) Damping factor.
 - (c) Deflection of the galvanometer coil in the absence of damping.