

Quantifying Uncertainty

Advanced Artificial Intelligence: Lecture

Heriberto Cuayáhuatl

<http://staff.lincoln.ac.uk/hcuayahuitl>

School of Computer Science, University of Lincoln



UNIVERSITY OF
LINCOLN

2 October 2019

About Me



- ▶ Dr. Heriberto Cuayahuitl
- ▶ PhD in informatics, University of Edinburgh, UK
- ▶ Research Interests:
 - ▶ Machine Intelligence
 - ▶ Dialogue Systems
 - ▶ Interactive Robots
- ▶ Surgery Hours: Wednesday 13:00-14:00, or arrange meeting
- ▶ Office Location: Isaac Newton Building, room INB3130
- ▶ Email: HCuayahuitl@lincoln.ac.uk

Overview

Topics:

- ▶ Introduction to Probability
- ▶ Probabilistic Reasoning

Readings:

1. Russel, S.& Norvig, P. (2013). “Artificial Intelligence: A Modern Approach”, **chapter13** (main reading)
2. Bishop, C. “Model-Based Machine Learning”, **chapter1** (optional reading)

Probability Basics

- ▶ Let Ω be the **sample space**, e.g., 6 possible rolls of a die, and $\omega \in \Omega$ a **sample point** or **possible world** or **atomic event**

Probability Basics

- ▶ Let Ω be the **sample space**, e.g., 6 possible rolls of a die, and $\omega \in \Omega$ a **sample point** or **possible world** or **atomic event**
- ▶ A **probability space** or **probability model** is a sample space with an assignment $P(\omega)$ for every $\omega \in \Omega$ s.t. $0 \leq P(\omega) \leq 1$
 $\sum_{\omega} P(\omega)=1$, e.g. $P(1)=P(2)=P(3)=P(4)=P(5)=P(6)=1/6$

Probability Basics

- ▶ Let Ω be the **sample space**, e.g., 6 possible rolls of a die, and $\omega \in \Omega$ a **sample point** or **possible world** or **atomic event**
- ▶ A **probability space** or **probability model** is a sample space with an assignment $P(\omega)$ for every $\omega \in \Omega$ s.t. $0 \leq P(\omega) \leq 1$
 $\sum_{\omega} P(\omega)=1$, e.g. $P(1)=P(2)=P(3)=P(4)=P(5)=P(6)=1/6$
- ▶ An **event** A is any subset of Ω , where $P(A)=\sum_{\omega \in A} P(\omega)$ e.g.
 $P(\text{die roll} < 4) = P(1) + P(2) + P(3) = 1/6 + 1/6 + 1/6 = 1/2$

Probability Basics

- ▶ Let Ω be the **sample space**, e.g., 6 possible rolls of a die, and $\omega \in \Omega$ a **sample point** or **possible world** or **atomic event**
- ▶ A **probability space** or **probability model** is a sample space with an assignment $P(\omega)$ for every $\omega \in \Omega$ s.t. $0 \leq P(\omega) \leq 1$
 $\sum_{\omega} P(\omega) = 1$, e.g. $P(1) = P(2) = P(3) = P(4) = P(5) = P(6) = 1/6$
- ▶ An **event** A is any subset of Ω , where $P(A) = \sum_{\omega \in A} P(\omega)$ e.g.
 $P(\text{die roll} < 4) = P(1) + P(2) + P(3) = 1/6 + 1/6 + 1/6 = 1/2$
- ▶ Variables in probability theory are called **random variables**, their names begin with an uppercase letter (e.g. $Die_1 = \{1, \dots, 6\}$, and have domain values (e.g. $\{true, false\}$)

Propositions

- ▶ Think of a proposition as an event (set of sample points) where the proposition is true

Propositions

- ▶ Think of a proposition as an event (set of sample points) where the proposition is true
- ▶ Given Boolean random variables A and B :
 - ▶ event a = set of sample points where $A(\omega) = \text{true}$
 - ▶ event $\neg a$ = set of sample points where $A(\omega) = \text{false}$
 - ▶ event $a \wedge b$ = points where $A(\omega) = \text{true}$ and $B(\omega) = \text{true}$

Propositions

- ▶ Think of a proposition as an event (set of sample points) where the proposition is true
- ▶ Given Boolean random variables A and B :
 - ▶ event a = set of sample points where $A(\omega) = \text{true}$
 - ▶ event $\neg a$ = set of sample points where $A(\omega) = \text{false}$
 - ▶ event $a \wedge b$ = points where $A(\omega) = \text{true}$ and $B(\omega) = \text{true}$
- ▶ Sample points are *defined* by the values of random variables, i.e. the sample space is the Cartesian product of their ranges

Propositions

- ▶ Think of a proposition as an event (set of sample points) where the proposition is true
- ▶ Given Boolean random variables A and B :
 - ▶ event a = set of sample points where $A(\omega) = \text{true}$
 - ▶ event $\neg a$ = set of sample points where $A(\omega) = \text{false}$
 - ▶ event $a \wedge b$ = points where $A(\omega) = \text{true}$ and $B(\omega) = \text{true}$
- ▶ Sample points are *defined* by the values of random variables, i.e. the sample space is the Cartesian product of their ranges
- ▶ Proposition = disjunction of atomic events in which it is true, e.g. $(a \vee b) \equiv (\neg a \wedge b) \vee (a \wedge \neg b) \vee (a \wedge b)$
 $\implies P(a \vee b) = P(\neg a \wedge b) + P(a \wedge \neg b) + P(a \wedge b)$

Syntax for Propositions

- ▶ **Boolean** random variables
e.g., *Cavity* (do I have a cavity?)
Cavity = *true* is a proposition, also written *cavity*

Syntax for Propositions

- ▶ **Boolean** random variables
e.g., *Cavity* (do I have a cavity?)
Cavity = *true* is a proposition, also written *cavity*
- ▶ **Discrete** random variables (*finite* or *infinite*)
e.g., *Weather* is one of $\langle \textit{sunny}, \textit{rain}, \textit{cloudy}, \textit{snow} \rangle$
Weather = *rain* is a proposition
Values must be exhaustive and mutually exclusive

Syntax for Propositions

- ▶ **Boolean** random variables
e.g., *Cavity* (do I have a cavity?)
Cavity = *true* is a proposition, also written *cavity*
- ▶ **Discrete** random variables (*finite* or *infinite*)
e.g., *Weather* is one of $\langle \textit{sunny}, \textit{rain}, \textit{cloudy}, \textit{snow} \rangle$
Weather = *rain* is a proposition
Values must be exhaustive and mutually exclusive
- ▶ **Continuous** random variables (*bounded* or *unbounded*)
e.g., *Temp* = 21.6; also allow, e.g., *Temp* < 22.0.

Prior Probability and (Joint) Probability Distribution

- ▶ **Prior** or **unconditional probabilities** of propositions, e.g. $P(\text{Cavity} = \text{true}) = 0.1$ and $P(\text{Weather} = \text{sunny}) = 0.72$ correspond to belief prior to arrival of any (new) evidence

¹Every question about a domain can be answered by the joint distribution because every event is a sum of sample points

Prior Probability and (Joint) Probability Distribution

- ▶ **Prior** or **unconditional probabilities** of propositions, e.g. $P(\text{Cavity} = \text{true}) = 0.1$ and $P(\text{Weather} = \text{sunny}) = 0.72$ correspond to belief prior to arrival of any (new) evidence
- ▶ **Probability distribution** gives values for all possible assignments: $\mathbf{P}(\text{Weather}) = \langle 0.72, 0.1, 0.08, 0.1 \rangle$ (*normalised*, i.e. sums to 1)

¹Every question about a domain can be answered by the joint distribution because every event is a sum of sample points

Prior Probability and (Joint) Probability Distribution

- ▶ **Prior** or **unconditional probabilities** of propositions, e.g. $P(\text{Cavity} = \text{true}) = 0.1$ and $P(\text{Weather} = \text{sunny}) = 0.72$ correspond to belief prior to arrival of any (new) evidence
- ▶ **Probability distribution** gives values for all possible assignments: $\mathbf{P}(\text{Weather}) = \langle 0.72, 0.1, 0.08, 0.1 \rangle$ (*normalised*, i.e. sums to 1)
- ▶ **Joint probability distribution**¹ gives the probability of every atomic event on a set of random variables (i.e. every sample point), e.g. $\mathbf{P}(\text{Weather}, \text{Cavity}) =$ a 4×2 matrix of values:

Weather =	sunny	rainy	cloudy	snowy
Cavity = true	0.144	0.02	0.016	0.02
Cavity = false	0.576	0.08	0.064	0.08

¹Every question about a domain can be answered by the joint distribution because every event is a sum of sample points

Conditional Probability (1/3)

- ▶ **Conditional or posterior probabilities** e.g.,
 $P(\text{cavity}|\text{toothache}) = 0.8$, i.e. *given that toothache is all I know ... NOT* “if toothache then 80% chance of cavity”

Conditional Probability (1/3)

- ▶ **Conditional or posterior probabilities** e.g.,
 $P(\text{cavity}|\text{toothache}) = 0.8$, i.e. *given that toothache is all I know ... NOT* “if toothache then 80% chance of cavity”
- ▶ Notation for conditional distributions:
 $\mathbf{P}(\text{Cavity}|\text{toothache}) =$
 $\langle P(\text{cavity}|\text{toothache}), P(\neg\text{cavity}|\text{toothache}) \rangle$

Conditional Probability (1/3)

- ▶ **Conditional or posterior probabilities** e.g.,
 $P(\text{cavity}|\text{toothache}) = 0.8$, i.e. *given that toothache is all I know ... NOT “if toothache then 80% chance of cavity”*
- ▶ Notation for conditional distributions:
 $\mathbf{P}(\text{Cavity}|\text{toothache}) =$
 $< P(\text{cavity}|\text{toothache}), P(\neg\text{cavity}|\text{toothache}) >$
- ▶ New evidence may be irrelevant, allowing simplification, e.g.
 $P(\text{cavity}|\text{toothache}, \text{Weather}) = P(\text{cavity}|\text{toothache}) = 0.8$

Conditional Probability (2/3)

- ▶ Conditional probability: $P(a|b) = \frac{P(a \wedge b)}{P(b)}$ if $P(b) \neq 0$

Conditional Probability (2/3)

- ▶ Conditional probability: $P(a|b) = \frac{P(a \wedge b)}{P(b)}$ if $P(b) \neq 0$
- ▶ The **product rule** gives an alternative formulation:
$$P(a \wedge b) = P(a|b)P(b) = P(b|a)P(a)$$

Conditional Probability (2/3)

- ▶ Conditional probability: $P(a|b) = \frac{P(a \wedge b)}{P(b)}$ if $P(b) \neq 0$
- ▶ The **product rule** gives an alternative formulation:
 $P(a \wedge b) = P(a|b)P(b) = P(b|a)P(a)$
- ▶ A general version holds for whole distributions, e.g.,
 $\mathbf{P}(\text{Weather}, \text{Cavity}) = \mathbf{P}(\text{Weather}|\text{Cavity})\mathbf{P}(\text{Cavity})$
(View as a 4×2 set of equations, *not* matrix multiplication)

$$P(W = \text{sunny} \wedge C = \text{true}) = P(W = \text{sunny}|C = \text{true}) P(C = \text{true})$$

$$P(W = \text{rain} \wedge C = \text{true}) = P(W = \text{rain}|C = \text{true}) P(C = \text{true})$$

$$P(W = \text{cloudy} \wedge C = \text{true}) = P(W = \text{cloudy}|C = \text{true}) P(C = \text{true})$$

$$P(W = \text{snow} \wedge C = \text{true}) = P(W = \text{snow}|C = \text{true}) P(C = \text{true})$$

$$P(W = \text{sunny} \wedge C = \text{false}) = P(W = \text{sunny}|C = \text{false}) P(C = \text{false})$$

$$P(W = \text{rain} \wedge C = \text{false}) = P(W = \text{rain}|C = \text{false}) P(C = \text{false})$$

$$P(W = \text{cloudy} \wedge C = \text{false}) = P(W = \text{cloudy}|C = \text{false}) P(C = \text{false})$$

$$P(W = \text{snow} \wedge C = \text{false}) = P(W = \text{snow}|C = \text{false}) P(C = \text{false}) .$$

Conditional Probability (3/3)

- ▶ The **chain rule** is derived by the successive application of the product rule:
 - ▶ $P(X_1, X_2) = P(X_2|X_1)P(X_1)$
 - ▶ $P(X_1, X_2, X_3) = P(X_3|X_2, X_1)P(X_2, X_1)$
 - ▶ $\quad\quad\quad = P(X_3|X_2, X_1)P(X_2|X_1)P(X_1)$
 - ▶ $P(X_1, X_2, X_3, X_4) = ?$
 - ▶ ...
 - ▶ $P(X_1, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_2, X_1) \dots$
 - ▶ $P(X_1, \dots, X_n) = \prod_{i=1}^n P(X_i|X_{i-1}, \dots, X_1)$

Conditional Probability (3/3)

- ▶ The **chain rule** is derived by the successive application of the product rule:

- ▶ $P(X_1, X_2) = P(X_2|X_1)P(X_1)$
- ▶ $P(X_1, X_2, X_3) = P(X_3|X_2, X_1)P(X_2, X_1)$
- ▶ $\quad\quad\quad = P(X_3|X_2, X_1)P(X_2|X_1)P(X_1)$
- ▶ $P(X_1, X_2, X_3, X_4) = ?$
- ▶ ...
- ▶ $P(X_1, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_2, X_1) \dots$
- ▶ $P(X_1, \dots, X_n) = \prod_{i=1}^n P(X_i|X_{i-1}, \dots, X_1)$

- ▶ **Bayes Rule:**
$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$
$$= \frac{P(B|A)P(A)}{\sum_i^n P(B|A_i)P(A_i)}$$

Marginal Probability

- ▶ Marginalisation is the process of summing up the probabilities of the other variables—i.e. taking them out of the equation.

Marginal Probability

- ▶ Marginalisation is the process of summing up the probabilities of the other variables—i.e. taking them out of the equation.
- ▶ General rule for any set of variables Y and Z :
$$P(Y) = \sum_{z \in Z} P(Y, z)$$

Marginal Probability

- ▶ Marginalisation is the process of summing up the probabilities of the other variables—i.e. taking them out of the equation.
- ▶ General rule for any set of variables Y and Z :
$$\mathbf{P}(Y) = \sum_{z \in Z} \mathbf{P}(Y, z)$$
- ▶ For example: $\mathbf{P}(Cavity) = \sum_{z \in \{Catch, Toothache\}} \mathbf{P}(Cavity, z)$

Marginal Probability

- ▶ Marginalisation is the process of summing up the probabilities of the other variables—i.e. taking them out of the equation.

- ▶ General rule for any set of variables Y and Z :

$$P(Y) = \sum_{z \in Z} P(Y, z)$$

- ▶ For example: $P(\text{Cavity}) = \sum_{z \in \{\text{Catch}, \text{Toothache}\}} P(\text{Cavity}, z)$

- ▶ Consider the following joint distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

Inference by Enumeration (1/3)

Start with the joint distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

For any proposition ϕ , sum the atomic events where it is true:

$$P(\phi) = \sum_{\omega: \omega \models \phi} P(\omega)$$

$$P(\text{toothache}) = 0.108 + 0.012 + 0.016 + 0.064 = 0.2$$

Inference by Enumeration (2/3)

Start with the joint distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

For any proposition ϕ , sum the atomic events where it is true:

$$P(\phi) = \sum_{\omega: \omega \models \phi} P(\omega)$$

$$P(\text{cavity} \vee \text{toothache}) =$$

$$0.108 + 0.012 + 0.072 + 0.008 + 0.016 + 0.064 = 0.28$$

Inference by Enumeration (3/3)

Start with the joint distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

We can also compute conditional probabilities:

$$\begin{aligned}P(\neg \text{cavity} | \text{toothache}) &= \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} \\&= \frac{0.016 + 0.064}{0.108 + 0.012 + 0.016 + 0.064} = 0.4\end{aligned}$$

Normalisation

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

The denominator can be viewed as a *normalization constant* α

$$\begin{aligned}\mathbf{P}(\text{Cavity}|\text{toothache}) &= \alpha \mathbf{P}(\text{Cavity}, \text{toothache}) \\ &= \alpha [\mathbf{P}(\text{Cavity}, \text{toothache}, \text{catch}) + \mathbf{P}(\text{Cavity}, \text{toothache}, \neg \text{catch})] \\ &= \alpha [< 0.108, 0.016 > + < 0.012, 0.064 >] \\ &= \alpha < 0.12, 0.08 > = < 0.6, 0.4 >\end{aligned}$$

General idea: compute distribution on query variable by fixing **evidence variables** and summing over **hidden variables**

Keywords

- ▶ **Probability Distribution**
- ▶ **Joint Probability**
- ▶ **Conditional Probability**
- ▶ **Prior Probability**
- ▶ **Posterior Probability**
- ▶ **Product Rule**
- ▶ **Sum Rule**
- ▶ **Chain Rule**
- ▶ **Marginalisation**
- ▶ **Normalisation**
- ▶ **Inference**
- ▶ **Bayes Rule**
- ▶ **Probabilistic Model**

Workshop:

- ▶ Exercises about quantifying uncertainty, Q&A
- ▶ Wednesday from 10:30-12hrs, room=INB1301

Readings:

1. Russel, S.& Norvig, P. (2013). "Artificial Intelligence: A Modern Approach", **chapter13** (main reading)
2. Bishop, C. "Model-Based Machine Learning", **chapter1** (optional reading)