


# Intro to Quantum Computing

*Quantum Circuits*



# Intro to Quantum Logic and Quantum Circuits

- Most of the stuff we will talk today can be found in *Nielsen & Chuang's* textbook (Chapter 4)

# Intro to Quantum Logic and Quantum Circuits

- Most of the stuff we will talk today can be found in *Nielsen & Chuang's* textbook (Chapter 4)
- But in much (much...) more detail.

# Intro to Quantum Logic and Quantum Circuits

- Most of the stuff we will talk today can be found in *Nielsen & Chuang's* textbook (Chapter 4)
- But in much (much...) more detail.
- Also, Preskill's Lecture notes (Chapter 5 [http://theory.caltech.edu/~preskill/ph219/chap5\\_15.pdf](http://theory.caltech.edu/~preskill/ph219/chap5_15.pdf)) contain a very thorough discussion and we will borrow some of the exposition from him

# What this is all about...

- Quantum Mechanical events are not *just* physical phenomena.

# What this is all about...

- Quantum Mechanical events are not *just* physical phenomena.
- Quantum Mechanics is a *Physical Theory*

# What this is all about...

- Quantum Mechanical events are not *just* physical phenomena.
- Quantum Mechanics is a *Physical Theory*
- Just like *Logic....*

# What this is all about...

- Quantum Mechanical events are not *just* physical phenomena.
- Quantum Mechanics is a *Physical Theory*
- Just like *Logic....*
- The rules of QM can also be interpreted as *executable operations in a programmable machine.*



# What this is all about...

- Quantum Mechanical events are not *just* physical phenomena.
- Quantum Mechanics is a *Physical Theory*
- Just like *Logic*....
- The rules of QM can also be interpreted as *executable operations in a programmable machine*.
- Just like a classical computing machine is a bunch of logical circuits, a quantum computing machine is a bunch of *quantum logical circuits (gates)*

# Quantum Logic

- Before we describe circuits (or gates), we will talk a bit about *quantum logic* and *quantum propositions*.

# Quantum Logic

- Before we describe circuits (or gates), we will talk a bit about *quantum logic* and *quantum propositions*.
- The evaluation of quantum propositions is a bit different than classical proposition that you learn in Discrete Maths.

# Quantum Logic

- Before we describe circuits (or gates), we will talk a bit about *quantum logic* and *quantum propositions*.
- The evaluation of quantum propositions is a bit different than classical proposition that you learn in Discrete Maths.
- (Today is Sunday) **or** (today I learn about QM) is a *classical* proposition that evaluations to *true*.


# Quantum Logic

- Before we describe circuits (or gates), we will talk a bit about *quantum logic* and *quantum propositions*.
- The evaluation of quantum propositions is a bit different than classical proposition that you learn in Discrete Maths.
- (Today is Sunday) **or** (today I learn about QM) is a *classical* proposition that evaluations to *true*.
- (Today is Sunday) **and** (today I learn about QM) is a classical proposition that evaluations to *false*.

# Speaking about Logic...



# Speaking about Logic...



# Intro to Quantum Logic

- Consider an electron and let's consider the following two propositions



# Intro to Quantum Logic

- Consider an electron and let's consider the following two propositions  
*A. The spin of the electron in the vertical direction is N-S (+1) (North pole up, South pole down).*

# Intro to Quantum Logic

- Consider an electron and let's consider the following two propositions
  - A. *The spin of the electron in the vertical direction is N-S (+1) (North pole up, South pole down).*
  - B. *The spin of the electron in the horizontal direction is N-S (+1) (North pole left, South pole right).*

# Intro to Quantum Logic

- Consider an electron and let's consider the following two propositions
  - A. The spin of the electron in the vertical direction is N-S (+1) (North pole up, South pole down).*
  - B. The spin of the electron in the horizontal direction is N-S (+1) (North pole left, South pole right).*
- How can we test these propositions?

# Intro to Quantum Logic

- Consider an electron and let's consider the following two propositions
  - A. The spin of the electron in the vertical direction is N-S (+1) (North pole up, South pole down).*
  - B. The spin of the electron in the horizontal direction is N-S (+1) (North pole left, South pole right).*
- How can we test these propositions?
- How we can test (***A or B***)? What about (***A and B***)?

# Intro to Quantum Logic

- Consider an electron and let's consider the following two propositions
  - A. The spin of the electron in the vertical direction is N-S (+1) (North pole up, South pole down).*
  - B. The spin of the electron in the horizontal direction is N-S (+1) (North pole left, South pole right).*
- How can we test these propositions?
- How we can test (**A or B**)? What about (**A and B**)?
- Let's just make the experiments and see what we get...

# Intro to Quantum Logic

- Let's consider the proposition (***A or B***).

# Intro to Quantum Logic

- Let's consider the proposition (***A or B***).
  1. Measure the spin of the electron in the vertical direction.

# Intro to Quantum Logic

- Let's consider the proposition (***A or B***).
  1. Measure the spin of the electron in the vertical direction.
  2. If it is **+1**, stop: the proposition (***A or B***) is **true**.



# Intro to Quantum Logic

- Let's consider the proposition (***A or B***).
  1. Measure the spin of the electron in the vertical direction.
  2. If it is **+1**, stop: the proposition (***A or B***) is **true**.
  3. Else, measure the spin of the electron in the horizontal direction.

# Intro to Quantum Logic

- Let's consider the proposition (***A or B***).
  1. Measure the spin of the electron in the vertical direction.
  2. If it is **+1**, stop: the proposition (***A or B***) is **true**.
  3. Else, measure the spin of the electron in the horizontal direction.
  4. If it is **+1**, stop: the proposition (***A or B***) is **true**

# Intro to Quantum Logic

- Let's consider the proposition (***A or B***).
  1. Measure the spin of the electron in the vertical direction.
  2. If it is **+1**, stop: the proposition (***A or B***) is **true**.
  3. Else, measure the spin of the electron in the horizontal direction.
  4. If it is **+1**, stop: the proposition (***A or B***) is **true**.
  5. Else it is **false**.

# Intro to Quantum Logic

- Let's consider the proposition (***A or B***).
  1. Measure the spin of the electron in the vertical direction.
  2. If it is **+1**, stop: the proposition (***A or B***) is **true**.
  3. Else, measure the spin of the electron in the horizontal direction.
  4. If it is **+1**, stop: the proposition (***A or B***) is **true**.
  5. Else it is **false**.
- Same protocol applies for measuring (***B or A***).

# Intro to Quantum Logic

- Let's consider the proposition (***A or B***).
  1. Measure the spin of the electron in the vertical direction.
  2. If it is **+1**, stop: the proposition (***A or B***) is **true**.
  3. Else, measure the spin of the electron in the horizontal direction.
  4. If it is **+1**, stop: the proposition (***A or B***) is **true**.
  5. Else it is **false**.
- Same protocol applies for measuring (***B or A***).
- The truth value of (***A or B***) is the same as (***B or A***)

# Intro to Quantum Logic

- Let's consider the proposition (***A or B***).
  1. Measure the spin of the electron in the vertical direction.
  2. If it is **+1**, stop: the proposition (***A or B***) is **true**.
  3. Else, measure the spin of the electron in the horizontal direction.
  4. If it is **+1**, stop: the proposition (***A or B***) is **true**
  5. Else it is **false**.
- Same protocol applies for measuring (***B or A***).
- The truth value of (***A or B***) is the same as (***B or A***)
- Easy-peasy, *right*?

# Intro to Quantum Logic



# Intro to Quantum Logic

- Assume that the electron has spin **+1** in the vertical direction.



# Intro to Quantum Logic

- Assume that the electron has spin **+1** in the vertical direction.
- If we measure it in the vertical direction, then we will find it **+1** and we win: (***A or B***) = ***true***.

# Intro to Quantum Logic

- Assume that the electron has spin **+1** in the vertical direction.
- If we measure it in the vertical direction, then we will find it **+1** and we win: **(A or B) = true**.

*But what if measure it instead in the horizontal direction?*

# Intro to Quantum Logic

- With probability 0.5 we will see that spin is **+1** (N-S) and with probability 0.5 the spin is **-1** (S-N).

# Intro to Quantum Logic

- With probability 0.5 we will see that spin is **+1** (N-S) and with probability 0.5 the spin is **-1** (S-N).
- If it is **+1**, we win: (***B or A***) is ***true***.

# Intro to Quantum Logic

- With probability 0.5 we will see that spin is **+1** (N-S) and with probability 0.5 the spin is **-1** (S-N).
- If it is **+1**, we win: (***B or A***) is ***true***.
- Else, we measure it's spin in the vertical direction.

# Intro to Quantum Logic

- With probability 0.5 we will see that spin is **+1** (N-S) and with probability 0.5 the spin is **-1** (S-N).
- If it is **+1**, we win: (***B or A***) is **true**.
- Else, we measure it's spin in the vertical direction.
- What is the new spin of the electron?

# Intro to Quantum Logic

- With probability 0.5 we will see that spin is **+1** (N-S) and with probability 0.5 the spin is **-1** (S-N).
- If it is **+1**, we win: (***B or A***) is **true**.
- Else, we measure it's spin in the vertical direction.
- What is the new spin of the electron?

# Intro to Quantum Logic

- With probability 0.5 we will see that spin is **+1** (N-S) and with probability 0.5 the spin is **-1** (S-N).
- If it is **+1**, we win: (***B or A***) is **true**.
- Else, we measure it's spin in the vertical direction.
- What is the new spin of the electron?
- **-1** in the horizontal direction!



# Intro to Quantum Logic

- With probability 0.5 we will see that spin is **+1** (N-S) and with probability 0.5 the spin is **-1** (S-N).
- If it is **+1**, we win: (***B or A***) is **true**.
- Else, we measure it's spin in the vertical direction.
- What is the new spin of the electron?
- **-1** in the horizontal direction!
- It has 50/50 chance to be **+1/-1** in the vertical direction!

# Intro to Quantum Logic

- With probability 0.5 we will see that spin is **+1** (N-S) and with probability 0.5 the spin is **-1** (S-N).
- If it is **+1**, we win: (***B or A***) is *true*.
- Else, we measure it's spin in the vertical direction.
- What is the new spin of the electron?
- **-1** in the horizontal direction!
- It has 50/50 chance to be **+1/-1** in the vertical direction!

***With probability 0.25 proposition (*B or A*) is false!***

# Intro to Quantum Logic

- What about (***A and B***)?

# Intro to Quantum Logic

- What about (***A and B***)?
- (think about it at home...)

# Intro to Quantum Logic

- What about (***A and B***)?
- (think about it at home...)
- The more certain we are about the first measurement, the less certain we are about the second measurement.

# Intro to Quantum Logic

- What about (***A and B***)?
- (think about it at home...)
- The more certain we are about the first measurement, the less certain we are about the second measurement.
- Sounds familiar?

# Intro to Quantum Logic

- What about (***A and B***)?
- (think about it at home...)
- The more certain we are about the first measurement, the less certain we are about the second measurement.



# Intro to Quantum Logic

- What about (***A and B***)?
- (think about it at home...)
- The more certain we are about the first measurement, the less certain we are about the second measurement.
- Sounds familiar?

***Heisenberg's Uncertainty Principle (once again...)***



# Quantum Circuits

- **Reminder**: quantum states can be described as **tensor products** of individual states, or they can be *entangled*.

# Quantum Circuits

- **Reminder**: quantum states can be described as **tensor products** of individual states, or they can be *entangled*.
- *Even* if the Qbits of a quantum state are *entangled*, we can apply *separable* operations to them.

# Quantum Circuits

- **Reminder**: quantum states can be described as **tensor products** of individual states, or they can be *entangled*.
- *Even* if the Qbits of a quantum state are *entangled*, we can apply *separable* operations to them.
- Let  $|\psi\rangle = a|00\rangle + b|01\rangle + c|10\rangle + d|11\rangle$

# Quantum Circuits


- **Reminder**: quantum states can be described as **tensor products** of individual states, or they can be *entangled*.
- *Even* if the Qbits of a quantum state are *entangled*, we can apply *separable* operations to them.
- Let  $|\psi\rangle = a|00\rangle + b|01\rangle + c|10\rangle + d|11\rangle$
- We can perform a *Hadamard* on the first Qbit only to get  $|\psi'\rangle = (H \otimes I)|\psi\rangle$ .

# Quantum Circuits

- **Reminder:** quantum states can be described as **tensor products** of individual states, or they can be *entangled*.
- *Even* if the Qbits of a quantum state are *entangled*, we can apply *separable* operations to them.
- Let  $|\psi\rangle = a|00\rangle + b|01\rangle + c|10\rangle + d|11\rangle$
- We can perform a *Hadamard* on the first Qbit only to get  $|\psi'\rangle = (H \otimes I)|\psi\rangle$ .
- We can describe it neatly graphically as a circuit:

# Quantum Circuits

- **Reminder:** quantum states can be described as **tensor products** of individual states, or they can be *entangled*.
- *Even* if the Qbits of a quantum state are *entangled*, we can apply *separable* operations to them.
- Let  $|\psi\rangle = a|00\rangle + b|01\rangle + c|10\rangle + d|11\rangle$
- We can perform a *Hadamard* on the first Qbit only to get  $|\psi'\rangle = (H \otimes I)|\psi\rangle$ .
- We can describe it neatly graphically as a circuit:




# Quantum Circuits

- Or, perform Pauli-**X** on the first Qbit only to get  $|\psi'\rangle = (X \otimes I)|\psi\rangle$ .

# Quantum Circuits


- Or, perform Pauli-**X** on the first Qbit only to get  $|\psi'\rangle = (X \otimes I)|\psi\rangle$ .
- We can describe it neatly as a circuit:





# Quantum Circuits

- Or, perform Pauli-**X** on the first Qbit only to get  $|\psi'\rangle = (X \otimes I)|\psi\rangle$ .
- We can describe it neatly as a circuit:



- We get  $|\psi'\rangle = a|10\rangle + b|11\rangle + c|00\rangle + d|01\rangle$
- **Etc.**

# Quantum Circuits

- Sometimes, these circuits are called *ansatzs* or *tensor networks*.

# Quantum Circuits

- Sometimes, these circuits are called *ansatzs* or *tensor networks*.
- We read the circuit from *left* to *right*.

# Quantum Circuits


- Sometimes, these circuits are called *ansatzs* or *tensor networks*.
- We read the circuit from *left* to *right*.
- The solid lines are *wires* and correspond to Qbits.

# Quantum Circuits

- Sometimes, these circuits are called *ansatzs* or *tensor networks*.
- We read the circuit from *left* to *right*.
- The solid lines are *wires* and correspond to Qbits.
- Boxes correspond to unitary operators.

# Quantum Circuits

- Sometimes, these circuits are called *ansatzs* or *tensor networks*.
- We read the circuit from *left* to *right*.
- The solid lines are *wires* and correspond to Qbits.
- Boxes correspond to unitary operators.
- The **CNOT** operator is denoted as



# Quantum Circuits

- **Remember**: the CNOT operator flips the value of the 2<sup>nd</sup> Qbit (*target* Qbit), if the 1<sup>st</sup> Qbit (**control** Qbit) is 1.

# Quantum Circuits

- **Remember**: the CNOT operator flips the value of the 2<sup>nd</sup> Qbit (*target* Qbit), if the 1<sup>st</sup> Qbit (**control** Qbit) is 1.
- Otherwise the 2<sup>nd</sup> Qbit is unchanged.




# Quantum Circuits

- **Remember**: the CNOT operator flips the value of the 2<sup>nd</sup> Qbit (*target* Qbit), if the 1<sup>st</sup> Qbit (**control** Qbit) is 1.
- Otherwise the 2<sup>nd</sup> Qbit is unchanged.
- $(x, y) \rightarrow (x, x \oplus y)$

# Quantum Circuits

- **Remember:** the CNOT operator flips the value of the 2<sup>nd</sup> Qbit (*target* Qbit), if the the 1<sup>st</sup> Qbit (**control** Qbit) is 1.
- Otherwise the 2<sup>nd</sup> Qbit is unchanged.
- $(x, y) \rightarrow (x, x \oplus y)$
- **Question:** *What does the following circuit do?*



# Quantum Circuits

- All Qbits are initialized to the left in the  $|0\rangle$  (or other) state.

# Quantum Circuits


- All Qbits are initialized to the left in the  $|0\rangle$  (or other) state.
- Then we add a bunch of *boxes* corresponding to *unitary transformations*.

# Quantum Circuits

- All Qbits are initialized to the left in the  $|0\rangle$  (or other) state.
- Then we add a bunch of *boxes* corresponding to *unitary transformations*.
- Finally, we have some *measurement gates* on some, or all of the Qbits.


# Quantum Circuits

- All Qbits are initialized to the left in the  $|0\rangle$  (or other) state.
- Then we add a bunch of *boxes* corresponding to *unitary transformations*.
- Finally, we have some *measurement gates* on some, or all of the Qbits.



# Quantum Circuits

- All Qbits are initialized to the left in the  $|0\rangle$  (or other) state.
- Then we add a bunch of *boxes* corresponding to *unitary transformations*.
- Finally, we have some *measurement gates* on some, or all of the Qbits.



# Matrix corresponding to a Quantum Circuit

- The Quantum circuit represents the *evolution* of our quantum system over time.



# Matrix corresponding to a Quantum Circuit


- The Quantum circuit represents the *evolution* of our quantum system over time.
- Also, appropriate matrix multiplication can describe the same evolution!

# Matrix corresponding to a Quantum Circuit


- The Quantum circuit represents the *evolution* of our quantum system over time.
- Also, appropriate matrix multiplication can describe the same evolution!
- Of course, because every step is a unitary transformation i.e., some matrix applied to the state of a Qbit (or more of them).

# Matrix corresponding to a Quantum Circuit


- The Quantum circuit represents the *evolution* of our quantum system over time.
- Also, appropriate matrix multiplication can describe the same evolution!
- Of course, because every step is a unitary transformation i.e., some matrix applied to the state of a Qbit (or more of them).
- **Question:** *What is the corresponding matrix product of the circuit?*



# More on the CNOT circuit




# More on the CNOT circuit



- What if we want to perform a CNOT on two *non-adjacent* Qbits?

# More on the CNOT circuit



- What if we want to perform a CNOT on two *non-adjacent* Qbits?

# More on the CNOT circuit

- *What if we want to perform a CNOT on two non-adjacent Qbits?*

# More on the CNOT circuit

- *What if we want to perform a CNOT on two non-adjacent Qbits?*
- In the first case, that is easy: ( $CNOT \otimes I_2$ ).



# More on the CNOT circuit

- *What if we want to perform a CNOT on two non-adjacent Qbits?*
- In the first case, that is easy: ( $CNOT \otimes I_2$ ).
- But in the second case?

# More on the CNOT circuit

- *What if we want to perform a CNOT on two non-adjacent Qbits?*
- In the first case, that is easy: ( $CNOT \otimes I_2$ ).
- But in the second case?
- Simply **SWAP, CNOT, SWAP** back!

# More on the CNOT circuit

- *What if we want to perform a CNOT on two non-adjacent Qbits?*
- In the first case, that is easy: ( $CNOT \otimes I_2$ ).
- But in the second case?
- Simply **SWAP, CNOT, SWAP** back!
- We know how to do the swap.....

# More on the CNOT circuit

- *What if we want to perform a CNOT on two non-adjacent Qbits?*
- In the first case, that is easy:  $(\mathbf{CNOT} \otimes I_2)$ .
- But in the second case?
- Simply **SWAP, CNOT, SWAP** back!
- We know how to do the swap...
- $(I_2 \otimes \mathbf{SWAP}) \times (\mathbf{CNOT} \otimes I_2) \times (I_2 \otimes \mathbf{SWAP})$

How many Quantum Gates are there?

# How many Quantum Gates are there?

- Many...

# How many Quantum Gates are there?

- Many...
- You might think.

# How many Quantum Gates are there?

- Many...
- You might think.
- Surprisingly, we only need the **CNOT** and **H**!



# How many Quantum Gates are there?

- Many...
- You might think.
- Surprisingly, we only need the **CNOT** and **H**!
- (And the **T** transform, change of phase  $T = \begin{bmatrix} 1 & 0 \\ 0 & e^{\frac{i\pi}{4}} \end{bmatrix}$ )

# How many Quantum Gates are there?

- Many...
- You might think.
- Surprisingly, we only need the **CNOT** and **H**!
- (And the **T** transform, change of phase  $T = \begin{bmatrix} 1 & 0 \\ 0 & e^{\frac{i\pi}{4}} \end{bmatrix}$ )
- Universal gates!

# How many Quantum Gates are there?

- Many...
- You might think.
- Surprisingly, we only need the **CNOT** and **H**!
- (And the **T** transform, change of phase  $T = \begin{bmatrix} 1 & 0 \\ 0 & e^{\frac{i\pi}{4}} \end{bmatrix}$ )
- Universal gates!
- **Gottesman-Kill Theorem**: Any quantum circuit consisting of **X,Y,Z,H,CNOT** can be *efficiently* simulated by a classical machine.

# Exercise to think about

- Construct a circuit that given two Qbits, both prepared in the  $|0\rangle$  state, outputs the Bell state  $\frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$