


Intro to Quantum Computing

No Cloning and Quantum Teleportation



Quantum Teleportation

Quantum Teleportation

- If you are not yet convinced that quantum mechanics is a strange theory...

Quantum Teleportation

- If you are not yet convinced that quantum mechanics is a strange theory...
- *Today you might finally change your mind* 😊

Quantum Teleportation

- If you are not yet convinced that quantum mechanics is a strange theory...
- *Today you might finally change your mind* 😊
- The question is how Alice can “**teleport**” the state of her Qbit $|\psi\rangle = a|0\rangle + b|1\rangle$ to Bob.

Quantum Teleportation

- If you are not yet convinced that quantum mechanics is a strange theory...
- *Today you might finally change your mind* 😊
- The question is how Alice can “**teleport**” the state of her Qbit $|\psi\rangle = a|0\rangle + b|1\rangle$ to Bob.
- What is the problem? One might wonder...

Quantum Teleportation

- If you are not yet convinced that quantum mechanics is a strange theory...
- *Today you might finally change your mind* 😊
- The question is how Alice can “**teleport**” the state of her Qbit $|\psi\rangle = a|0\rangle + b|1\rangle$ to Bob.
- What is the problem? One might wonder...
- Just send him a, b .

Quantum Teleportation

- If you are not yet convinced that quantum mechanics is a strange theory...
- *Today you might finally change your mind* 😊
- The question is how Alice can “**teleport**” the state of her Qbit $|\psi\rangle = a|0\rangle + b|1\rangle$ to Bob.
- What is the problem? One might wonder...
- Just send him a, b .
- But she doesn't know them...

Quantum Teleportation

- If you are not yet convinced that quantum mechanics is a strange theory...
- *Today you might finally change your mind* 😊
- The question is how Alice can “**teleport**” the state of her Qbit $|\psi\rangle = a|0\rangle + b|1\rangle$ to Bob.
- What is the problem? One might wonder...
- Just send him a, b .
- But she doesn't know them...
- ***And measurement, definitely doesn't help!***

Quantum Teleportation



Quantum Teleportation



Quantum Teleportation



$$|\psi\rangle = a|0\rangle + b|1\rangle$$



No Cloning Theorem

- Before we go into that, we will talk about the ***No-Cloning Theorem***.

No Cloning Theorem

- Before we go into that, we will talk about the ***No-Cloning Theorem***.
- ***You cannot clone a Qbit.***

No Cloning Theorem

- Before we go into that, we will talk about the ***No-Cloning Theorem***.
- ***You cannot clone a Qbit.***
- Well, if we could, then “teleporting” it to Bob would be a piece of cake.

No Cloning Theorem

- Before we go into that, we will talk about the *No-Cloning Theorem*.
- *You cannot clone a Qbit.*
- Well, if we could, then “teleporting” it to Bob would be a piece of cake.
- On the other hand, teleporting classical bit is easy. It’s called *One-Time Pad*.

No Cloning Theorem

- Before we go into that, we will talk about the *No-Cloning Theorem*.
- *You cannot clone a Qbit.*
- Well, if we could, then “teleporting” it to Bob would be a piece of cake.
- On the other hand, teleporting classical bit is easy. It’s called *One-Time Pad*.
- So, assume we have a single Qbit in some state $|\psi\rangle = a|0\rangle + b|1\rangle$.

No Cloning Theorem

- Before we go into that, we will talk about the *No-Cloning Theorem*.
- *You cannot clone a Qbit.*
- Well, if we could, then “teleporting” it to Bob would be a piece of cake.
- On the other hand, teleporting classical bit is easy. It’s called *One-Time Pad*.
- So, assume we have a single Qbit in some state $|\psi\rangle = a|0\rangle + b|1\rangle$.
- Suppose we do not know what these a, b really are

No Cloning Theorem

- Before we go into that, we will talk about the *No-Cloning Theorem*.
- *You cannot clone a Qbit.*
- Well, if we could, then “teleporting” it to Bob would be a piece of cake.
- On the other hand, teleporting classical bit is easy. It’s called *One-Time Pad*.
- So, assume we have a single Qbit in some state $|\psi\rangle = a|0\rangle + b|1\rangle$.
- Suppose we do not know what these a, b really are
- But you still want to *copy* it.

No Cloning Theorem

- So suppose that someone has another Qbit, for example in $|0\rangle$.

No Cloning Theorem


- So suppose that someone has another Qbit, for example in $|0\rangle$.
- What we want to do is to do *something* on the combined system consisting of $|\psi\rangle$ and $|0\rangle$

No Cloning Theorem

- So suppose that someone has another Qbit, for example in $|0\rangle$.
- What we want to do is to do *something* on the combined system consisting of $|\psi\rangle$ and $|0\rangle$ such that at the end both Qbits are in the state $|\psi\rangle$.

No Cloning Theorem

- So suppose that someone has another Qbit, for example in $|0\rangle$.
- What we want to do is to do *something* on the combined system consisting of $|\psi\rangle$ and $|0\rangle$ such that at the end both Qbits are in the state $|\psi\rangle$.



No Cloning Theorem

- The No Cloning Theorem says that there is no such unitary transformation U that can achieve this.

No Cloning Theorem

- The No Cloning Theorem says that there is no such unitary transformation U that can achieve this.
- Assume there was such U .

No Cloning Theorem

- The No Cloning Theorem says that there is no such unitary transformation U that can achieve this.
- Assume there was such U .
- Then

$$(a|0\rangle + b|1\rangle) \otimes |0\rangle \rightarrow (a|0\rangle + b|1\rangle) \otimes (a|0\rangle + b|1\rangle)$$

No Cloning Theorem

- The No Cloning Theorem says that there is no such unitary transformation U that can achieve this.
- Assume there was such U .

• Then

$$(a|0\rangle + b|1\rangle) \otimes |0\rangle \rightarrow (a|0\rangle + b|1\rangle) \otimes (a|0\rangle + b|1\rangle)$$

- *For all possible values of a, b that describe a valid quantum state.*

No Cloning Theorem

- The No Cloning Theorem says that there is no such unitary transformation U that can achieve this.
- Assume there was such U .

- Then

$$(a|0\rangle + b|1\rangle) \otimes |0\rangle \rightarrow (a|0\rangle + b|1\rangle) \otimes (a|0\rangle + b|1\rangle)$$

- *For all possible values of a, b that describe a valid quantum state.*
- Then it must also work when $a = 1, b = 0$ and when $a = 0, b = 1$

No Cloning Theorem

- The No Cloning Theorem says that there is no such unitary transformation U that can achieve this.
- Assume there was such U .

• Then

$$(a|0\rangle + b|1\rangle) \otimes |0\rangle \rightarrow (a|0\rangle + b|1\rangle) \otimes (a|0\rangle + b|1\rangle)$$

- *For all possible values of a, b that describe a valid quantum state.*
- Then it must also work when $a = 1, b = 0$ and when $a = 0, b = 1$
- Which means that under U we have $|00\rangle \rightarrow |00\rangle$ **and** $|10\rangle \rightarrow |11\rangle$

No Cloning Theorem

- Let's expand $(a|0\rangle + b|1\rangle) \otimes |0\rangle = a|00\rangle + b|10\rangle$

No Cloning Theorem

- Let's expand $(a|0\rangle + b|1\rangle) \otimes |0\rangle = a|00\rangle + b|10\rangle$
- According to the previous observation, this should be mapped to $a|00\rangle + b|11\rangle$.

No Cloning Theorem

- Let's expand $(a|0\rangle + b|1\rangle) \otimes |0\rangle = a|00\rangle + b|10\rangle$
- According to the previous observation, this should be mapped to $a|00\rangle + b|11\rangle$.
- Which, by the first equation, should be equal to $a^2|00\rangle + ab|01\rangle + ab|10\rangle + b^2|11\rangle$.

No Cloning Theorem

- Let's expand $(a|0\rangle + b|1\rangle) \otimes |0\rangle = a|00\rangle + b|10\rangle$
- According to the previous observation, this should be mapped to $a|00\rangle + b|11\rangle$.
- Which, by the first equation, should be equal to $a^2|00\rangle + ab|01\rangle + ab|10\rangle + b^2|11\rangle$.
- But the two expressions are equal only if $a = 1$ or $b = 1$, and $ab = 0$.

No Cloning Theorem

- Let's expand $(a|0\rangle + b|1\rangle) \otimes |0\rangle = a|00\rangle + b|10\rangle$
- According to the previous observation, this should be mapped to $a|00\rangle + b|11\rangle$.
- Which, by the first equation, should be equal to $a^2|00\rangle + ab|01\rangle + ab|10\rangle + b^2|11\rangle$.
- But the two expressions are equal only if $a = 1$ or $b = 1$, and $ab = 0$.
- Which is obviously a contradiction.

No Cloning Theorem

- Let's expand $(a|0\rangle + b|1\rangle) \otimes |0\rangle = a|00\rangle + b|10\rangle$
- According to the previous observation, this should be mapped to $a|00\rangle + b|11\rangle$.
- Which, by the first equation, should be equal to $a^2|00\rangle + ab|01\rangle + ab|10\rangle + b^2|11\rangle$.
- But the two expressions are equal only if $a = 1$ or $b = 1$, and $ab = 0$.
- Which is obviously a contradiction.
- \Rightarrow we cannot clone an unknown quantum state $|\psi\rangle$

The problem of Quantum Teleportation

- Alice wants to send to Bob $|\psi\rangle = a|0\rangle + b|1\rangle$

The problem of Quantum Teleportation

- Alice wants to send to Bob $|\psi\rangle = a|0\rangle + b|1\rangle$
- The amplitudes are of course *unknown* to Alice.

The problem of Quantum Teleportation

- Alice wants to send to Bob $|\psi\rangle = a|0\rangle + b|1\rangle$
- The amplitudes are of course *unknown* to Alice.
- Bob is far away and (crucially) they do not share a Quantum Channel...

The problem of Quantum Teleportation

- Alice wants to send to Bob $|\psi\rangle = a|0\rangle + b|1\rangle$
- The amplitudes are of course *unknown* to Alice.
- Bob is far away and (crucially) they do not share a Quantum Channel...
- Only a conventional telephone line.

The problem of Quantum Teleportation

- Alice wants to send to Bob $|\psi\rangle = a|0\rangle + b|1\rangle$
- The amplitudes are of course *unknown* to Alice.
- Bob is far away and (crucially) they do not share a Quantum Channel...
- Only a conventional telephone line.

***How can she possibly teleport $|\psi\rangle$ to Bob using only
Classical Information?***

The problem of Quantum Teleportation

- What they *can* do is share an entangled pair of Qbits $|\phi\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$.

The problem of Quantum Teleportation

- What they *can* do is share an entangled pair of Qbits $|\phi\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$.
- And, as you might imagine, they will heavily use their entangled state.

The problem of Quantum Teleportation

- What they *can* do is share an entangled pair of Qbits $|\phi\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$.
- And, as you might imagine, they will heavily use their entangled state.
- Indeed, Alice will perform some measurement on her unknown Qbit $|\psi\rangle$ and her part of the entangled state $|\phi\rangle$.

The problem of Quantum Teleportation

- What they *can* do is share an entangled pair of Qbits $|\phi\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$.
- And, as you might imagine, they will heavily use their entangled state.
- Indeed, Alice will perform some measurement on her unknown Qbit $|\psi\rangle$ and her part of the entangled state $|\phi\rangle$.
- The result of her measurement will be b_1, b_2 respectively.

The problem of Quantum Teleportation

- What they *can* do is share an entangled pair of Qbits $|\phi\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$.
- And, as you might imagine, they will heavily use their entangled state.
- Indeed, Alice will perform some measurement on her unknown Qbit $|\psi\rangle$ and her part of the entangled state $|\phi\rangle$.
- The result of her measurement will be b_1, b_2 respectively.
- She picks up her phone and tells Bob “Hey dude, I measured b_1, b_2 ”

The problem of Quantum Teleportation

- What they *can* do is share an entangled pair of Qbits $|\phi\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$.
- And, as you might imagine, they will heavily use their entangled state.
- Indeed, Alice will perform some measurement on her unknown Qbit $|\psi\rangle$ and her part of the entangled state $|\phi\rangle$.
- The result of her measurement will be b_1, b_2 respectively.
- She picks up her phone and tells Bob “Hey dude, I measured b_1, b_2 ”
- Bob will immediately deduce what a, b are...

The problem of Quantum Teleportation

- Technically, depending on the 4 possible values of $\mathbf{b}_1, \mathbf{b}_2$, *Bob performs one of possible four Unitary transformation on his part of the entangled pair of Qbits he shares with Alice.*

The problem of Quantum Teleportation

- Technically, depending on the 4 possible values of $\mathbf{b}_1, \mathbf{b}_2$, Bob performs one of possible four Unitary transformation on his part of the entangled pair of Qbits he shares with Alice.
- The output of this Unitary transformation is, amazingly, $|\psi\rangle = a|0\rangle + b|1\rangle$.

The problem of Quantum Teleportation

- Technically, depending on the 4 possible values of b_1, b_2 , Bob performs one of possible four Unitary transformation on his part of the entangled pair of Qbits he shares with Alice.
- The output of this Unitary transformation is, amazingly, $|\psi\rangle = a|0\rangle + b|1\rangle$.
- *Alice has successfully teleported her unknown Qbit to Bob.*

The problem of Quantum Teleportation

- Technically, depending on the 4 possible values of b_1, b_2 , Bob performs one of possible four Unitary transformation on his part of the entangled pair of Qbits he shares with Alice.
- The output of this Unitary transformation is, amazingly, $|\psi\rangle = a|0\rangle + b|1\rangle$.
- *Alice has successfully teleported her unknown Qbit to Bob.*
- Actually, she has teleported classical information that allows Bob to reconstruct $|\psi\rangle$.

The problem of Quantum Teleportation

- Technically, depending on the 4 possible values of b_1, b_2 , Bob performs one of possible four Unitary transformation on his part of the entangled pair of Qbits he shares with Alice.
- The output of this Unitary transformation is, amazingly, $|\psi\rangle = a|0\rangle + b|1\rangle$.
- *Alice has successfully teleported her unknown Qbit to Bob.*
- Actually, she has teleported classical information that allows Bob to reconstruct $|\psi\rangle$.
- In the process, she destroys her Qbit, but we do not care: we only care that she teleports it somehow to Bob.

First Approach

- Assume they have some means of ***Quantum communication***.

First Approach


- Assume they have some means of ***Quantum communication***.
- In particular, assume that they have a CNOT gate that extends from Alice's place, all the way to Bob's.

First Approach

- Assume they have some means of ***Quantum communication***.
- In particular, assume that they have a CNOT gate that extends from Alice's place, all the way to Bob's.
- So, Alice applies a CNOT to Bob's Qbit

First Approach

- Assume they have some means of ***Quantum communication***.
- In particular, assume that they have a CNOT gate that extends from Alice's place, all the way to Bob's.
- So, Alice applies a CNOT to Bob's Qbit



First Approach

- i.e. $\text{CNOT}(a|00\rangle + b|10\rangle) = a|00\rangle + b|11\rangle$

First Approach

- I.e. $\text{CNOT}(a|00\rangle + b|10\rangle) = a|00\rangle + b|11\rangle$
- So they end up in an entangled state (CNOT entangles the Qbits).

First Approach

- I.e. $\text{CNOT}(a|00\rangle + b|10\rangle) = a|00\rangle + b|11\rangle$
- So they end up in an entangled state (CNOT entangles the Qbits).
- Now, Alice measures her Qbit.

First Approach

- I.e. $\text{CNOT}(a|00\rangle + b|10\rangle) = a|00\rangle + b|11\rangle$
- So they end up in an entangled state (CNOT entangles the Qbits).
- Now, Alice measures her Qbit.
- *In which basis?*

First Approach

- I.e. $\text{CNOT}(a|00\rangle + b|10\rangle) = a|00\rangle + b|11\rangle$
- So they end up in an entangled state (CNOT entangles the Qbits).
- Now, Alice measures her Qbit.
- *In which basis?*
- Let's say she measures it in $|0\rangle, |1\rangle$ basis.

First Approach

- I.e. $\text{CNOT}(a|00\rangle + b|10\rangle) = a|00\rangle + b|11\rangle$
- So they end up in an entangled state (CNOT entangles the Qbits).
- Now, Alice measures her Qbit.
- *In which basis?*
- Let's say she measures it in $|0\rangle, |1\rangle$ basis.
- If she sees 0, the new state is $|00\rangle$.
- If she sees 1, the new state is $|11\rangle$.

First Approach

- I.e. $\text{CNOT}(a|00\rangle + b|10\rangle) = a|00\rangle + b|11\rangle$
- So they end up in an entangled state (CNOT entangles the Qbits).
- Now, Alice measures her Qbit.
- *In which basis?*
- Let's say she measures it in $|0\rangle, |1\rangle$ basis.
- If she sees 0, the new state is $|00\rangle$.
- If she sees 1, the new state is $|11\rangle$.
- Bob ends up with either state $|0\rangle$, or state $|1\rangle$, which have nothing to do with $|\psi\rangle = a|0\rangle + b|1\rangle$

First Approach

- I.e. $\text{CNOT}(a|00\rangle + b|10\rangle) = a|00\rangle + b|11\rangle$
- So they end up in an entangled state (CNOT entangles the Qbits).
- Now, Alice measures her Qbit.
- *In which basis?*
- Let's say she measures it in $|0\rangle, |1\rangle$ basis.
- If she sees 0, the new state is $|00\rangle$.
- If she sees 1, the new state is $|11\rangle$.
- Bob ends up with either state $|0\rangle$, or state $|1\rangle$, which have nothing to do with $|\psi\rangle = a|0\rangle + b|1\rangle$ 😞

First Approach

- Here is a better idea.

First Approach

- Here is a better idea.
- Alice measures in the $|+\rangle, |-\rangle$ basis.

First Approach

- Here is a better idea.
- Alice measures in the $|+\rangle, |-\rangle$ basis.
- We need to rewrite Alice's Qbit in the $+, -$ basis:

First Approach

- Here is a better idea.
- Alice measures in the $|+\rangle, |-\rangle$ basis.
- We need to rewrite Alice's Qbit in the $+, -$ basis:

$$a|00\rangle + b|11\rangle = a\left(\frac{1}{\sqrt{2}}|+\rangle + \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |0\rangle + b\left(\frac{1}{\sqrt{2}}|+\rangle - \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |1\rangle$$

First Approach

- Here is a better idea.
- Alice measures in the $|+\rangle, |-\rangle$ basis.
- We need to rewrite Alice's Qbit in the $+, -$ basis:

$$a|00\rangle + b|11\rangle = a\left(\frac{1}{\sqrt{2}}|+\rangle + \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |0\rangle + b\left(\frac{1}{\sqrt{2}}|+\rangle - \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |1\rangle$$

First Approach

- Here is a better idea.
- Alice measures in the $|+\rangle, |-\rangle$ basis.
- We need to rewrite Alice's Qbit in the $+, -$ basis:

$$a|00\rangle + b|11\rangle = a\left(\frac{1}{\sqrt{2}}|+\rangle + \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |0\rangle + b\left(\frac{1}{\sqrt{2}}|+\rangle - \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |1\rangle$$

First Approach

- Here is a better idea.
- Alice measures in the $|+\rangle, |-\rangle$ basis.
- We need to rewrite Alice's Qbit in the $+, -$ basis:

$$a|00\rangle + b|11\rangle = a\left(\frac{1}{\sqrt{2}}|+\rangle + \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |0\rangle + b\left(\frac{1}{\sqrt{2}}|+\rangle - \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |1\rangle$$

- Now we make the calculations...

First Approach

- Here is a better idea.
- Alice measures in the $|+\rangle, |-\rangle$ basis.
- We need to rewrite Alice's Qbit in the $+, -$ basis:

$$a|00\rangle + b|11\rangle = a\left(\frac{1}{\sqrt{2}}|+\rangle + \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |0\rangle + b\left(\frac{1}{\sqrt{2}}|+\rangle - \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |1\rangle$$

- Now we make the calculations...

$$a|00\rangle + b|11\rangle = \frac{1}{\sqrt{2}}|+\rangle[a|0\rangle + b|1\rangle] + \frac{1}{\sqrt{2}}|-\rangle[a|0\rangle - b|1\rangle]$$

First Approach

- Here is a better idea.
- Alice measures in the $|+\rangle, |-\rangle$ basis.
- We need to rewrite Alice's Qbit in the $+, -$ basis:

$$a|00\rangle + b|11\rangle = a\left(\frac{1}{\sqrt{2}}|+\rangle + \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |0\rangle + b\left(\frac{1}{\sqrt{2}}|+\rangle - \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |1\rangle$$

- Now we make the calculations...

$$a|00\rangle + b|11\rangle = \frac{1}{\sqrt{2}}|+\rangle[a|0\rangle + b|1\rangle] + \frac{1}{\sqrt{2}}|-\rangle[a|0\rangle - b|1\rangle]$$

- *This is much better!*

First Approach

- Here is a better idea.
- Alice measures in the $|+\rangle, |-\rangle$ basis.
- We need to rewrite Alice's Qbit in the $+, -$ basis:

$$a|00\rangle + b|11\rangle = a\left(\frac{1}{\sqrt{2}}|+\rangle + \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |0\rangle + b\left(\frac{1}{\sqrt{2}}|+\rangle - \frac{1}{\sqrt{2}}|-\rangle\right) \otimes |1\rangle$$

- Now we make the calculations...

$$a|00\rangle + b|11\rangle = \frac{1}{\sqrt{2}}|+\rangle[a|0\rangle + b|1\rangle] + \frac{1}{\sqrt{2}}|-\rangle[a|0\rangle - b|1\rangle]$$

- ***This is much better!***

- If Alice measures $+$ (i.e., 0), Bob's bit is in $|\psi\rangle$!

First Approach

- But what if Alice measures — (i.e., 1)?

First Approach

- But what if Alice measures — (i.e., 1)?
- Then Bob knows his Qbit is in state $a|0\rangle - b|1\rangle$

First Approach

- But what if Alice measures — (i.e., 1)?
- Then Bob knows his Qbit is in state $a|0\rangle - b|1\rangle$
- Apply the **Z**-Unitary transform to his Qbit!

First Approach

- But what if Alice measures — (i.e., 1)?
- Then Bob knows his Qbit is in state $a|0\rangle - b|1\rangle$
- Apply the **Z**-Unitary transform to his Qbit!
- $Z(a|0\rangle - b|1\rangle) = a|0\rangle + b|1\rangle$

First Approach

- But what if Alice measures — (i.e., 1)?
- Then Bob knows his Qbit is in state $a|0\rangle - b|1\rangle$
- Apply the **Z**-Unitary transform to his Qbit!
- $Z(a|0\rangle - b|1\rangle) = a|0\rangle + b|1\rangle$
- And that is it!

First Approach: The algorithm

- Alice has $|\psi\rangle = a|0\rangle + b|1\rangle$. Bob has $|0\rangle$.
- They share a CNOT gate with which they entangle their Qbits in the composite state $|\psi'\rangle = a|00\rangle + b|11\rangle$.
- Alice measures her Qbit in the $|+\rangle, |-\rangle$ basis.
- She calls and tells Bob if she observed 0 or 1.
- If she observed 0, Bob's Qbit is in $|\psi\rangle$.
- If she observes 1, Bob apply Z to his Qbit and now this is again in $|\psi\rangle$.

First Approach: The algorithm


But how they can create their entangled state $|\psi'\rangle = a|00\rangle + b|11\rangle$ without any means of Quantum communications (i.e., the CNOT gate) between them?

Exercise to think about (from last lecture)

- Construct a circuit that given two Qbits, both prepared in the $|0\rangle$ state, outputs the Bell state $\frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$


Exercise to think about (from last lecture)

- Construct a circuit that given two Qbits, both prepared in the $|0\rangle$ state, outputs the Bell state $\frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle = |\Phi^+\rangle$




Exercise to think about (from last lecture)

- What happens if we apply the same circuit to input bits $|0\rangle, |1\rangle$?




Exercise to think about (from last lecture)

- What happens if we apply the same circuit to input bits $|0\rangle, |1\rangle$?
- The output is $\frac{1}{\sqrt{2}}|01\rangle + \frac{1}{\sqrt{2}}|10\rangle = |\Psi^+\rangle$.




Exercise to think about (from last lecture)

- What happens if we apply the same circuit to input bits $|1\rangle, |0\rangle$?




Exercise to think about (from last lecture)

- What happens if we apply the same circuit to input bits $|1\rangle, |0\rangle$?
- The output is $\frac{1}{\sqrt{2}}|00\rangle - \frac{1}{\sqrt{2}}|11\rangle = |\Phi^-\rangle$




Exercise to think about (from last lecture)

- What happens if we apply the same circuit to input bits $|1\rangle, |1\rangle$?



Exercise to think about (from last lecture)

- What happens if we apply the same circuit to input bits $|\mathbf{1}\rangle, |\mathbf{1}\rangle$?
- The output is $\frac{1}{\sqrt{2}} |\mathbf{01}\rangle - \frac{1}{\sqrt{2}} |\mathbf{10}\rangle = |\Psi^-\rangle$.



Exercise to think about (from last lecture)

- All these four states are called **Bell states**.

Exercise to think about (from last lecture)

- All these four states are called **Bell states**.
- They form an ***orthonormal*** basis for the 4-dimensional vector space describing composite systems of two Qbits

Second approach: no Quantum channel

- Now, Alice and Bob will try to create the state $|\psi'\rangle = a|00\rangle + b|11\rangle$ but without using a Quantum communication channel between them.

Second approach: no Quantum channel

- Now, Alice and Bob will try to create the state $|\psi'\rangle = a|00\rangle + b|11\rangle$ but without using a Quantum communication channel between them.
- Only telephone lines!

Second approach: no Quantum channel

- Now, Alice and Bob will try to create the state $|\psi'\rangle = a|00\rangle + b|11\rangle$ but without using a Quantum communication channel between them.
- Only telephone lines!
- But in their rooms, they can do whatever they want.

Second approach: no Quantum channel

- Now, Alice and Bob will try to create the state $|\psi'\rangle = a|00\rangle + b|11\rangle$ but without using a Quantum communication channel between them.
- Only telephone lines!
- But in their rooms, they can do whatever they want.
- What they can do is meet beforehand in a secret location and share a Bell's state $|\Phi^+\rangle$.

Second approach: no Quantum channel

- Now, Alice and Bob will try to create the state $|\psi'\rangle = a|00\rangle + b|11\rangle$ but without using a Quantum communication channel between them.
- Only telephone lines!
- But in their rooms, they can do whatever they want.
- What they can do is meet beforehand in a secret location and share a Bell's state $|\Phi^+\rangle$.
- **Then they move far apart.**

Second approach: no Quantum channel

- Now, Alice and Bob will try to create the state $|\psi'\rangle = a|00\rangle + b|11\rangle$ but without using a Quantum communication channel between them.
- Only telephone lines!
- But in their rooms, they can do whatever they want.
- What they can do is meet beforehand in a secret location and share a Bell's state $|\Phi^+\rangle$.
- **Then they move far apart.**
- Alice applies a CNOT to her Qbit $|\psi\rangle$ and her Qbit from the entangled state $|\Phi^+\rangle$.

Second approach: no Quantum channel

- Now, Alice and Bob will try to create the state $|\psi'\rangle = a|00\rangle + b|11\rangle$ but without using a Quantum communication channel between them.
- Only telephone lines!
- But in their rooms, they can do whatever they want.
- What they can do is meet beforehand in a secret location and share a Bell's state $|\Phi^+\rangle$.
- **Then they move far apart.**
- Alice applies a CNOT to her Qbit $|\psi\rangle$ and her Qbit from the entangled state $|\Phi^+\rangle$.
- No problem with that, she can do it.

Second approach: no Quantum channel

- What is the composite state describing Alice's and Bob's Qbits?

Second approach: no Quantum channel

- What is the composite state describing Alice's and Bob's Qbits?

$$(a|0\rangle + b|1\rangle) \otimes |\Phi^+\rangle =$$

Second approach: no Quantum channel

- What is the composite state describing Alice's and Bobs Qbits?

$$\begin{aligned} & (a|0\rangle + b|1\rangle) \otimes |\Phi^+\rangle = \\ &= \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle + \frac{b}{\sqrt{2}}|100\rangle + \frac{b}{\sqrt{2}}|111\rangle = |C\rangle \end{aligned}$$

Second approach: no Quantum channel

- What is the composite state describing Alice's and Bob's Qbits?

$$\begin{aligned} & (a|0\rangle + b|1\rangle) \otimes |\Phi^+\rangle = \\ &= \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle + \frac{b}{\sqrt{2}}|100\rangle + \frac{b}{\sqrt{2}}|111\rangle = |C\rangle \end{aligned}$$

- Alice applies CNOT to her two Qbits: the control bit is $|\psi\rangle$.

Second approach: no Quantum channel

- What is the composite state describing Alice's and Bob's Qbits?

$$\begin{aligned} & (a|0\rangle + b|1\rangle) \otimes |\Phi^+\rangle = \\ &= \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle + \frac{b}{\sqrt{2}}|100\rangle + \frac{b}{\sqrt{2}}|111\rangle = |C\rangle \end{aligned}$$

- Alice applies CNOT to her two Qbits: the control bit is $|\psi\rangle$.
- So, $|C\rangle$ becomes

Second approach: no Quantum channel

- What is the composite state describing Alice's and Bob's Qbits?

$$\begin{aligned} & (a|0\rangle + b|1\rangle) \otimes |\Phi^+\rangle = \\ &= \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle + \frac{b}{\sqrt{2}}|100\rangle + \frac{b}{\sqrt{2}}|111\rangle = |C\rangle \end{aligned}$$

- Alice applies CNOT to her two Qbits: the control bit is $|\psi\rangle$.
- So, $|C\rangle$ becomes

$$|C'\rangle = \frac{a}{\sqrt{2}}|000\rangle$$

Second approach: no Quantum channel

- What is the composite state describing Alice's and Bobs Qbits?

$$\begin{aligned} & (a|0\rangle + b|1\rangle) \otimes |\Phi^+\rangle = \\ &= \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle + \frac{b}{\sqrt{2}}|100\rangle + \frac{b}{\sqrt{2}}|111\rangle = |C\rangle \end{aligned}$$

- Alice applies CNOT to her two Qbits: the control bit is $|\psi\rangle$.
- So, $|C\rangle$ becomes

$$|C'\rangle = \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle$$

Second approach: no Quantum channel

- What is the composite state describing Alice's and Bobs Qbits?

$$\begin{aligned} & (a|0\rangle + b|1\rangle) \otimes |\Phi^+\rangle = \\ &= \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle + \frac{b}{\sqrt{2}}|100\rangle + \frac{b}{\sqrt{2}}|111\rangle = |C\rangle \end{aligned}$$

- Alice applies CNOT to her two Qbits: the control bit is $|\psi\rangle$.
- So, $|C\rangle$ becomes

$$|C'\rangle = \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle + \frac{b}{\sqrt{2}}|110\rangle$$

Second approach: no Quantum channel

- What is the composite state describing Alice's and Bob's Qbits?

$$\begin{aligned} & (a|0\rangle + b|1\rangle) \otimes |\Phi^+\rangle = \\ &= \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle + \frac{b}{\sqrt{2}}|100\rangle + \frac{b}{\sqrt{2}}|111\rangle = |C\rangle \end{aligned}$$

- Alice applies CNOT to her two Qbits: the control bit is $|\psi\rangle$.
- So, $|C\rangle$ becomes

$$|C'\rangle = \frac{a}{\sqrt{2}}|000\rangle + \frac{a}{\sqrt{2}}|011\rangle + \frac{b}{\sqrt{2}}|110\rangle + \frac{b}{\sqrt{2}}|101\rangle$$

Second approach: no Quantum channel

- What happens if Alice measures her second Qbit (the one from $|\Phi^+\rangle$)?

Second approach: no Quantum channel

- What happens if Alice measures her second Qbit (the one from $|\Phi^+\rangle$)?
- With probability $\frac{1}{2}$ she sees 0 and with probability $\frac{1}{2}$ she sees 1.

Second approach: no Quantum channel

- What happens if Alice measures her second Qbit (the one from $|\Phi^+\rangle$)?
- With probability $\frac{1}{2}$ she sees 0 and with probability $\frac{1}{2}$ she sees 1.
- If she sees 0, the new state becomes

$$|C'\rangle = a|00\rangle + b|11\rangle$$

Second approach: no Quantum channel

- What happens if Alice measures her second Qbit (the one from $|\Phi^+\rangle$)?
- With probability $\frac{1}{2}$ she sees 0 and with probability $\frac{1}{2}$ she sees 1.
- If she sees 0, the new state becomes
$$|C'\rangle = a|00\rangle + b|11\rangle$$
- If she sees 1, the new state becomes
$$|C'\rangle = a|01\rangle + b|10\rangle$$

Second approach: no Quantum channel

- What happens if Alice measures the second Qbit (the one from $|\Phi^+\rangle$)?
- With probability $\frac{1}{2}$ she sees 0 and with probability $\frac{1}{2}$ she sees 1.
- If she sees 0, the new state becomes
$$|C'\rangle = a|00\rangle + b|11\rangle$$
- If she sees 1, the new state becomes
$$|C'\rangle = a|01\rangle + b|10\rangle$$
- She picks up the phone and tells Bob if she got 0 or 1.

Second approach: no Quantum channel

- What happens if Alice measures the second Qbit (the one from $|\Phi^+\rangle$)?
- With probability $\frac{1}{2}$ she sees 0 and with probability $\frac{1}{2}$ she sees 1.
- If she sees 0, the new state becomes
$$|C'\rangle = a|00\rangle + b|11\rangle$$
- If she sees 1, the new state becomes
$$|C'\rangle = a|01\rangle + b|10\rangle$$
- She picks up the phone and tells Bob if she got 0 or 1.
- If she gets 0, Bob leaves his Qbit alone.

Second approach: no Quantum channel

- What happens if Alice measures the second Qbit (the one from $|\Phi^+\rangle$)?
- With probability $\frac{1}{2}$ she sees 0 and with probability $\frac{1}{2}$ she sees 1.
- If she sees 0, the new state becomes
$$|C'\rangle = a|00\rangle + b|11\rangle$$
- If she sees 1, the new state becomes
$$|C'\rangle = a|01\rangle + b|10\rangle$$
- She picks up the phone and tells Bob if she got 0 or 1.
- If she gets 0, Bob leaves his Qbit alone. *They share the desired state!*

Second approach: no Quantum channel

- What happens if Alice measures the second Qbit (the one from $|\Phi^+\rangle$)?
- With probability $\frac{1}{2}$ she sees 0 and with probability $\frac{1}{2}$ she sees 1.
- If she sees 0, the new state becomes
$$|C'\rangle = a|00\rangle + b|11\rangle$$
- If she sees 1, the new state becomes
$$|C'\rangle = a|01\rangle + b|10\rangle$$
- She picks up the phone and tells Bob if she got 0 or 1.
- If she gets 0, Bob leaves his Qbit alone. *They share the desired state!*
- Else, Bob applies a bit-flip transform X to his bit to get $|C'\rangle = a|00\rangle + b|11\rangle$.

Second approach: no Quantum channel

- What happens if Alice measures the second Qbit (the one from $|\Phi^+\rangle$)?
- With probability $\frac{1}{2}$ she sees 0 and with probability $\frac{1}{2}$ she sees 1.
- If she sees 0, the new state becomes
$$|C'\rangle = a|00\rangle + b|11\rangle$$
- If she sees 1, the new state becomes
$$|C'\rangle = a|01\rangle + b|10\rangle$$
- She picks up the phone and tells Bob if she got 0 or 1.
- If she gets 0, Bob leaves his Qbit alone. *They share the desired state!*
- Else, Bob applies a bit-flip transform X to his bit to get $|C'\rangle = a|00\rangle + b|11\rangle$. *Again they share the desired state.*

Exercise to think about

- Construct the circuit for the Quantum Teleportation protocol.