


Intro to Quantum Computing


Maths of Qbits Part III



Today

- A *very gentle* intro to Postulates (axioms) of Quantum Mechanics
- At least through the lenses of Qbits
- Intro to Operators
- Projection operators, Pauli matrices. The Hadamard matrix

Today



Quantum Mechanics

Quantum Mechanics

- QM is not a physical theory.

Quantum Mechanics

- QM is not a physical theory.
- It is a framework for the development of physical theories.

Quantum Mechanics

- QM is not a physical theory.
- It is a framework for the development of physical theories.
- In CS vocabulary:

Quantum Mechanics

- QM is not a physical theory.
- It is a framework for the development of physical theories.
- In CS vocabulary:

QM is the operating system that all physical theories run as application software

Quantum Mechanics

- With one exception!

Quantum Mechanics

- With one exception!
- *Gravity* has not (yet) been ported into the QM Operating System

Quantum Mechanics

- With one exception!
- *Gravity* has not (yet) been ported into the QM Operating System
- And it is unknown if it will ever be.


Quantum Mechanics

- With one exception!
- *Gravity* has not (yet) been ported into the QM Operating System
- And it is unknown if it will ever be.
- One example of successful portability is *Quantum Electrodynamics*

Quantum Mechanics

- With one exception!
- **Gravity** has not (yet) been ported into the QM Operating System
- And it is unknown if it will ever be.
- One example of successful portability is **Quantum Electrodynamics**
- Richard Feynman: [this theory is] *“The jewel of physics”*

Quantum Mechanics



The four postulates of QM

The four postulates of QM

1. *State space*: how a quantum state can be described?

The four postulates of QM

1. **State space**: how a quantum state can be described?
2. **Evolution**: how a quantum state is allowed to change over time?

The four postulates of QM

1. **State space**: how a quantum state can be described?
2. **Evolution**: how a quantum state is allowed to change over time?
3. **Measurement**: the effect on a quantum state of interaction with a classical system (apparatus) that yields *classical* information.

The four postulates of QM

1. **State space**: how a quantum state can be described?
2. **Evolution**: how a quantum state is allowed to change over time?
3. **Measurement**: the effect on a quantum state of interaction with a classical system (apparatus) that yields *classical* information.
4. **Composition**: How quantum systems can be composed?

State space of Quantum System

- *Postulate 1*

Associated to any isolated physical system is a (complex) vector space equipped with an inner product (i.e., a Hilbert space) known as the state space of the system. The system is completely described by its state vector, which is a unit vector in the system's state space.

State space of Quantum System

- *Postulate 1*

Associated to any isolated physical system is a (complex) vector space equipped with an inner product (i.e., a Hilbert space) known as the state space of the system. The system is completely described by its state vector, which is a unit vector in the system's state space.

- We are interested in Qbits.

State space of Quantum System

- **Postulate 1**

Associated to any isolated physical system is a (complex) vector space equipped with an inner product (i.e., a Hilbert space) known as the state space of the system. The system is completely described by its state vector, which is a unit vector in the system's state space.

- We are interested in Qbits.
- Their associated state space is R^2 (or C^2 in general).

State space of Quantum System

- *Postulate 1*

Associated to any isolated physical system is a (complex) vector space equipped with an inner product (i.e., a Hilbert space) known as the state space of the system. The system is completely described by its state vector, which is a unit vector in the system's state space.

- We are interested in Qbits.
- Their associated state space is R^2 (or C^2 in general).
- There are generalization (qtrits etc.)

State space of Quantum System

- *Postulate 1*

Associated to any isolated physical system is a (complex) vector space equipped with an inner product (i.e., a Hilbert space) known as the state space of the system. The system is completely described by its state vector, which is a unit vector in the system's state space.

- We are interested in Qbits.
- Their associated state space is R^2 (or C^2 in general).
- There are generalization (qtrits etc.)
- We have discussed physical realizations.

Evolution of a Quantum System

- **Postulate 2**

The evolution over time of a quantum system is described by Schrödinger equation

$$i\hbar \frac{d|\psi\rangle}{dt} = \mathbf{H}|\psi\rangle$$

Where \hbar is Planck's constant and \mathbf{H} is a Hermitian operator (Hamiltonian)

Evolution of a Quantum System

- *Postulate 2*

The evolution over time of a quantum system is described by Schrödinger equation

$$i\hbar \frac{d|\psi\rangle}{dt} = \mathbf{H}|\psi\rangle$$

Where \hbar is Planck's constant and \mathbf{H} is a Hermitian operator (Hamiltonian)

- That's great, but this concerns (mostly) continuous evolution of time.

Evolution of a Quantum System

- *Postulate 2*

The evolution over time of a quantum system is described by Schrödinger equation

$$i\hbar \frac{d|\psi\rangle}{dt} = \mathbf{H}|\psi\rangle$$

Where \hbar is Planck's constant and \mathbf{H} is a Hermitian operator (Hamiltonian)

- That's great, but this concerns (mostly) continuous evolution of time.
- In Computer Science we are interested in discrete time intervals.

Evolution of a Quantum System

- *Postulate 2*

The evolution over time of a quantum system is described by Schrödinger equation

$$i\hbar \frac{d|\psi\rangle}{dt} = \mathbf{H}|\psi\rangle$$

Where \hbar is Planck's constant and \mathbf{H} is a Hermitian operator (Hamiltonian)

- That's great, but this concerns (mostly) continuous evolution of time.
- In Computer Science we are interested in discrete time intervals.
- Thus...

Evolution of a Quantum System

- *Postulate 2'*

The evolution over time of a quantum system from time t_1 to the next time step t_2 is described by

$$|\psi_{t_2}\rangle = \mathbf{U}|\psi_{t_1}\rangle$$

Where \mathbf{U} is a unitary transformation

Evolution of a Quantum System

- *Postulate 2'*

The evolution over time of a quantum system from time t_1 to the next time step t_2 is described by

$$|\psi_{t_2}\rangle = \mathbf{U}|\psi_{t_1}\rangle$$

Where \mathbf{U} is a unitary transformation

- It follows directly by previous equation

Evolution of a Quantum System

- *Postulate 2'*

The evolution over time of a quantum system from time t_1 to the next time step t_2 is described by

$$|\psi_{t_2}\rangle = \mathbf{U}|\psi_{t_1}\rangle$$

Where \mathbf{U} is a unitary transformation

- It follows directly by previous equation
- Some authors in Quantum Computing call Postulate 2' as the *Fundamental Postulate of Quantum state evolution*

Why Unitary Matrices?

Why Unitary Matrices?

- A unitary matrix is the answer to the following question:

Why Unitary Matrices?

- A unitary matrix is the answer to the following question:

What is the most general type of matrix that maps a unit vector in ℓ_2 norm to another unit vector in ℓ_2 norm?

Why Unitary Matrices?

- A unitary matrix is the answer to the following question:

What is the most general type of matrix that maps a unit vector in ℓ_2 norm to another unit vector in ℓ_2 norm?

- We want to transform $|\psi_1\rangle = \alpha|0\rangle + \beta|1\rangle$ to $|\psi_2\rangle = \gamma|0\rangle + \delta|1\rangle$.

Why Unitary Matrices?

- A unitary matrix is the answer to the following question:

What is the most general type of matrix that maps a unit vector in ℓ_2 norm to another unit vector in ℓ_2 norm?

- We want to transform $|\psi_1\rangle = \alpha|0\rangle + \beta|1\rangle$ to $|\psi_2\rangle = \gamma|0\rangle + \delta|1\rangle$.
- The transformation is $|\psi_2\rangle = U|\psi_1\rangle$.

Why Unitary Matrices?

- A unitary matrix is the answer to the following question:

What is the most general type of matrix that maps a unit vector in ℓ_2 norm to another unit vector in ℓ_2 norm?

- We want to transform $|\psi_1\rangle = \alpha|0\rangle + \beta|1\rangle$ to $|\psi_2\rangle = \gamma|0\rangle + \delta|1\rangle$.
- The transformation is $|\psi_2\rangle = U|\psi_1\rangle$.
- The amplitudes of $|\psi_2\rangle$ must have the property that $\gamma^2 + \delta^2 = 1$.

Why Unitary Matrices?

- A unitary matrix is the answer to the following question:

What is the most general type of matrix that maps a unit vector in ℓ_2 norm to another unit vector in ℓ_2 norm?

- In other words

Why Unitary Matrices?

- A unitary matrix is the answer to the following question:

What is the most general type of matrix that maps a unit vector in ℓ_2 norm to another unit vector in ℓ_2 norm?

- In other words

$$\begin{bmatrix} \gamma \\ \delta \end{bmatrix} = \mathbf{U} \begin{bmatrix} \alpha \\ \beta \end{bmatrix}, \text{ where } \begin{bmatrix} \gamma \\ \delta \end{bmatrix} = |\psi_2\rangle \text{ and } \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = |\psi_1\rangle \text{ such that}$$

$$\left\| |\psi_2\rangle \right\|_2 = \left\| |\psi_1\rangle \right\|_2 = 1 \text{ (unit length vectors in } \ell_2 \text{)}$$

What is a unitary matrix?

What is a unitary matrix?

- In the real space, a unitary matrix M is a *square* matrix whose inverse equal its transpose

$$M^{-1} = M^T$$

What is a unitary matrix?

- In the real space, a unitary matrix \mathbf{M} is a **square** matrix whose inverse equal its transpose

$$\mathbf{M}^{-1} = \mathbf{M}^T$$

- In the complex space a square matrix \mathbf{M} is a unitary matrix if its inverse is its conjugate transpose

$$\mathbf{M}^{-1} = \mathbf{M}^\dagger$$

What is a unitary matrix?

- In the real space, a unitary matrix \mathbf{M} is a **square** matrix whose inverse equal its transpose

$$\mathbf{M}^{-1} = \mathbf{M}^T$$

- In the complex space a square matrix \mathbf{M} is a unitary matrix if its inverse is its conjugate transpose

$$\mathbf{M}^{-1} = \mathbf{M}^\dagger$$

- It follows that $\mathbf{M}^T \mathbf{M} = \mathbf{M}^{-1} \mathbf{M} = \mathbf{I}$.

What is a unitary matrix?

- In the real space, a unitary matrix \mathbf{M} is a **square** matrix whose inverse equal its transpose

$$\mathbf{M}^{-1} = \mathbf{M}^T$$

- In the complex space a square matrix \mathbf{M} is a unitary matrix if its inverse is its conjugate transpose

$$\mathbf{M}^{-1} = \mathbf{M}^\dagger$$

- It follows that $\mathbf{M}^T \mathbf{M} = \mathbf{M}^{-1} \mathbf{M} = \mathbf{I}$.
- All unitary matrices are orthogonal and even diagonalizable.

What is a unitary matrix?

- In the real space, a unitary matrix \mathbf{M} is a **square** matrix whose inverse equal its transpose

$$\mathbf{M}^{-1} = \mathbf{M}^T$$

- In the complex space a square matrix \mathbf{M} is a unitary matrix if its inverse is its conjugate transpose

$$\mathbf{M}^{-1} = \mathbf{M}^\dagger$$

- It follows that $\mathbf{M}^T \mathbf{M} = \mathbf{M}^{-1} \mathbf{M} = \mathbf{I}$.
- All unitary matrices are orthogonal and even diagonalizable.
- All eigenvalues of unitary matrices have absolute value one.

What is a unitary matrix?

- Most importantly: *They preserve inner products:*

What is a unitary matrix?

- Most importantly: *They preserve inner products:*
- Let $\mathbf{M}|\mathbf{u}\rangle = |\mathbf{u}'\rangle$ and $\mathbf{M}|\mathbf{v}\rangle = |\mathbf{v}'\rangle$. Then we have that

What is a unitary matrix?

- Most importantly: *They preserve inner products:*
- Let $M|u\rangle = |u'\rangle$ and $M|v\rangle = |v'\rangle$. Then we have that

$$\langle u'|v'\rangle = (M|u\rangle)^T (M|v\rangle)$$

What is a unitary matrix?

- Most importantly: *They preserve inner products:*
- Let $M|u\rangle = |u'\rangle$ and $M|v\rangle = |v'\rangle$. Then we have that

$$\begin{aligned}\langle u'|v'\rangle &= (M|u\rangle)^T (M|v\rangle) \\ &= (\langle u|M^T)(M|v\rangle)\end{aligned}$$

What is a unitary matrix?

- Most importantly: *They preserve inner products:*
- Let $M|u\rangle = |u'\rangle$ and $M|v\rangle = |v'\rangle$. Then we have that

$$\begin{aligned}\langle u'|v'\rangle &= (M|u\rangle)^T (M|v\rangle) \\ &= (\langle u|M^T)(M|v\rangle) \\ &= \langle u|M^T M|v\rangle\end{aligned}$$

What is a unitary matrix?

- Most importantly: *They preserve inner products:*
- Let $M|u\rangle = |u'\rangle$ and $M|v\rangle = |v'\rangle$. Then we have that

$$\begin{aligned}\langle u'|v'\rangle &= (M|u\rangle)^T (M|v\rangle) \\ &= (\langle u|M^T)(M|v\rangle) \\ &= \langle u|M^T M|v\rangle \\ &= \langle u|I|v\rangle\end{aligned}$$

What is a unitary matrix?

- Most importantly: *They preserve inner products:*
- Let $M|u\rangle = |u'\rangle$ and $M|v\rangle = |v'\rangle$. Then we have that

$$\begin{aligned}\langle u'|v'\rangle &= (M|u\rangle)^T (M|v\rangle) \\ &= (\langle u|M^T)(M|v\rangle) \\ &= \langle u|M^T M|v\rangle \\ &= \langle u|I|v\rangle \\ &= \langle u|v\rangle\end{aligned}$$

Unitary matrices: Example

- Consider the following matrix

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = U$$

Unitary matrices: Example

- Consider the following matrix

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = \mathbf{U} \text{ and apply it to } |\mathbf{0}\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Unitary matrices: Example

- Consider the following matrix

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = \mathbf{U} \text{ and apply it to } |\mathbf{0}\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ to get } |\boldsymbol{\phi}\rangle = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ 1 \end{bmatrix}$$

Unitary matrices: Example

- Consider the following matrix

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = \mathbf{U} \text{ and apply it to } |\mathbf{0}\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ to get } |\boldsymbol{\phi}\rangle = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} \text{, i.e., a coin flip!}$$

Unitary matrices: Example

- Consider the following matrix

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = \mathbf{U} \text{ and apply it to } |\mathbf{0}\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ to get } |\boldsymbol{\phi}\rangle = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ 1 \end{bmatrix} \text{ i.e., a coin flip!}$$

- Now apply \mathbf{U} to $|\boldsymbol{\phi}\rangle$

Unitary matrices: Example

- Consider the following matrix

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = \mathbf{U} \text{ and apply it to } |\mathbf{0}\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ to get } |\boldsymbol{\phi}\rangle = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ 1 \end{bmatrix} \text{, i.e., a coin flip!}$$

- Now apply \mathbf{U} to $|\boldsymbol{\phi}\rangle$ to get $\mathbf{U}|\boldsymbol{\psi}\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix} = |\mathbf{1}\rangle$

Unitary matrices: Example

- Consider the following matrix

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = \mathbf{U} \text{ and apply it to } |\mathbf{0}\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ to get } |\boldsymbol{\phi}\rangle = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ 1 \end{bmatrix} \text{ i.e., a coin flip!}$$

- Now apply \mathbf{U} to $|\boldsymbol{\phi}\rangle$ to get $\mathbf{U}|\boldsymbol{\psi}\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix} = |\mathbf{1}\rangle$ i.e., a deterministic outcome!

Unitary matrices: Example

- Consider the following matrix

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = \mathbf{U} \text{ and apply it to } |\mathbf{0}\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ to get } |\boldsymbol{\phi}\rangle = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ 1 \end{bmatrix} \text{ i.e., a coin flip!}$$

- Now apply \mathbf{U} to $|\boldsymbol{\phi}\rangle$ to get $\mathbf{U}|\boldsymbol{\psi}\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix} = |\mathbf{1}\rangle$ i.e., a deterministic outcome!
- This is an example of *Quantum Interference*
-

Unitary matrices: Example

- Consider the following matrix

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = \mathbf{U} \text{ and apply it to } |\mathbf{0}\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ to get } |\boldsymbol{\phi}\rangle = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ 1 \end{bmatrix} \text{ i.e., a coin flip!}$$

- Now apply \mathbf{U} to $|\boldsymbol{\phi}\rangle$ to get $\mathbf{U}|\boldsymbol{\psi}\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix} = |\mathbf{1}\rangle$ i.e., a deterministic outcome!
- This is an example of ***Quantum Interference***
- The two “paths” leading to $|\mathbf{0}\rangle$ interfere destructively (cancel out).

Measurement, Projection and Collapse

- Let $|\psi\rangle = a|0\rangle + b|1\rangle$ be the state of a qbit.

Measurement, Projection and Collapse

- Let $|\psi\rangle = a|0\rangle + b|1\rangle$ be the state of a qbit.
- The probability that we will observe **0** or **1** is given by

$$P(0) = |a|^2 \text{ and } P(1) = b^2$$

Measurement, Projection and Collapse

- Let $|\psi\rangle = a|0\rangle + b|1\rangle$ be the state of a qbit.
- The probability that we will observe **0** or **1** is given by

$$P(0) = |a|^2 \text{ and } P(1) = b^2$$

- The total probability is $a^2 + b^2 = \langle\psi|\psi\rangle = 1$.

Measurement, Projection and Collapse

- Let $|\psi\rangle = a|0\rangle + b|1\rangle$ be the state of a qbit.
- The probability that we will observe **0** or **1** is given by

$$P(0) = |a|^2 \text{ and } P(1) = b^2$$

- The total probability is $a^2 + b^2 = \langle\psi|\psi\rangle = 1$.
- We can more conveniently express these probabilities through the projection operators.

Measurement, Projection and Collapse

- Let $|\psi\rangle = a|0\rangle + b|1\rangle$ be the state of a qbit.
- The probability that we will observe **0** or **1** is given by
$$P(0) = |a|^2 \text{ and } P(1) = b^2$$
- The total probability is $a^2 + b^2 = \langle\psi|\psi\rangle = 1$.
- We can more conveniently express these probabilities through the projection operators.
- Remember: a projection is a linear transformation P from a vector space to itself such that $P^2 = P$.

Measurement, Projection and Collapse

- Let $|\psi\rangle = a|0\rangle + b|1\rangle$ be the state of a qbit.
- The probability that we will observe **0** or **1** is given by
$$P(0) = |a|^2 \text{ and } P(1) = b^2$$
- The total probability is $a^2 + b^2 = \langle\psi|\psi\rangle = 1$.
- We can more conveniently express these probabilities through the projection operators.
- Remember: a projection is a linear transformation P from a vector space to itself such that $P^2 = P$.
- I.e., if we apply it twice, it is the same as if we apply it once.

Projection Operators

- Let P_0 be the projector operator that projects onto $|\mathbf{0}\rangle$:

Projection Operators

- Let P_0 be the projector operator that projects onto $|\mathbf{0}\rangle$:

$$P_0 = \begin{bmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}$$

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply $(\langle\psi|P_0)(P_0|\psi\rangle)$

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply

$$(\langle\psi|P_0)(P_0|\psi\rangle) = \langle\psi|P_0P_0|\psi\rangle$$

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply

$$(\langle\psi|P_0)(P_0|\psi\rangle) = \langle\psi|P_0P_0|\psi\rangle = \langle\psi|P_0^2|\psi\rangle$$

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply

$$(\langle\psi|P_0)(P_0|\psi\rangle) = \langle\psi|P_0P_0|\psi\rangle = \langle\psi|P_0^2|\psi\rangle = \langle\psi|P_0|\psi\rangle = ||P_0|\psi\rangle||^2$$

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply

$$(\langle\psi|P_0)(P_0|\psi\rangle) = \langle\psi|P_0P_0|\psi\rangle = \langle\psi|P_0^2|\psi\rangle = \langle\psi|P_0|\psi\rangle = ||P_0|\psi\rangle||^2$$

Which is equal to a^2

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply

$$(\langle\psi|P_0)(P_0|\psi\rangle) = \langle\psi|P_0P_0|\psi\rangle = \langle\psi|P_0^2|\psi\rangle = \langle\psi|P_0|\psi\rangle = ||P_0|\psi\rangle||^2$$

Which is equal to a^2

- Similarly we can define the projection operator P_1 onto $|1\rangle$.

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply

$$(\langle\psi|P_0)(P_0|\psi\rangle) = \langle\psi|P_0P_0|\psi\rangle = \langle\psi|P_0^2|\psi\rangle = \langle\psi|P_0|\psi\rangle = ||P_0|\psi\rangle||^2$$

Which is equal to a^2

- Similarly we can define the projection operator P_1 onto $|1\rangle$.
- These projection operators sum up to 1: $P_0 + P_1 = \mathbf{1}$.

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply
 $(\langle\psi|P_0)(P_0|\psi\rangle) = \langle\psi|P_0P_0|\psi\rangle = \langle\psi|P_0^2|\psi\rangle = \langle\psi|P_0|\psi\rangle = ||P_0|\psi\rangle||^2$

Which is equal to a^2

- Similarly we can define the projection operator P_1 onto $|1\rangle$.
- These projection operators sum up to 1: $P_0 + P_1 = \mathbf{1}$.
- The total probability $P(0) + P(1)$ is

Projection Operators

- Let P_0 be the projector operator that projects onto $|0\rangle$:

$$P_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

- The probability that we will observe the value 0 is simply
 $(\langle\psi|P_0)(P_0|\psi\rangle) = \langle\psi|P_0P_0|\psi\rangle = \langle\psi|P_0^2|\psi\rangle = \langle\psi|P_0|\psi\rangle = ||P_0|\psi\rangle||^2$

Which is equal to a^2

- Similarly we can define the projection operator P_1 onto $|1\rangle$.
- These projection operators sum up to 1: $P_0 + P_1 = \mathbf{1}$.
- The total probability $P(0) + P(1)$ is
 $\langle\psi|P_0|\psi\rangle + \langle\psi|P_1|\psi\rangle = \langle\psi|(P_0 + P_1)|\psi\rangle = \langle\psi|\psi\rangle = 1.$

Projection Operators

- We can define P_0 and P_1 in terms of *outer products*:

Projection Operators

- We can define P_0 and P_1 in terms of outer products:
- $P_0 = |\mathbf{0}\rangle\langle\mathbf{0}|$
- $P_1 = |\mathbf{1}\rangle\langle\mathbf{1}|$

Projection Operators

- We can define P_0 and P_1 in terms of outer products:
- $P_0 = |\mathbf{0}\rangle\langle\mathbf{0}|$
- $P_1 = |\mathbf{1}\rangle\langle\mathbf{1}|$
- Equivalent to $\boldsymbol{v}\boldsymbol{v}^T$ for some vector $\boldsymbol{v} \in \boldsymbol{R}^n$.

Projection Operators

- We can define P_0 and P_1 in terms of outer products:
- $P_0 = |\mathbf{0}\rangle\langle\mathbf{0}|$
- $P_1 = |\mathbf{1}\rangle\langle\mathbf{1}|$
- Equivalent to $\mathbf{v}\mathbf{v}^T$ for some vector $\mathbf{v} \in \mathbf{R}^n$.
- It is a $n \times n$ matrix.

Projection Operators

- We can define P_0 and P_1 in terms of outer products:
- $P_0 = |0\rangle\langle 0|$
- $P_1 = |1\rangle\langle 1|$
- Equivalent to $\boldsymbol{v}\boldsymbol{v}^T$ for some vector $\boldsymbol{v} \in \boldsymbol{R}^n$.
- It is a $n \times n$ matrix.
- After we have observed the truth value 0, the qbit has gone from the superposition $|\boldsymbol{\psi}\rangle = \boldsymbol{a}|0\rangle + \boldsymbol{b}|1\rangle$ to the definite state $|0\rangle$.

Projection Operators

- We can define P_0 and P_1 in terms of outer products:
- $P_0 = |\mathbf{0}\rangle\langle\mathbf{0}|$
- $P_1 = |\mathbf{1}\rangle\langle\mathbf{1}|$
- Equivalent to $\mathbf{v}\mathbf{v}^T$ for some vector $\mathbf{v} \in \mathbf{R}^n$.
- It is a $n \times n$ matrix.
- After we have observed the truth value 0, the qbit has gone from the superposition $|\psi\rangle = \mathbf{a}|\mathbf{0}\rangle + \mathbf{b}|\mathbf{1}\rangle$ to the definite state $|\mathbf{0}\rangle$.
- We must normalize $|\mathbf{0}\rangle = P_0|\psi\rangle$ so that the probabilities sum up to 1.

Projection Operators

- We must normalize $|\mathbf{0}\rangle = \mathbf{P}_0|\psi\rangle$ so that the probabilities sum up to 1:

Projection Operators

- We must normalize $|\mathbf{0}\rangle = P_0|\psi\rangle$ so that the probabilities sum up to 1:

$$|\mathbf{0}\rangle = \frac{P_0|\psi\rangle}{\langle\psi|P_0|\psi\rangle}$$

Projection Operators

- We must normalize $|\mathbf{0}\rangle = P_0|\psi\rangle$ so that the probabilities sum up to 1:

$$|\mathbf{0}\rangle = \frac{P_0|\psi\rangle}{\langle\psi|P_0|\psi\rangle}$$

- This is called the ***collapse of the wave function***.

The Pauli Matrices

- They are denoted by X, Y, Z and they are important operators acting on one qbit.
- $X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$
- It has the following effect on the standard basis:
- $X|0\rangle = |1\rangle$
- $X|1\rangle = |0\rangle$
- **Flip operator.**

The Pauli Matrices

- $Y = i \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$
- It has the following effect on the standard basis:
- $Y|0\rangle = i|1\rangle$
- $Y|1\rangle = -i|0\rangle$
- Keep the bit unchanged but induces a phase change (in the complex plane)

The Pauli Matrices

- $Z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$
- It has the following effect on the standard basis:
- $Z|0\rangle = |0\rangle$
- $Z|1\rangle = -|1\rangle$
- The effect of Z (and Y) is fundamentally quantum: it leaves the truth value (bit) unchanged but induces a phase change of -1 if the qbit is true.
- We can write $Y = iXZ$ which switches the two bit values and also adds a phase of $+i, -i$.

The Pauli Matrices



The Hadamard matrix

- It is defined as

$$H = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

- It has the following effect on the two basic states:
- $H|0\rangle = \frac{1}{\sqrt{2}} (|0\rangle + |1\rangle)$
- $H|1\rangle = \frac{1}{\sqrt{2}} (|0\rangle - |1\rangle)$.
- It puts the basis into a superposition.
- Useful when we want to prepare a totally random state.
- H also interferes the superposition so we can recover the original states.

Back to the state space

- Suppose we have now two (q)bits x_1, x_2 .

Back to the state space

- Suppose we have now two (q)bits x_1, x_2 .
- These bits can be in one of the following four possible states

Back to the state space

- Suppose we have now two (q)bits x_1, x_2 .
- These bits can be in one of the following four possible states

1. $|x_1\rangle|x_2\rangle = |0\rangle|0\rangle$

Back to the state space

- Suppose we have now two (q)bits x_1, x_2 .
- These bits can be in one of the following four possible states

1. $|x_1\rangle|x_2\rangle = |0\rangle|0\rangle$

2. $|x_1\rangle|x_2\rangle = |0\rangle|1\rangle$

Back to the state space

- Suppose we have now two (q)bits x_1, x_2 .
- These bits can be in one of the following four possible states

1. $|x_1\rangle|x_2\rangle = |0\rangle|0\rangle$

2. $|x_1\rangle|x_2\rangle = |0\rangle|1\rangle$

3. $|x_1\rangle|x_2\rangle = |1\rangle|0\rangle$

Back to the state space

- Suppose we have now two (q)bits x_1, x_2 .
- These bits can be in one of the following four possible states

1. $|x_1\rangle|x_2\rangle = |0\rangle|0\rangle$

2. $|x_1\rangle|x_2\rangle = |0\rangle|1\rangle$

3. $|x_1\rangle|x_2\rangle = |1\rangle|0\rangle$

4. $|x_1\rangle|x_2\rangle = |1\rangle|1\rangle$

Back to the state space

- Suppose we have now two (q)bits x_1, x_2 .
- These bits can be in one of the following four possible states
 1. $|x_1\rangle|x_2\rangle = |0\rangle|0\rangle$
 2. $|x_1\rangle|x_2\rangle = |0\rangle|1\rangle$
 3. $|x_1\rangle|x_2\rangle = |1\rangle|0\rangle$
 4. $|x_1\rangle|x_2\rangle = |1\rangle|1\rangle$
- We write, for convenience, $|x_1x_2\rangle = |01\rangle$ etc.

Back to the state space

- Suppose we have now two (q)bits x_1, x_2 .
- These bits can be in one of the following four possible states
 1. $|x_1\rangle|x_2\rangle = |0\rangle|0\rangle$
 2. $|x_1\rangle|x_2\rangle = |0\rangle|1\rangle$
 3. $|x_1\rangle|x_2\rangle = |1\rangle|0\rangle$
 4. $|x_1\rangle|x_2\rangle = |1\rangle|1\rangle$
- We write, for convenience, $|x_1x_2\rangle = |01\rangle$ etc.
- We can even write $|x_1x_2\rangle = |3\rangle$ meaning $|x_1x_2\rangle = |11\rangle$ etc.

Operations on multiple (q)bits

- Imagine an instruction of the form $x_2 := x_1$.

Operations on multiple (q)bits

- Imagine an instruction of the form $x_2 := x_1$.
- What is the mapping of old states to new?

Operations on multiple (q)bits

- Imagine an instruction of the form $x_2 := x_1$.
- What is the mapping of old states to new?
- $|01\rangle$ maps to $|00\rangle$.

Operations on multiple (q)bits

- Imagine an instruction of the form $x_2 := x_1$.
- What is the mapping of old states to new?
- $|01\rangle$ maps to $|00\rangle$.
- $|10\rangle$ maps to $|11\rangle$.

Operations on multiple (q)bits

- Imagine an instruction of the form $x_2 := x_1$.
- What is the mapping of old states to new?
- $|01\rangle$ maps to $|00\rangle$.
- $|10\rangle$ maps to $|11\rangle$.
- The other two states remain the same.

Operations on multiple (q)bits

- Imagine an instruction of the form $x_2 := x_1$.
- What is the mapping of old states to new?
- $|01\rangle$ maps to $|00\rangle$.
- $|10\rangle$ maps to $|11\rangle$.
- The other two states remain the same.
- So, each state is a basis vector in the $2^2 = 4$ – dimensional space.

Operations on multiple (q)bits

- Imagine an instruction of the form $x_2 := x_1$.
- What is the mapping of old states to new?
- $|01\rangle$ maps to $|00\rangle$.
- $|10\rangle$ maps to $|11\rangle$.
- The other two states remain the same.
- So, each state is a basis vector in the $2^2 = 4$ – dimensional space.
- For example

$$|00\rangle = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, |01\rangle = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, |10\rangle = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, |11\rangle = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

Operations on multiple (q)bits

- Imagine an instruction of the form $x_2 := x_1$.
- How can we describe this transformation?
- *We need a 4 times 4 matrix (a system of two (q)bits has 4 amplitudes).*

Operations on multiple (q)bits

- Imagine an instruction of the form $x_2 := x_1$.
- How can we describe this transformation?
- *We need a 4 times 4 matrix (a system of two (q)bits has 4 amplitudes)*

$$U = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}$$

Composition of Quantum Systems

- *Postulate 4*

The state space of a composite physical system $|\psi\rangle$ is the tensor product of the state spaces of the constituent components physical systems, i.e., $|\psi\rangle = |\psi_1\rangle \otimes |\psi_2\rangle \otimes \cdots \otimes |\psi_n\rangle$

I.e., if $|\psi\rangle$ is a 2-qbit system consisting of $|q_1\rangle, |q_2\rangle$ then $|\psi\rangle = |q_1\rangle \otimes |q_2\rangle$