VTPEH 6270 - Check Point 06

Jinchan Sun

2026-04-09

Contents

1 Introduction **1**

2 Material & Methods **2**

3 Results **3**

4 Conclusions **8**

5 References **8**

6 AI Use Disclosure Statement **8**

1 Introduction

This report explores county-level health indicators using the CDC PLACES 2025 dataset. The dataset includes health outcomes, prevention measures, and risk factors across U.S. counties. Due to file size limitations, a reduced sample of the original dataset (PLACES_sample.csv) is used for all analyses to ensure reproducibility of the results. The goal of this report is to explore potential associations between selected health indicators and demographic or behavioral characteristics at the county level, using county-level aggregated data.

1.1 Research Question

Is adult obesity prevalence associated with physical inactivity at the county level?

1.2 Scientific Plausibility and Context

Physical inactivity is widely recognized as a major public health concern. Global health authorities identify insufficient physical activity as an important risk factor for non-communicable diseases, including obesity (World Health Organization 2024). A recent systematic review reported consistent evidence that sedentary behavior and low physical activity are associated with increased abdominal obesity and overall adiposity in adults (Silveira et al. 2022). Earlier research has further suggested a possible vicious cycle, in which physical inactivity contributes to weight gain, and increased body weight may in turn reduce physical activity levels over time (Pietiläinen et al. 2007).

Based on existing literature, we expect that higher levels of physical inactivity will be positively associated with higher adult obesity prevalence at the county level. This analysis focuses on examining associations rather than causal relationships. County-level analysis allows us to identify geographic patterns and inform local public health interventions.

2 Material & Methods

2.1 Data

The data used in this analysis come from the CDC PLACES 2025 dataset, which provides model-based estimates of health outcomes, prevention measures, and risk factors at the county level across the United States. The dataset is publicly available from the CDC data portal: https://data.cdc.gov/500-Cities-Places/PLACES-Local-Data-for-Better-Health-County-Data-20/swc5-untb/about_data

PLACES data are derived using small-area estimation methods based on the Behavioral Risk Factor Surveillance System (BRFSS) and U.S. Census data, allowing for consistent health estimates across all U.S. counties :contentReferenceoaicite:0.

Due to file size limitations, a reduced sample of the dataset (PLACES_sample.csv) was used for all analyses to ensure reproducibility. This sample dataset retains the key variables of interest, including adult obesity prevalence and physical inactivity.

Before analysis, the dataset was inspected and cleaned as needed. Relevant variables were selected, and observations with missing values for the variables of interest were removed. No additional transformations were applied.

2.2 Statistical Analyses

To examine the association between physical inactivity and adult obesity prevalence at the county level, a combination of exploratory, descriptive, and inferential statistical approaches was used.

First, exploratory data analysis was conducted to understand the distribution of key variables. Histograms were used to assess the distribution of adult obesity prevalence across counties. Scatterplots were also generated to visualize relationships between obesity prevalence and other variables, including total population and physical inactivity.

Next, descriptive analyses were performed to compare obesity prevalence across different levels of physical inactivity. Counties were categorized into “high” and “low” physical inactivity groups based on the median value of physical inactivity. Summary statistics, including mean, standard deviation, and median obesity prevalence, were calculated for each group. A boxplot was used to visualize differences between these groups.

To formally assess the association between physical inactivity and obesity prevalence, a linear regression model was fitted, with obesity prevalence as the outcome variable and physical inactivity as the explanatory variable. A scatterplot with a fitted regression line was used to visualize this relationship.

Model assumptions were evaluated using diagnostic methods. Linearity and homoscedasticity were assessed using a residual plot, where residuals were plotted against fitted values. The distribution of residuals was visually inspected to assess normality.

In addition, a simulation analysis was conducted to explore how effect size, sample size, and variability influence the observed association between physical inactivity and obesity. Synthetic datasets were generated under varying conditions, and summary statistics were compared across simulated scenarios. A heatmap was used to visualize how these parameters affected differences in physical inactivity between obesity groups.

2.3 Code

All code used for data cleaning, visualization, and statistical analysis is available in the following GitHub repository:

https://github.com/Irene-sjc/R-project

The repository contains the R scripts used in this project, the sample dataset used for reproducible analyses, and the output figures generated from the code. Due to file size limitations, the full CDC PLACES 2025 dataset is not included in the repository. Instead, a reduced sample dataset is provided so that the analysis can be reproduced on any machine.

3 Results

```r
# Load package
library(here)
library(ggplot2)
library(dplyr)
library(tidyr)

# Load data
data <- read.csv(here("data", "raw", "PLACES_sample.csv"), stringsAsFactors = FALSE)

# Make sure key columns are usable
data$Data_Value <- suppressWarnings(as.numeric(data$Data_Value))
data$Year <- suppressWarnings(as.numeric(data$Year))

# Filter data
data_clean <- data %>%
  filter(
    Data_Value_Type == "Crude prevalence",
    Year == 2023
  )

# Keep only the two measures needed
selected_long <- data_clean %>%
  filter(Measure %in% c(
    "No leisure-time physical activity among adults",
    "Obesity among adults"
  )) %>%
  select(LocationName, Measure, Data_Value)

# Convert to wide format
selected <- selected_long %>%
  pivot_wider(
    names_from = Measure,
    values_from = Data_Value,
    values_fn = mean
  ) %>%
  rename(
    County = LocationName,
    Physical_Inactivity = `No leisure-time physical activity among adults`,
```

```r
    Obesity = `Obesity among adults`
  ) %>%
  drop_na()

# Create high/low inactivity group
median_inactivity <- median(selected$Physical_Inactivity, na.rm = TRUE)

selected <- selected %>%
  mutate(
    Inactivity_Group = ifelse(
      Physical_Inactivity > median_inactivity,
      "High Inactivity",
      "Low Inactivity"
    )
  )

# Fit model
model <- lm(Obesity ~ Physical_Inactivity, data = selected)
```

3.1 Visualization

Figure 1 shows the association between physical inactivity and obesity prevalence across counties.

```r
ggplot(selected, aes(x = Physical_Inactivity, y = Obesity)) +
  geom_point(alpha = 0.4) +
  geom_smooth(method = "lm", se = FALSE, color = "red") +
  labs(
    x = "Physical inactivity (%)",
    y = "Obesity prevalence (%)"
  ) +
  theme_minimal()
```



A positive association is observed in the scatterplot, with the regression line indicating that counties with higher levels of physical inactivity tend to have higher obesity prevalence.

Figure 2 compares obesity prevalence between counties with high and low levels of physical inactivity.

```
ggplot(selected, aes(x = Inactivity_Group, y = Obesity)) +
  geom_boxplot(fill = "steelblue") +
  labs(
    x = "Physical inactivity group",
    y = "Obesity prevalence (%)"
  ) +
  theme_minimal()
```



The boxplot shows that counties in the high physical inactivity group tend to have higher obesity prevalence compared to those in the low inactivity group.

3.2 Statistical Analyses

Counties in the high physical inactivity group generally show higher obesity prevalence compared to those in the low inactivity group.

```r
summary(model)
```

```
##
## Call:
## lm(formula = Obesity ~ Physical_Inactivity, data = selected)
##
## Residuals:
##     Min      1Q  Median      3Q     Max
## -5.8621 -0.8113  0.2094  1.9413  6.3172
##
## Coefficients:
##                     Estimate Std. Error t value Pr(>|t|)
## (Intercept)          27.0309     4.4671   6.051 1.68e-05 ***
## Physical_Inactivity   0.3793     0.1361   2.787   0.0132 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
```

```
## Residual standard error: 3.283 on 16 degrees of freedom
## Multiple R-squared:  0.3269, Adjusted R-squared:  0.2848
## F-statistic:  7.77 on 1 and 16 DF,  p-value: 0.01318
```

The linear regression analysis showed a statistically significant positive association between physical inactivity and obesity prevalence. The estimated coefficient for physical inactivity was 0.379 ($p = 0.013$), indicating that a one-percentage-point increase in physical inactivity is associated with an approximately 0.38 percentage-point increase in obesity prevalence at the county level.The model explained a moderate proportion of the variation in obesity prevalence ($R^2 = 0.327$). Overall, these results suggest that higher levels of physical inactivity are associated with higher levels of obesity across counties.

Figure 3 presents the residual plot used to assess model assumptions.

```r
ggplot(
  data.frame(
    Fitted = fitted(model),
    Residuals = resid(model)
  ),
  aes(x = Fitted, y = Residuals)
) +
  geom_point(alpha = 0.4) +
  geom_hline(yintercept = 0, color = "red") +
  labs(
    x = "Fitted values",
    y = "Residuals"
  ) +
  theme_minimal()
```



The residuals are distributed around zero without a clear pattern, suggesting that the assumptions of linearity and homoscedasticity are reasonably satisfied.

Overall, the findings support a positive association between physical inactivity and adult obesity prevalence. As the analysis is based on observational county-level data, the results should be interpreted as associations rather than causal relationships.

4 Conclusions

The results of this analysis are consistent with the expectations outlined in the introduction and with findings from existing literature. A positive association between physical inactivity and adult obesity prevalence was observed across counties. Both the scatterplot and the linear regression model indicated that higher levels of physical inactivity are associated with higher levels of obesity.

These findings align with prior research suggesting that insufficient physical activity is an important risk factor for obesity and related chronic conditions. The observed relationship also supports the idea of a potential cycle in which physical inactivity contributes to weight gain, and higher body weight may in turn reduce physical activity over time.

However, this analysis has several limitations. The data are aggregated at the county level and are based on modeled estimates, which may not reflect individual-level variation. In addition, the cross-sectional design does not allow for causal inference, and other factors not included in the analysis may also influence obesity prevalence.

Overall, the findings suggest that promoting physical activity at the population level may be an important strategy for reducing obesity prevalence. Future research could incorporate additional variables or longitudinal data to better understand the mechanisms underlying this relationship.

5 References

Pietiläinen, K. H., J. Kaprio, P. Borg, M. Koskenvuo, and A. Rissanen. 2007. "Physical Inactivity and Obesity: A Vicious Circle." *Obesity* 15 (2): 409–14. https://doi.org/10.1038/oby.2007.72.

Silveira, E. A., M. C. Romualdo, M. Noll, T. C. de Souza, C. L. Santos, and C. de Oliveira. 2022. "Sedentary Behavior, Physical Inactivity, Abdominal Obesity and Obesity in Adults and Older Adults: A Systematic Review and Meta-Analysis." *Clinical Nutrition ESPEN* 48: 63–75. https://doi.org/10.1016/j.clnesp.2022.06.001.

World Health Organization. 2024. "Physical Activity." https://www.who.int/news-room/fact-sheets/detail/physical-activity.

6 AI Use Disclosure Statement

Yes. This document was generated with assistance from ChatGPT, which was used to help debug and improve code written by the author. The primary code and analysis were completed by the author.