

The Immune System and Cancer

How the body detects, controls, and sometimes loses to cancer cells

The immune system is constantly patrolling for emerging abnormal cells and can often eliminate precancerous and early cancer cells before they ever become clinically relevant. Cancer only becomes a problem when those abnormal cells acquire ways to hide from, suppress, or outpace immune surveillance.

Big Picture: Cancer Immunosurveillance

- Throughout life, cells accumulate DNA damage from replication errors, inflammation, radiation, and toxins.
- Most damaged cells are repaired, forced into senescence, or eliminated by apoptosis; immune cells help recognize and clear those that slip through intrinsic checkpoints.
- This ongoing “cancer watch” is called immunosurveillance and is thought to prevent many would-be cancers from ever establishing.

A useful way to frame it is the “three E’s” model of cancer–immune interaction: Elimination → Equilibrium → Escape.

- **Elimination:** immune system detects and destroys nascent transformed cells.
- **Equilibrium:** small populations of tumor cells persist but are held in check.
- **Escape:** subclones evolve that evade control, leading to clinically apparent cancer.

In plain terms:

Your immune system isn't just for fighting colds — it's also a built-in cancer patrol. Most abnormal cells get caught and destroyed immediately (elimination). Some survive but get “held down” for years without growing (equilibrium). A few eventually find ways to escape and become a real tumor.

Step 1 – Handling Stressed and Precancerous Cells

Intrinsic Cellular Safeguards

Before the immune system even acts, cells have internal anti-cancer programs:

- DNA damage sensors (like p53 pathways) stop the cell cycle, trigger repair, or induce apoptosis when damage is excessive.
- Oncogene activation and telomere dysfunction can push cells into senescence, a non-dividing but metabolically active state.

Immune cells then help “clean up”:

- Senescent and apoptotic cells expose “eat me” signals (like phosphatidylserine) that attract phagocytes such as macrophages and dendritic cells.
- These phagocytes engulf debris and can present abnormal peptides from those cells to T cells, priming adaptive responses against similar abnormal cells.

Stress Ligands on Precancerous Cells

Cells under oncogenic or genotoxic stress upregulate surface molecules that flag them as “altered self”:

- Many precancerous cells increase ligands for activating receptors on natural killer (NK) cells, such as NKG2D ligands (e.g., MICA/B in humans).
- At the same time they may show reduced or altered major histocompatibility complex class I (MHC I), which also signals NK cells that something is wrong.

NK cells then:

- Recognize the imbalance between “normal self” signals and “stress” signals.
- Release perforin and granzymes to induce apoptosis in the stressed cell, often before it is fully malignant.

Innate Inflammation Around Damaged Tissue

When there is chronic irritation or damage (e.g., due to smoking or infection), innate immune cells constantly sample the environment:

- Pattern-recognition receptors on macrophages and dendritic cells detect danger signals from damaged or dying precancerous cells.
- This can trigger local inflammation that both helps clear abnormal cells and, paradoxically, if prolonged, can supply growth factors and DNA-damaging reactive species that promote further mutations.

So at the precancerous stage, you have a combined effect of intrinsic checkpoints, NK-cell surveillance, and phagocytic clearance constraining clonal expansion.

In plain terms:

Every cell has an internal alarm system (p53) that stops badly damaged cells from dividing, or self-destructs them. If a cell dodges that, “natural killer” (NK) immune cells act like bouncers — they check for a mismatched ID badge (weird stress signals plus missing normal markers) and kill on the spot.

Step 2 – Detecting and Targeting Early Cancer Cells

Once cells cross into true malignancy (sustained proliferation, invasion potential), the immune response becomes more organized and adaptive.

Tumor Antigens and Immune Recognition

Cancer cells differ from normal cells in ways the immune system can “see”:

- They often express neoantigens: novel peptides derived from mutated proteins presented on MHC I molecules.
- They may also overexpress normal self proteins (e.g., differentiation antigens) at abnormal levels or in unusual contexts, creating tumor-associated antigens.

Dendritic cells (DCs) are central here:

- DCs ingest dying tumor or precancerous cells, process the proteins, and present peptide fragments on MHC molecules.
- They migrate to lymph nodes and prime naïve CD8 and CD4 T cells against these antigens, turning a local event into system-wide “memory” of what tumor cells look like.

Cytotoxic T Cells as Precision Killers

Activated CD8 T cells (cytotoxic T lymphocytes, CTLs):

- Patrol tissues and scan MHC I on cells for the specific tumor peptide they recognize.
- When they find a match, they form an immune synapse and release perforin and granzymes, triggering apoptosis of that single target cell.
- They can also engage death receptors (e.g., Fas–FasL) to induce programmed cell death.

This is a highly specific system: each CTL clone recognizes a particular peptide–MHC combination, allowing discrimination of transformed vs healthy neighboring cells.

Helper T Cells and Orchestration

CD4 T helper cells support the anti-tumor response by:

- Providing cytokines (e.g., interferon-gamma) that increase MHC expression on tumor and stromal cells, making them more visible to CTLs and NK cells.
- Enhancing the activity of macrophages and promoting B-cell antibody production against tumor-associated antigens.

Antibodies and Complement

B cells and antibodies play a supporting but important role:

- Antibodies can bind surface antigens on tumor cells, flagging them for antibody-dependent cellular cytotoxicity (ADCC) by NK cells and other effector cells.
- Complement activation on antibody-coated tumor cells can contribute to lysis or opsonization for phagocytosis.

In plain terms:

Once a cell is fully cancerous, it often makes odd new proteins (neoantigens). Dendritic cells “photograph” these and show them to T cells in the lymph nodes, training a targeted killing squad (CD8 T cells) that can recognize and destroy only cells carrying that specific marker — precision, not carpet-bombing.

Step 3 – Keeping Small Tumors in “Equilibrium”

Not all malignant clones are wiped out at once. Some are held in a dynamic steady state:

- The immune system may contain small clusters of tumor cells so they divide and die at roughly the same rate, never forming a detectable mass.
- This equilibrium can last for years, with continuous selective pressure favoring tumor variants that better evade or resist immunity.

Mechanistically:

- CD8 and CD4 T cells, NK cells, and macrophages keep attacking, producing interferon-gamma and cytotoxic mediators that restrain growth.
- Tumor cells under this pressure may down-modulate immunogenic antigens or MHC I, or alter signaling pathways to resist apoptosis.

This is analogous to low-level chronic infection control: the host and the tumor “coexist” until either immune function diminishes or the tumor evolves sufficient escape mechanisms.

In plain terms:

Sometimes the immune system doesn't fully win or lose — it just keeps a tiny tumor in check for years, like a stalemate, while the cancer cells keep trying new tricks to break free.

Step 4 – How Cancers Evade and Suppress Immunity

Clinically apparent cancer usually implies the tumor has navigated around one or more layers of immune defense. Key escape strategies include:

1. Hiding Their Identity

- Downregulating MHC I so CTLs can no longer recognize tumor antigens effectively.
- Losing or editing the most immunogenic neoantigens, leaving only low-visibility subclones.
- Upregulating non-classical MHC molecules that inhibit NK cells despite reduced classical MHC I.

2. Hitting the “Brakes” on T Cells

T cells naturally have inhibitory checkpoints to prevent autoimmunity; tumors hijack these:

- Tumor and stromal cells often overexpress ligands such as PD-L1 that bind PD-1 on T cells, sending a “stand down” signal.
- Many tumors upregulate other checkpoints like CTLA-4 signaling pathways that dampen T-cell activation.

This leads to T-cell exhaustion:

- T cells remain present but less functional, with reduced cytokine production, proliferation, and cytotoxicity over time.

3. Creating an Immunosuppressive Microenvironment

- Tumors secrete immunosuppressive cytokines like TGF-beta and IL-10 that blunt effector T cells and NK cells while promoting regulatory T cells (Tregs) and myeloid-derived suppressor cells (MDSCs).
- Tumor-associated macrophages often shift toward a pro-tumor phenotype that supports angiogenesis, matrix remodeling, and suppression of local immune responses.
- Abnormal vasculature and high interstitial pressure make it physically harder for effector T cells to infiltrate the tumor core.

4. Outpacing the Response

- If proliferation is very rapid, immune cells may simply not kill fast enough to prevent net growth, especially in the setting of systemic immune compromise (e.g., age, immunosuppressive drugs, chronic infection).

In plain terms:

Cancer cells evolve like a fugitive learning to avoid cameras. They hide their ID (lower visibility), disable the “kill switch” T cells normally respect (checkpoints), and poison the neighborhood with signals that calm down nearby immune cells.

Putting It Together: From Prevention to Therapy

In simple terms, the immune system acts at multiple levels:

- **Preventive surveillance:** clearing stressed, senescent, and precancerous cells via intrinsic checkpoints, NK cells, and phagocytes.
- **Targeted killing of cancer cells:** CTLs, NK cells, antibodies, and helper T cells coordinate to recognize neoantigens and destroy malignant cells.
- **Long-term control:** maintaining some tumors in a low-level equilibrium until they either get eliminated or evolve escape mechanisms.

Modern immunotherapies are essentially tools to re-tilt this balance in favor of the host:

- Checkpoint inhibitors block inhibitory pathways like PD-1/PD-L1 or CTLA-4, “releasing the brakes” on T cells so they can resume attacking tumor cells.
- CAR T cells and other engineered cell therapies enhance T-cell recognition of tumor antigens.
- Cancer vaccines and oncolytic viruses aim to increase the visibility and immunogenicity of tumor cells and improve priming of adaptive responses.

Sources

- How The Immune System Fights Cancer – Immunology Explained (immunologyexplained.aai.org)
- The immune system and cancer – Cancer Research UK (cancerresearchuk.org)
- Roles of the immune system in cancer: from tumor initiation to... – PMC ([pmc.ncbi.nlm.nih.gov/articles/PMC6169832/](https://pubmed.ncbi.nlm.nih.gov/articles/PMC6169832/))
- Harnessing the immune system against cancer – PMC ([pmc.ncbi.nlm.nih.gov/articles/PMC8605995/](https://pubmed.ncbi.nlm.nih.gov/articles/PMC8605995/))
- Host Response to Tumors – MSD Manual Professional Edition ([msdmanuals.com](https://www.msdmanuals.com))

- How Does the Body Fight Cancer? 7 Key Ways Your Immune System... (int.livhospital.com)