15. Atlantic Salmon (Salmo salar)

Sensitivity Assessment

No existing MarESA or FeAST sensitivity assessments for salmon were available, therefore a full literature search and MarESA assessment were conducted. The OSPAR status assessment in 2022 covered the majority of the identified pressures and was used to supplement the MarESA results. A recent review of marine stressors impacting salmon (Gillson *et al.*, 2022) identified major sensitivities and supporting literature. To identify key considerations in the Celtic Sea, an external expert in the subject area was also consulted. The following sensitivity assessment is therefore based on the MarESA protocol and OSPAR assessment supplemented with key literature.

Table A11.15. Sensitivity assessment for Atlantic Salmon (*Salmo salar***).** Associated sectors include activities related to offshore renewable energy (O), Fishing (F), or shipping (S). NR = not relevant, NA = not assessed, NEv = no evidence, H = high, M = medium, L = low, NS = not sensitive.

Pressures		Associated sector(s)	Resistance				Resilience				Sensitivity			
Classification	Pressure type	Sector(s)	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
Physical	Physical loss (to land or freshwater habitat)		None	Н	Н	Н	VL	Н	Н	Н	Н	Н	Н	Н
	Physical change (to another seabed type)	O, F	None	Н	Н	Н	VL	Н	Н	Н	Н	Н	Н	Н

Pressures		Associated sector(s)	Resistance				Resilience				Sensitivity			
Classification	Pressure type		Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
	Physical change (to another sediment type)	O, F									NEv	NR	NR	NR
	Habitat structure change-removal of substratum (extraction)	0									NEv	NR	NR	NR
	Abrasion/disturbance of substratum surface or seabed	O, F									NEv	NR	NR	NR
Physical	Penetration or disturbance of substratum subsurface	O, F									NEv	NR	NR	NR
	Changes in suspended solids (water clarity)	O, F									NA	NR	NR	NR

Pressures		Associated	Resistance				Resilience				Sensitivity			
Classification	Pressure type	sector(s)	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
	Smothering and siltation changes (light)	0									NEv	NR	NR	NR
	Smothering and siltation changes (heavy)	0									NEv	NR	NR	NR
	Underwater noise	O, F, S									M	L	М	L
	Electromagnetic energy	0	М	М	M	L	L	L	M	L	М	L	M	L
	Barrier to species movement	O, F	L	Н	Н	Н	L	Н	Н	Н	Н	Н	Н	Н
	Death or injury by collision	O, F, S	L	Н	Н	Н	L	Н	Н	Н	Н	Н	Н	Н
Hydrological	Water flow changes	0									NEv	NR	NR	NR

Pressures		Associated	Resistance				Resilience				Sensitivity			
Classification	Pressure type	sector(s)	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
Chemical	Transition elements & organo-metal contamination	O, F, S									Sensitive	Н	Н	Н
	Hydrocarbon & PAH contamination	O, F, S									Sensitive	Н	Н	Н
	Synthetic compound contamination	O, F, S									Sensitive	Н	Н	Н
Chemical	Introduction of other substances	O, F, S									Sensitive	Н	Н	Н
	Deoxygenation	0									Sensitive	Н	Н	Н
Biological	Introduction or spread of invasive non-indigenous species	O, F, S									Sensitive	Н	Н	Н

		Associated sector(s)	Resistance				Resilience				Sensitivity			
Classification	Pressure type		Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
	Removal of target species	F	L	Н	Н	Н	VL	Н	Н	Н	Н	Н	Н	Н
	Removal of non-target species	F	L	Н	Н	Н	VL	Н	Н	Н	Н	Н	Н	Н

Key references for sensitivity assessment

- 1. Freyhof, J. (2014). Salmo salar (Europe assessment). The IUCN Red List of Threatened Species 2014: e.T19855A2532398. Accessed on 23 April 2024.
- Gillson, J.P., Bašić, T., Davison, P.I., Riley, W.D., Talks, L., Walker, A.M., & Russell, I.C. (2022). A review of marine stressors impacting Atlantic salmon *Salmo salar*, with an assessment of the major threats to English stocks. *Reviews in Fish Biology and Fisheries*, 32(3), 879-919. https://doi.org/10.1007/s11160-022-09714-x
- OSPAR quality Status Report (2023). Status Assessment 2022 Atlantic salmon https://oap.ospar.org/en/ospar-assessments/committee-assessments/biodiversity-committee/status-assessments/atlantic-salmon/

Full Reference list from MarESA Screening

- 1. Andrews, C.D., Payne, J.F., & Rise, M L. (2014). Identification of a gene set to evaluate the potential effects of loud sounds from seismic surveys on the ears of fishes: a study with *Salmo salar. Journal of Fish Biology*, 84(6), 1793-1819. https://doi.org/10.1111/jfb.12398
- Ciborowski, K.L., Consuegra, S., Garcia de Leaniz, C., Wang, J., Beaumont, M.A., & Jordan, W.C. (2007). Stocking may increase mitochondrial DNA diversity but fails to halt the decline of endangered Atlantic salmon populations. *Conservation Genetics*, 8, 1355-1367. http://dx.doi.org/10.1007/s10592-007-9286-2
- 3. Elliott, S.A., Deleys, N., Beaulaton, L., Rivot, E., Réveillac, E., & Acou, A. (2023). Fisheries-dependent and-Independent data used to model the distribution of diadromous fish at-sea. *Data in Brief*, 48, 109107. http://dx.doi.org/10.1016/j.dib.2023.109107
- Fairchild, W.L., Swansburg, E. O., Arsenault, J.T., & Brown, S. B. (1999). Does an association between pesticide use and subsequent declines in catch of Atlantic salmon (*Salmo salar*) represent a case of endocrine disruption?. *Environmental Health Perspectives*, 107(5), 349-358. http://dx.doi.org/10.2307/3434538
- Fausch, K. D. (2007). Introduction, establishment and effects of non-native salmonids:
 Considering the risk of rainbow trout invasion in the United Kingdom. *Journal of Fish Biology*,
 71, 1-32. http://dx.doi.org/10.1111/j.1095-8649.2007.01682.x

- Finnegan, A. K., & Stevens, J. R. (2008). Assessing the long-term genetic impact of historical stocking events on contemporary populations of Atlantic salmon, *Salmo salar*. *Fisheries Management and Ecology*, 15(4), 315-326. http://dx.doi.org/10.1111/j.1365-2400.2008.00616.x
- Gagnon, M. M., & Holdway, D. A. (2000). EROD induction and biliary metabolite excretion following exposure to the water accommodated fraction of crude oil and to chemically dispersed crude oil. *Archives of environmental contamination and toxicology*, 38, 70-77. http://dx.doi.org/10.1007/s002449910009
- Gillson, J. P., Bašić, T., Davison, P. I., Riley, W. D., Talks, L., Walker, A. M., & Russell, I. C. (2022).
 A review of marine stressors impacting Atlantic salmon *Salmo salar*, with an assessment of the major threats to English stocks. *Reviews in Fish Biology and Fisheries*, 32(3), 879-919.

 http://dx.doi.org/10.1007/s11160-022-09714-x
- Greco, L., Capri, E., & Rustad, T. (2007). Biochemical responses in *Salmo salar* muscle following exposure to ethynylestradiol and tributyltin. *Chemosphere*, 68(3), 564-571. http://dx.doi.org/10.1016/j.chemosphere.2006.12.033
- 10. Halttunen, E., Gjelland, K.Ø., Glover, K.A., Johnsen, I.A., Serra-Llinares, R.M., Skaala, Ø., Nilsen, R., Bjørn, P-A., Karlsen, Ø., Finstad, B., & Skilbrei, O.T. (2018). Migration of Atlantic salmon post-smolts in a fjord with high infestation pressure of salmon lice. *Marine Ecology Progress Series*, 592, 243-256.

http://dx.doi.org/10.3354/meps12403

- Horreo, J.L., Griffiths, A. M., Machado-Schiaffino, G., Stevens, J.R., & Garcia-Vazquez, E. (2019). Northern areas as refugia for temperate species under current climate warming:
 Atlantic salmon (Salmo salar L.) as a model in northern Europe. Journal of Fish Biology, 95(1), 304-310.
 - http://dx.doi.org/10.1111/jfb.13825
- 12. Jonsson, B., Jonsson, N., & Albretsen, J. (2016). Environmental change influences the life history of salmon Salmo salar in the North Atlantic Ocean. *Journal of Fish Biology*, 88(2), 618-637.

http://dx.doi.org/10.1111/jfb.12854

- 13. Kennedy, R.J., & Crozier, W.W. (2010). Evidence of changing migratory patterns of wild Atlantic salmon *Salmo salar* smolts in the River Bush, Northern Ireland, and possible associations with climate change. *Journal of Fish Biology*, *76*(7), 1786-1805. http://dx.doi.org/10.1111/j.1095-8649.2010.02617.x
- 14. Le Cam, S., Perrier, C., Besnard, A.L., Bernatchez, L., & Evanno, G. (2015). Genetic and phenotypic changes in an Atlantic salmon population supplemented with non-local individuals: a longitudinal study over 21 years. *Proceedings of the Royal Society B: Biological Sciences*, 282(1802), 20142765. http://dx.doi.org/10.1098/rspb.2014.2765
- 15. Lothian, A.J., Newton, M., Barry, J., Walters, M., Miller, R.C., & Adams, C.E. (2018). Migration pathways, speed and mortality of Atlantic salmon (*Salmo salar*) smolts in a Scottish river and the near-shore coastal marine environment. *Ecology of Freshwater Fish*, 27(2), 549-558. http://dx.doi.org/10.1111/eff.12369
- Lundebye, A.K., Berntssen, M.H.G., Bonga, S.W., & Maage, A. (1999). Biochemical and physiological responses in Atlantic salmon (*Salmo salar*) following dietary exposure to copper and cadmium. *Marine Pollution Bulletin*, 39(1-12), 137-144.
 http://dx.doi.org/10.1016/S0025-326X(98)00208-2
- 17. Lyssimachou, A., Jenssen, B.M., & Arukwe, A. (2006). Brain cytochrome P450 aromatase gene isoforms and activity levels in Atlantic salmon after waterborne exposure to nominal environmental concentrations of the pharmaceutical ethynylestradiol and antifoulant tributyltin. *Toxicological Sciences*, 91(1), 82-92. http://dx.doi.org/10.1093/toxsci/kfj136
- Meucci, V., & Arukwe, A. (2005). Detection of vitellogenin and zona radiata protein expressions in surface mucus of immature juvenile Atlantic salmon (*Salmo salar*) exposed to waterborne nonylphenol. *Aquatic toxicology*, 73(1), 1-10. http://dx.doi.org/10.1016/j.aquatox.2005.03.021
- 19. Mortensen, A.S., & Arukwe, A. (2006). The persistent DDT metabolite, 1, 1-dichloro-2, 2-bis (p-chlorophenyl) ethylene, alters thyroid hormone-dependent genes, hepatic cytochrome P4503A, and pregnane× receptor gene expressions in Atlantic salmon (*Salmo salar*) parr. Environmental Toxicology and Chemistry: An International Journal, 25(6), 1607-1615. http://dx.doi.org/10.1897/05-376R1.1

- 20. Nedwell, J.R., Turnpenny, A.W., Lovell, J.M., & Edwards, B. (2006). An investigation into the effects of underwater piling noise on salmonids. *The Journal of the Acoustical Society of America*, 120(5), 2550-2554. http://dx.doi.org/10.1121/1.2335573
- 21. Parrish, D.L., Behnke, R.J., Gephard, S.R., McCormick, S.D., & Reeves, G.H. (1998). Why aren't there more Atlantic salmon (*Salmo salar*)?. *Canadian Journal of Fisheries and Aquatic Sciences*, 55(S1), 281-287. http://dx.doi.org/10.1139/cjfas-55-S1-281
- 22. Peyronnet, A., Friedland, K.D., & Ó Maoileidigh, N. (2008). Different ocean and climate factors control the marine survival of wild and hatchery Atlantic salmon *Salmo salar* in the north-east Atlantic Ocean. *Journal of Fish Biology*, 73(4), 945-962. http://dx.doi.org/10.1111/j.1095-8649.2008.01984.x
- 23. Solberg, C.B., Sæthre, L., & Julshamn, K. (2002). The effect of copper-treated net pens on farmed salmon (*Salmo salar*) and other marine organisms and sediments. *Marine Pollution Bulletin*, 45(1-12), 126-132. https://doi.org/10.1016/S0025-326X(01)00296-X
- 24. Soto, D. X., Trueman, C. N., Samways, K. M., Dadswell, M. J., & Cunjak, R. A. (2018). Ocean warming cannot explain synchronous declines in North American Atlantic salmon populations. *Marine Ecology Progress Series*, 601, 203-213. http://dx.doi.org/10.3354/meps12674
- 25. Stagg, R.M., Robinson, C., McIntosh, A.M., Moffat, C.F., & Bruno, D.W. (1998). The effects of the 'Braer'oil spill, Shetland Isles, Scotland, on P4501A in fanned Atlantic salmon (*Salmo salar*) and the common dab (*Limanda limanda*). *Marine environmental research*, 46(1-5), 301-306. http://dx.doi.org/10.1016/S0141-1136(98)00014-2
- 26. Stankevičiūtė, M., Sauliutė, G., Makaras, T., Markuckas, A., Virbickas, T., & Baršienė, J. (2018). Responses of biomarkers in Atlantic salmon (*Salmo salar*) following exposure to environmentally relevant concentrations of complex metal mixture (Zn, Cu, Ni, Cr, Pb, Cd). Part II. *Ecotoxicology*, 27, 1069-1086. http://dx.doi.org/10.1007/s10646-018-1960-2
- 27. Sundt-Hansen, L., Huisman, J., Skoglund, H., & Hindar, K. (2015). Farmed Atlantic salmon Salmo salar L. parr may reduce early survival of wild fish. *Journal of Fish Biology*, 86(6), 1699-1712. http://dx.doi.org/10.1111/jfb.12677
- 28. Thorstad, E.B., Whoriskey, F., Uglem, I., Moore, A., Rikardsen, A.H., & Finstad, B. (2012). A critical life stage of the Atlantic salmon *Salmo salar*: Behaviour and survival during the smolt *Appendix 11 Sensitivity Analyses 15 Atlantic salmon*

and initial post-smolt migration. *Journal of fish biology*, 81(2), 500-542. http://dx.doi.org/10.1111/j.1095-8649.2012.03370.x

- 29. Verspoor, E., Knox, D., & Marshall, S. (2016). Assessment of interbreeding and introgression of farm genes into a small Scottish Atlantic salmon *Salmo salar* stock: ad hoc samples—ad hoc results?. *Journal of Fish Biology*, 89(6), 2680-2696. http://dx.doi.org/10.1111/jfb.13173
- 30. Vollset, K.W., Krontveit, R.I., Jansen, P.A., Finstad, B., Barlaup, B. T., Skilbrei, O.T., Krkošek, M., Romunstad, P., Aunsmo, A., Jensen, A.J., & Dohoo, I. (2016). Impacts of parasites on marine survival of Atlantic salmon: A meta-analysis. *Fish and Fisheries*, 17(3), 714-730. http://dx.doi.org/10.1111/faf.12141

Literature Search

Web of Science search terms

("Salmo salar" OR "S. salar") AND ("angl*" OR "beam" OR "bottom trawl*" OR "by-catch" OR "dredge*" OR "fish*" OR "gear" OR "gillnet*" OR "hook*" OR "injury" OR "net*" OR "otter trawl*" OR "remov*" OR "aggregate*" OR "anchor*" OR "ballast" OR "barrier*" OR "beach*" OR "launch*" OR "moor*" OR "noise" OR "ship*" OR "steaming" OR "collision*" OR "construction" OR "electro*" OR "turbine*" OR "renewable*" OR "wave" OR "wind" OR "wind farm*" OR "anoxia" OR "copper" OR "current*" OR "deoxy*" OR "disease*" OR "disturbance" OR "endocrine disru*" OR "eutrophication" OR "exposure" OR "heavy metals" OR "hydrocarbon" OR "hypoxia OR litter*" OR "non-native*" OR "nitrate*" OR "nitrite*" OR "noise" OR "radionuclide" OR "nutrient*" OR "oil" OR "PAH*" OR "PCB*" OR "regime" OR "sedimentation" OR "silt*" OR "tributyltin" OR "turbid*")

Database

ISI Web of Science

Search date

7th February 2024 - 5,035 results

https://www.webofscience.com/wos/woscc/summary/a439c152-8f07-499e-b2f2-bd3f80ea38f7-d614734f/relevance/1

Search output and screening process

Workflow follows the Rapid Evidence Assessment. Web of Science only exports the first 1,000 results. The titles of these 1,000 papers were screened for relevance i.e. must describe salmon and mention of one of the listed sectors and/or pressures from MARESA. A second screening of abstracts was then conducted. The large number of initial results was due to the fact that salmon are widely used in aquaculture, however many of these reports were not of relevance to wild salmon under natural marine conditions. >4,000 pieces of literature were unscreened, hence the importance of the review papers and OSPAR assessment.

Outcome from screening

54 papers passed the first screening and 30 of these passed the second. These 30 papers formed the basis of the sensitivity assessment, supplemented with the latest reviews, OSPAR report and IUCN Red List assessments