

# 18. Turbot (*Scophthalmus maximus*)

## Sensitivity Assessment

**Table A11.18. Sensitivity assessment for Turbot (*Scophthalmus maximus*).** Associated sectors include activities related to offshore renewable energy (O), Fishing (F), or shipping (S). NR = not relevant, NA = not assessed, NEv = no evidence, H = high, M = medium, L = low, NS = not sensitive.

Pressures		Associated sector(s)	Resistance				Resilience				Sensitivity			
Classification	Pressure type		Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
Physical	Physical loss (to land or freshwater habitat)	O	L	H	H	H	M	M	M	M	M	M	M	M
	Physical change (to another seabed type)	O, F	L	H	H	H	M	M	M	M	M	M	M	M
	Physical change (to another sediment type)	O, F	L	H	H	H	M	M	M	M	M	M	M	M

Pressures		Associated sector(s)	Resistance				Resilience				Sensitivity			
Classification	Pressure type		Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
	Habitat structure change-removal of substratum (extraction)	O	L	H	H	H	M	M	M	M	M	M	M	M
	Abrasion/disturbance of substratum surface or seabed	O, F	M	M	M	M	H	NR	NR	NR	L	NR	NR	NR
	Penetration or disturbance of substratum subsurface	O, F	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Changes in suspended solids (water clarity)	O, F	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR

Pressures		Associated sector(s)	Resistance				Resilience				Sensitivity			
Classification	Pressure type		Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
Physical	Smothering and siltation changes (light)	O	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Smothering and siltation changes (heavy)	O	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Underwater noise	O, F, S	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Electromagnetic energy	O	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Barrier to species movement	O, F	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Death or injury by collision	O, F, S	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Hydrological	Water flow changes	O	M	M	H	M	M	M	M	M	M	M	H	M

Appendix 11 Sensitivity Analyses - 18 Turbot

Pressures		Associated sector(s)	Resistance				Resilience				Sensitivity			
Classification	Pressure type		Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
Chemical	Transition elements & organo-metal contamination	O, F, S	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Hydrocarbon & PAH contamination	O, F, S	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Synthetic compound contamination	O, F, S	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Introduction of other substances	O, F, S	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
	Deoxygenation	O	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR
Biological	Introduction or spread of invasive non-indigenous species	O, F, S	NEv	NR	NR	NR	NEv	NR	NR	NR	NEv	NR	NR	NR

Pressures		Associated sector(s)	Resistance				Resilience				Sensitivity			
Classification	Pressure type		Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC
	Removal of target species	F	L	H	H	H	M	H	H	H	M	H	H	H
	Removal of non-target species	F	L	H	H	H	M	H	H	H	M	H	H	H

## References for sensitivity assessment

1. Basaran, F., & Samsun, N. (2004). Survival rates of black sea turbot (*Psetta maxima maeotica*, L. 1758) broodstock captured by gill nets from different depths and their adaptation culture conditions. *Aquaculture International*, 12, 321-331.  
<https://doi.org/10.1023/B:AQUI.0000036183.39217.2a>
2. Burel, C., Person-Le Ruyet, J., Gaumet, F., Le Roux, A., Severe, A., & Boeuf, G. (1996). Effects of temperature on growth and metabolism in juvenile turbot. *Journal of Fish Biology*, 49(4), 678-692. <https://doi.org/10.1006/jfbi.1996.0196>
3. Caputo, V., Candi, G., Colella, S., & Arneri, E. (2001). Reproductive biology of turbot (*Psetta maxima*) and brill (*Scophthalmus rhombus*) (Teleostei, Pleuronectiformes) in the Adriatic Sea. *Italian Journal of Zoology*, 68(2), 107-113.  
<https://doi.org/10.1080/11250000109356393>
4. Cardinale, M., Chanet, B., Martínez Portela, P., Munroe, T.A., Nimmegeers, S., Shlyakhov, V., Turan, C. & Vansteenbrugge, L. (2021). *Scophthalmus maximus*. *The IUCN Red List of Threatened Species* 2021: e.T198731A144939322.  
<https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T198731A144939322.en>
5. Ellis, T., Howell, B. R., & Hayes, J. (1997). Morphological differences between wild and hatchery-reared turbot. *Journal of Fish Biology*, 50(5), 1124-1128.  
<https://doi.org/10.1006/jfbi.1996.0364>
6. Florin, A. B., & Höglund, J. (2007). Absence of population structure of turbot (*Psetta maxima*) in the Baltic Sea. *Molecular Ecology*, 16(1), 115-126.  
<https://doi.org/10.1111/j.1365-294X.2006.03120.x>
7. Haynes, P. S., Brophy, D., McGrath, D., O'Callaghan, R., Comerford, S., & Casburn, P. (2010). Annual and spatial variation in the abundance length and condition of juvenile turbot (*Psetta maxima* L.) on nursery grounds on the west coast of Ireland: 2000–2007. *Journal of Sea Research*, 64(4), 494-504.  
<https://doi.org/10.1016/j.seares.2010.01.009>
8. Imsland, A. K., Sunde, L. M., Folkvord, A., & Stefansson, S. O. (1996). The interaction of temperature and fish size on growth of juvenile turbot. *Journal of Fish Biology*, 49(5), 926-940. <https://doi.org/10.1111/j.1095-8649.1996.tb00090.x>
9. Kerambrun, E., Henry, F., Rabhi, K., & Amara, R. (2014). Effects of chemical stress and food limitation on the energy reserves and growth of turbot, *Scophthalmus*

*maximus*. *Environmental Science and Pollution Research*, 21, 13488-13495.  
<https://doi.org/10.1007/s11356-014-3281-1>

10. Laikre, L., Palm, S., & Ryman, N. (2005). Genetic population structure of fishes: implications for coastal zone management. *AMBIO: A Journal of the Human Environment*, 34(2), 111-119. [https://doi.org/10.1639/0044-7447\(2005\)034\[0111:GPSOFI\]2.0.CO;2](https://doi.org/10.1639/0044-7447(2005)034[0111:GPSOFI]2.0.CO;2)
11. Lennquist, A., Hilvarsson, A., & Förlin, L. (2010). Responses in fish exposed to medetomidine, a new antifouling agent. *Marine Environmental Research*, 69, S43-S45.  
<https://doi.org/10.1016/j.marenvres.2010.01.004>
12. Maruya, K. A., Vidal-Dorsch, D. E., Bay, S. M., Kwon, J. W., Xia, K., & Armbrust, K. L. (2012). Organic contaminants of emerging concern in sediments and flatfish collected near outfalls discharging treated wastewater effluent to the Southern California Bight. *Environmental Toxicology and Chemistry*, 31(12), 2683-2688.  
<https://doi.org/10.1002/etc.2003>
13. Mhadhbi, L., Boumaiza, M., & Beiras, R. (2010). A standard ecotoxicological bioassay using early life stages of the marine fish *Psetta maxima*. *Aquatic living resources*, 23(2), 209-216.  
<https://doi.org/10.1051/alr/2010014>
14. Panayotova, M., Raykov, V., Maximov, V., Radu, G., & Anton, E. (2012). Distribution, abundance and population structure of Turbot (*Psetta maxima* L.) in the Bulgarian and Romanian Black Sea area in spring 2010. *Comptes rendus de l'Académie bulgare des Sciences*, 65(1), 63-66.
15. Person-Le Ruyet, J., Lacut, A., Le Bayon, N., Le Roux, A., Pichavant, K., & Quémener, L. (2003). Effects of repeated hypoxic shocks on growth and metabolism of turbot juveniles. *Aquatic Living Resources*, 16(1), 25-34.  
[https://doi.org/10.1016/S0990-7440\(02\)00002-5](https://doi.org/10.1016/S0990-7440(02)00002-5)
16. Pichavant, K., Person-Le-Ruyet, J., Bayon, N. L., Severe, A., Roux, A. L., & Boeuf, G. (2001). Comparative effects of long-term hypoxia on growth, feeding and oxygen consumption in juvenile turbot and European sea bass. *Journal of Fish Biology*, 59(4), 875-883.  
<https://doi.org/10.1006/jfbi.2001.1702>

17. Polak-Juszczak, L. (2015). Selenium and mercury molar ratios in commercial fish from the Baltic Sea: additional risk assessment criterion for mercury exposure. *Food control*, 50, 881-888.
18. <https://doi.org/10.1016/j.foodcont.2014.10.046>
19. Pranovi, F., Monti, M. A., Caccin, A., Brigolin, D., & Zucchetto, M. (2015). Permanent trawl fishery closures in the Mediterranean Sea: An effective management strategy?. *Marine Policy*, 60, 272-279. <https://doi.org/10.1016/j.marpol.2015.07.003>
20. Rijnsdorp, A. D., van Overzee, H. M., & Poos, J. J. (2012). Ecological and economic trade-offs in the management of mixed fisheries: a case study of spawning closures in flatfish fisheries. *Marine Ecology Progress Series*, 447, 179-194.  
<https://doi.org/10.3354/meps09519>
21. Sparrevohn, C. R., & Støttrup, J. G. (2008). Diet, abundance, and distribution as indices of turbot (*Psetta maxima* L.) release habitat suitability. *Reviews in Fisheries Science*, 16(1-3), 338-347. <https://doi.org/10.1080/10641260701686846>
22. Ustups, D., Bergström, U., Florin, A. B., Kruze, E., Zilniece, D., Elferts, D., Knospina, E., & Uzars, D. (2016). Diet overlap between juvenile flatfish and the invasive round goby in the central Baltic Sea. *Journal of Sea Research*, 107, 121-129.  
<https://doi.org/10.1016/j.seares.2015.06.021>
23. Valente, L. M., Moutou, K. A., Conceicao, L. E., Engrola, S., Fernandes, J. M., & Johnston, I. A. (2013). What determines growth potential and juvenile quality of farmed fish species?. *Reviews in Aquaculture*, 5, S168-S193.  
<https://doi.org/10.1111/raq.12020>
24. van Bussel, C. G., Mahlmann, L., Kroeckel, S., Schroeder, J. P., & Schulz, C. (2013). The effect of high ortho-phosphate water levels on growth, feed intake, nutrient utilization and health status of juvenile turbot (*Psetta maxima*) reared in intensive recirculating aquaculture systems (RAS). *Aquacultural engineering*, 57, 63-70.  
<https://doi.org/10.1016/j.aquaeng.2013.08.003>
25. van Bussel, C. G., Schroeder, J. P., Wuertz, S., & Schulz, C. (2012). The chronic effect of nitrate on production performance and health status of juvenile turbot (*Psetta maxima*). *Aquaculture*, 326, 163-167.  
<https://doi.org/10.1016/j.aquaculture.2011.11.019>
26. van der Hammen, T., Poos, J. J., van Overzee, H. M., Heessen, H. J., Magnusson, A., & Rijnsdorp, A. D. (2013). Population ecology of turbot and brill: What can we



learn from two rare flatfish species?. *Journal of Sea Research*, 84, 96-108.

<https://doi.org/10.1016/j.seares.2013.07.001>

27. Wang, X. A., Ma, A., Huang, Z., Sun, Z., Cui, W., Qu, J., & Yu, H. (2019). Estimation of genetic parameters for upper thermal tolerances and growth-related traits in turbot *Scophthalmus maximus*. *Journal of oceanology and limnology*, 37(5), 1736-1745.

<https://doi.org/10.1007/s00343-019-7267-1>

28. Weltzien, F. A., Planas, M., & Fyhn, H. J. (1999). Temperature dependency of early growth of turbot (*Scophthalmus maximus* L.) and its implications for developmental progress. *Journal of experimental marine biology and ecology*, 242(2), 201-210.

[https://doi.org/10.1016/S0022-0981\(99\)00099-4](https://doi.org/10.1016/S0022-0981(99)00099-4)

Literature search

### Web of Science search terms

AB=("Scophthalmus maximus" OR "S. maximus") AND AB=("angl\*" OR "beam" OR "bottom trawl\*" OR "by-catch" OR "dredge\*" OR "fish\*" OR "gear" OR "gillnet\*" OR "hook\*" OR "injury" OR "net\*" OR "otter trawl\*" OR "remov\*" OR "aggregate\*" OR "anchor\*" OR "ballast" OR "barrier\*" OR "beach\*" OR "launch\*" OR "moor\*" OR "noise" OR "ship\*" OR "steaming" OR "collision\*" OR "construction" OR "electro\*" OR "turbine\*" OR "renewable\*" OR "wave" OR "wind" OR "wind farm\*" OR "anoxia" OR "copper" OR "current\*" OR "deoxy\*" OR "disease\*" OR "disturbance" OR "endocrine disru\*" OR "eutrophication" OR "exposure" OR

"heavy metals" OR "hydrocarbon" OR "hypoxia" OR "litter\*" OR "non-native\*" OR "nitrate\*" OR "nitrite\*" OR "noise" OR "radionuclide" OR "nutrient\*" OR "oil" OR "PAH\*" OR "PCB\*" OR "regime" OR "sedimentation" OR "silt\*" OR "tributyltin" OR "turbid\*")

## **Database**

ISI Web of Science

<https://www.webofscience.com/wos/woscc/summary/fa3ef578-304a-4a10-9430-eb1fea555b6e-cb38492f/relevance/1>

## **Search date**

28th February 2023 - 856 results

## **Search output and screening process**

Abstracts screened for relevance i.e. must describe turbot and mention of one of the listed sectors and/or pressures from MARESA. Workflow follows the Rapid Evidence Assessment approach. The title and all auxiliary information (including abstract) were downloaded from ISI Web of Science in a .ris and excel format. In Excel, abstracts were read and listed to either pass or fail the initial screening process with a reason provided.

## **Outcome from screening**

As turbot is grown in aquaculture in some areas, many of the resultant papers were related to non-natural conditions. 26 papers passed the screening and formed the basis of the sensitivity assessment, supplemented with the latest IUCN Red List assessment (2021).