34. Kelp forest

Sensitivity Assessment

Sensitivity scores for 12 kelp level 5 biotopes, as described under The Marine Habitat Classification for Britain and Ireland (JNCC, 2022), within the Celtic Sea were obtained from MarLIN website (www.marlin.ac.uk). See case report (Appendix 10) for details of kelp biotopes used to assess this feature. The resistance, resilience and sensitivity scores for each pressure comprise those scores for the kelp biotope(s) most sensitive to that pressure.

Table A11.34. Sensitivity assessment for kelp forest. Associated sectors include activities related to offshore renewable energy (O), Fishing (F), or shipping (S). NR = not relevant, NA = not assessed, NEv = no evidence, H = high, M = medium, L = low, VL = very low, N = none, NS = not sensitive. Refs = References.

Pressures		Associated	Resistance				Resilier	nce			Sensitiv	ity		Biotope associated	Refs	
Classification	Pressure type	sector(s)	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	Score	QoE	AoE	DoC	with score	incis
Physical	Physical loss (to land or freshwater habitat)	0	N	Н	Н	Н	VL	Н	Н	Н	Н	Н	Н	Н	All	1-12
	Physical change (to another seabed type)	O, F	N	н	Н	Н	VL	Н	Н	Н	Н	Н	Н	Н	AII	1-12
	Physical change (to another sediment type)	O, F	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	All	1-12

	Habitat structure change-removal of substratum (extraction)	0	N	Н	Н	н	M	Н	Н	Н	M	Н	Н	н	5, 6	3, 11
	Abrasion/disturbance of substratum surface or seabed	O, F	L	Н	Н	Н	М	Н	Н	Н	М	Н	Н	Н	3, 4, 5, 6, 7, 8, 9, 10, 12	3-8, 10- 12
	Penetration or disturbance of substratum subsurface	O, F	NR	All	1-12											
Physical	Changes in suspended solids (water clarity)	O, F	N	Н	Н	н	М	Н	Н	Н	М	Н	Н	Н	4, 5, 6, 7, 8, 9	3, 4, 7, 10-12
	Smothering and siltation changes (light)	О	М	L	NR	NR	Н	н	L	Н	L	L	L	L	9	7
	Smothering and siltation changes (heavy)	0	L	М	Н	Н	Н	Н	Н	Н	L	М	Н	Н	2, 3, 4, 7, 8, 9, 10, 11, 12	1, 2, 4- 8, 10, 12

	Underwater noise	O, F, S	NR	All	1-12											
	Electromagnetic energy	О	NR	All	1-12											
	Barrier to species movement	O, F	NR	All	1-12											
	Death or injury by collision	O, F, S	NR	All	1-12											
Hydrological	Water flow changes	0	М	L	NR	NR	Н	Н	L	Н	L	L	NR	NR	1	9
Chemical	Transition elements & organo-metal contamination	O, F, S	NA	NR	NR	NR	NA	NR	NR	NR	NA	NR	NR	NR	All	1-12
	Hydrocarbon & PAH contamination	O, F, S	NA	NR	NR	NR	NA	NR	NR	NR	NA	NR	NR	NR	All	1-12
	Synthetic compound contamination	O, F, S	NA	NR	NR	NR	NA	NR	NR	NR	NA	NR	NR	NR	All	1-12

	Introduction of other substances	O, F, S	NA	NR	NR	NR	NA	NR	NR	NR	NA	NR	NR	NR	All	1-12
	Deoxygenation	О	М	Н	М	Н	Н	Н	М	Н	L	Н	М	Н	2, 5, 6	1, 3, 11
Biological	Introduction or spread of invasive non-indigenous species	0	L	L	NR	NR	VL	Н	Н	Н	Н	L	NR	NR	2, 3, 9	1, 7, 8
	Removal of target species	O, F, S	N	Н	Н	Н	М	Н	Н	Н	М	Н	Н	Н	1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12	
	Removal of non-target species	F	L	н	н	н	М	Н	н	н	М	Н	Н	Н	All	1-12

References for kelp forest sensitivity assessment

- Hill, J.M., Tyler-Walters, H.,, Burdett, E.G. & Jasper, C. (2023). Laminaria digitata on moderately exposed sublittoral fringe rock. In Tyler-Walters H. (ed) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line]. Plymouth: Marine Biological Association of the United Kingdom. [cited 29-04-2024].
 - Available from: https://www.marlin.ac.uk/habitats/detail/93
- Jasper, C., Lloyd, K.A., & Mardle, M.J. (2022). Saccharina latissima on very sheltered infralittoral rock. In Tyler-Walters H. (ed) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line].
 Plymouth: Marine Biological Association of the United Kingdom. [cited 29-04-2024].
 - Available from: https://www.marlin.ac.uk/habitats/detail/89
- Stamp, T.E., Burdett, E.G., Tyler-Walters, H., & Lloyd, K.A. (2023a). Laminaria hyperborea on tide-swept, infralittoral mixed substrata. In Tyler-Walters H. (ed) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line]. Plymouth: Marine Biological Association of the United Kingdom. [cited 29-04-2024].
 - Available from: https://www.marlin.ac.uk/habitats/detail/1045
- Stamp, T.E., Hiscock, K., Garrard, S.L., Burdett, E.G. & Tyler-Walters, H.
 (2023b). Laminaria hyperborea forest with a faunal cushion (sponges and polyclinids) and foliose red seaweeds on very exposed upper infralittoral rock. In Tyler-Walters H. and Hiscock K. (eds) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line].

 Plymouth: Marine Biological Association of the United Kingdom. [cited 29-04-2024].
 - Available from: https://www.marlin.ac.uk/habitats/detail/44
- Stamp, T.E., Hiscock, K. & Williams, E., Lloyd, K.A.,, Mardle, M.J., & Tyler-Walters, H., (2022a). Saccharina latissima and/or Saccorhiza polyschides on exposed infralittoral rock. In Tyler-Walters H. (ed) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line].

- Plymouth: Marine Biological Association of the United Kingdom. [cited 29-04-2024]. Available from: https://www.marlin.ac.uk/habitats/detail/237
- Stamp, T.E., Williams, E., Lloyd, K.A., Mardle, M.J., & Tyler-Walters, H.
 (2022b). Saccorhiza polyschides and other opportunistic kelps on disturbed sublittoral fringe rock. In Tyler-Walters H. (ed) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line].

 Plymouth: Marine Biological Association of the United Kingdom. [cited 29-04-2024].

Available from: https://www.marlin.ac.uk/habitats/detail/118

Available from: https://www.marlin.ac.uk/habitats/detail/141

- Stamp, T.E., Lloyd, K.A., & Mardle, M.J. (2022c). Mixed Laminaria hyperborea
 and Saccharina latissima park on sheltered lower infralittoral rock. In TylerWalters H. (ed) Marine Life Information Network: Biology and Sensitivity Key
 Information Reviews, [on-line]. Plymouth: Marine Biological Association of
 the United Kingdom. [cited 29-04-2024].
- Stamp, T.E., Marshall, C.E., Williams, E., Lloyd, K.A., & Mardle, M.J. (2022d).
 Laminaria digitata, ascidians and bryozoans on tide-swept sublittoral fringe
 rock. In Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews*, [on-line].
 Plymouth: Marine Biological Association of the United Kingdom. [cited 29 04-2024]. Available from: https://www.marlin.ac.uk/habitats/detail/273
- Stamp, T.E., Tyler-Walters, H., & Burdett, E.G. (2023c). Alaria esculenta on exposed sublittoral fringe bedrock. In Tyler-Walters H. and Hiscock K. (eds) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line]. Plymouth: Marine Biological Association of the United Kingdom. [cited 29-04-2024].

Available from: https://www.marlin.ac.uk/habitats/detail/165

10. Stamp, T.E., Tyler-Walters, H., & Burdett, E.G. (2023d). Laminaria hyperborea and foliose red seaweeds on moderately exposed infralittoral rock. In Tyler-Walters H. (ed) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line]. Plymouth: Marine Biological Association of

the United Kingdom. [cited 29-04-2024].

Available from: https://www.marlin.ac.uk/habitats/detail/292

11. Stamp, T.E., Tyler-Walters, H., Burdett, E.G. & Lloyd, K.A. (2023e). Laminaria hyperborea on tide-swept, infralittoral rock. In Tyler-Walters H. (ed) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line]. Plymouth: Marine Biological Association of the United Kingdom. [cited 29-04-2024].

Available from: https://www.marlin.ac.uk/habitats/detail/1044

12. Tyler-Walters, H., Stamp, T.E., Burdett, E.G. & Lloyd, K.A. (2023). *Laminaria hyperborea* with dense foliose red seaweeds on exposed infralittoral rock. In Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews,* [on-line]. Plymouth: Marine Biological Association of the United Kingdom. [cited 29-04-2024].

Available from: https://www.marlin.ac.uk/habitats/detail/171