
WEST OF SCOTLAND, ROCKALL AND NORTH SEA ANGLERFISH



Sub-areas 4 and 6 and Division 3.a

ADVICE FOR 2024


ICES advises that when the MSY approach is applied, catches should be no more than 9,881 t in each of the years 2023 and 2024.



Assessment area

KEY POINTS

ICES Assessment Category	3 (rfb rule)	
Management Plan	NSMAP (target species)	
Advice Basis	ICES MSY approach	Catch=9,881 t
Ranges	F _{MSY} Range Lower	Not available
	F _{MSY}	Not available
	F _{MSY} Range Upper	Not available
Landing Obligation	From 2019	No derogations or exemptions apply




MANAGEMENT IN 2023

- Two species of Lophius occur in this area. Black anglerfish (Lophius budegassa) usually make up around 10% of the international landings but in some years this may be as much as 30%. The two species are assessed and managed as a single stock.
- There are three TAC areas that cover anglerfish in this area:
 - Union and UK waters of areas 2.a and 4
 - Norwegian waters of area 4
 - O Union, UK and international waters of areas 6, 5.b, 12, 14.

Ireland only has a quota for the 6, 5.b, 12, 14 area and generally relies on swaps to land its catches. France has the largest EU quota but usually takes less than a third of this. The UK has full quota uptake and is responsible for the vast majority of the international landings.

2023 Quota Allocations for area 6, 5.b, 12 and 14



• There is a special condition that states that up to 20% of the area 6, 5b, 12, 14 quota may be fished in UK and Union waters of 2a and 4.

KEY STOCK CONSIDERATIONS

- The advice is based on a survey index trend (decreasing) together with a fishing pressure proxy (F is above the reference point), a biomass safeguard (biomass is above the reference point) and precautionary multiplier. This would result in a reduction in catch advice of 36% but a stability clause limits the reduction in advice to 30%.
- Irish vessels mainly catch anglerfish in mixed fisheries with various gadoids and megrim near the Stanton Banks; with hake and other gadoids along the continental slope; and with megrim to the south of Rockall.
- Discard levels in this stock are estimated to be low (1.2% in 2021).