

# Test Version

2020-04-06

*Load packages*

*Load and transform shapefile*

```
###1.Load seabed habitats shapefile
```

```
sb2019 <-readOGR(paste0(Data_location,"Template_Ecosystem/data/MSFD"),"MSFD_BBHT_Ireland2019")
```

```
## OGR data source with driver: ESRI Shapefile
## Source: "C:\Template_Ecosystem\data\MSFD", layer: "MSFD_BBHT_Ireland2019"
## with 100401 features
## It has 7 fields
```

```
#####
####"Mercator" Coordinate Reference Systems used in the habitat shapefile####
#####
```

```
###2. Transform to "WGS84" Coordinate Reference Systems
```

```
#####
#### spTransform() function from sp package used to transform from "Mercator" to "WGS84" CRS##
#####
```

```
sb2019 <-
spTransform(sb2019,
"+proj=longlat +datum=WGS84 +no_defs +ellps=WGS84 +towgs84=0,0,0")
```

## Seabed Habitats Mapping around Ireland

### Summary

Biozone	area_km2	perc
Atlantic upper bathyal	0	0%
Atlanto-Mediterranean mid bathyal	0	0%
Deep circalittoral	23256	97.9%
Infralittoral	0	0%
Shallow circalittoral	167	0.7%
Unclassified	343	1.4%

Substrate	area_km2	perc
Coarse sediment	474	2%
Coarse substrate	1227	5.2%
Fine mud	56	0.2%
Muddy sand	4650	19.6%
Rock	75	0.3%
Rock or other hard substrata	881	3.7%
Sand	15929	67%
Sandy mud	48	0.2%
sandy Mud/muddy Sand	83	0.3%
Seabed	0	0%
Unclassified substrate	0	0%
Unclassified substrate	343	1.4%

Folk_5	area_km2	perc
Coarse sediment	1701	7.2%
Mud to muddy Sand	4837	20.4%
Rock	956	4%
Sand	15929	67%
Unclassified	343	1.4%

EUNIS	area_km2	perc
A3: Infralittoral rock and other hard substrata	0	0%
A4: Circalittoral rock and other hard substrata	956	4%
A5.13: Infralittoral coarse sediment	0	0%
A5.14: Circalittoral coarse sediment	1	0%
A5.15: Deep circalittoral coarse sediment	1700	7.2%
A5.25 or A5.26: Circalittoral fine sand or Circalittoral muddy sand	4	0%
A5.27: Deep circalittoral sand	15925	67%
A5.35: Circalittoral sandy mud	25	0.1%
A5.37: Deep circalittoral mud	4812	20.2%
A6.5: Deep-sea mud	0	0%
A6: Deep-sea bed	0	0%
Unclassified	343	1.4%

MSFD_BBHT	area_km2	perc
Circalittoral coarse sediment	1	0%
Circalittoral mud	25	0.1%
Circalittoral rock and biogenic reef	137	0.6%
Circalittoral sand	4	0%
Infralittoral coarse sediment	0	0%
Infralittoral rock and biogenic reef	0	0%
Offshore circalittoral coarse sediment	1700	7.2%
Offshore circalittoral mud	4812	20.2%
Offshore circalittoral rock and biogenic reef	819	3.4%
Offshore circalittoral sand	15925	67%
Unclassified	343	1.4%
Upper bathyal sediment	0	0%


MSFD_BBHT	area_km2	perc
Upper bathyal sediment or Upper bathyal rock and biogenic reef	0	0%




# Copernicus Marine Service Data


## Ocean Physics Realanalysis

### Temperature




### Temperature







## Sea floor potential temperature



## Sea floor potential temperature




## Sea floor potential temperature







## Salinity




## Salinity






## Ocean BioGeoChemistry NON ASSIMILATIVE Hindcast


### Mass Concentration of Chlorophyll in Sea Water




### Mass Concentration of Chlorophyll in Sea Water




## Mass Concentration of Chlorophyll in Sea Water




### Mole Concentration of Phytoplankton Expressed as Carbon in Sea Water



### Mole Concentration of Phytoplankton Expressed as Carbon in Sea Water




## In situ Concentration of Phytoplankton Expressed as Carbon in Sea Water




# An operational zooplankton data service

## Observed Counts




## Data Interpolation and Variational Analysis

Acartia







*Calanus finmarchicus*




*Calanus\_helgolandicus*




*Metridia\_lucens*



*Temora\_longicornis*



## Large\_copepods



## Small\_copepods

