

Multi-resolution Spatial Statistics

lao Hoi (Isaac) Leong Supervisor: Dr Tingjin Chu

Motivation

Many spatial datasets are collected at incompatible locations and resolutions being the average of the local neighbourhood.

If this mismatch is ignored, simple upsampling can blur spatial structure and give uncertainty statements that are overconfident. This is especially common in astronomy, where different instruments and beam sizes naturally produce measurements on different supports.

This project is motivated by scientific settings where the covariate is intrinsically aggregated by the measurement process, so the analysis should respect the support on which it is observed. One example is the following two datasets as shown in Figure 1 captured for astronomical research.




Figure 1: Observation locations for two metallicity datasets with support mismatch: MGS is sampled at finer resolution, while GAL is measured on a much coarser footprint. Both record gas-phase metallicity across the same galaxy using different indicators.

A recent research by Zheng et al. (2025) proposed to use Basic Area Unit (BAU) level modelling to solve the problem. We followed similar ideas, but employ a frequentist approach in contrast to theirs Bayesian approach.

Background

Consider the following spatial process

$$y(\mathbf{s}) = \beta_0 + \beta_1 d(\mathbf{s}) + \beta_2 x(\mathbf{s}) + \varepsilon(\mathbf{s}), \quad (1)$$

where both $y(\cdot)$ and $d(\cdot)$ are observed at $\{\mathbf{s}_1, \dots, \mathbf{s}_n\}$, while $x(\cdot)$ is observed at other locations at $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ with different resolution in a 100×100 domain. We assume that $\{\mathbf{s}_1, \dots, \mathbf{s}_n\}$ has a resolution of 5×5 while $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ has a resolution of 3×3 for simplicity.

We divide the 3×3 area to 1×1 basic area unit (BAU) by modelling the coarsing 3×3 resolution as an average of the BAU. Then $\mathbf{x}_u = [x(\mathbf{u}_1) \ x(\mathbf{u}_2) \ \dots \ x(\mathbf{u}_m)]^\top$ and similar for \mathbf{x}_v which denotes $x(\cdot)$ at all BAUs, we can put it in matrix form as,

$$\mathbf{x}_u = W\mathbf{x}_v, \text{ where } W = \underbrace{\begin{bmatrix} \frac{1}{9} & \dots & \frac{1}{9} & 0 & 0 & 0 & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \frac{1}{9} & \dots & \frac{1}{9} & 0 & \dots & \dots & \dots \\ \vdots & \vdots & \vdots & 0 & 0 & 0 & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots & \frac{1}{9} & \dots & \frac{1}{9} \end{bmatrix}}_{\mathbb{R}^{m \times (9 \times m)}}$$

Then model $x(\cdot)$ at BAU level, as $x(\mathbf{v}) = \mathbf{r}(\mathbf{v})^\top \boldsymbol{\alpha} + \eta(\mathbf{v})$ would give the following distribution,

$$\mathbf{x}_v \sim \text{MVN}(\alpha_0 \mathbf{1}, \Sigma_\eta(\boldsymbol{\xi})) \implies \mathbf{x}_u \sim \text{MVN}(\alpha_0 \mathbf{1}, W\Sigma_\eta(\boldsymbol{\xi})W^\top).$$




Figure 2: Approach for going from observed aggregated data and underlying BAU grids.

MLE

Since the aggregated data \mathbf{x}_u is observed, we can use likelihood methods to estimate the unknown parameters α_0 and $\boldsymbol{\xi}$. Sepcifically, the log-likelihood is:

$$L(\alpha_0, \boldsymbol{\xi}) = -\frac{m}{2} \log(2\pi) - \frac{1}{2} \log(|W\Sigma_\eta(\boldsymbol{\xi})W^\top|) - \frac{1}{2} (\mathbf{x}_u - \alpha_0 W\mathbf{1})^\top (W\Sigma_\eta(\boldsymbol{\xi})W^\top)^{-1} (\mathbf{x}_u - \alpha_0 W\mathbf{1}) \quad (2)$$

Then, by setting the gradient with respect to α_0 we obtain the following estimator for $\hat{\alpha}_0(\boldsymbol{\xi})$:

$$\hat{\alpha}_0(\boldsymbol{\xi}') = \frac{\mathbf{1}^\top (W\Sigma_\eta(\boldsymbol{\xi})W^\top)^{-1} \mathbf{x}_u}{\mathbf{1}^\top (W\Sigma_\eta(\boldsymbol{\xi})W^\top)^{-1} \mathbf{1}}$$

Then, assume the covariance function admits a separable variance (partial sill) parameter σ^2 , then, the covaraince matrix $\Sigma(\boldsymbol{\xi})$ can be expressed as

$$\Sigma_\eta(\boldsymbol{\xi}) = \sigma^2 R_\eta(\boldsymbol{\xi}')$$

where $\boldsymbol{\xi}'$ is just $\boldsymbol{\xi}$ with σ^2 excluded. Then, by recognising the same trick as above, we can rewrite the log-likelihood in Equation (2) as:

$$L(\alpha_0, \sigma^2, \boldsymbol{\xi}') = -\frac{m}{2} \log(2\pi) - \frac{1}{2} \log(|\sigma^2 W R_\eta(\boldsymbol{\xi}') W^\top|) - \frac{1}{2} (\mathbf{x}_u - \alpha_0 \mathbf{1})^\top (\sigma^2 W R_\eta(\boldsymbol{\xi}') W^\top)^{-1} (\mathbf{x}_u - \alpha_0 \mathbf{1}) \quad (3)$$

Then by setting the gradient of Equation (3) with respect to σ^2 , we obtain the following estimator for σ^2 :

$$\hat{\sigma}^2(\boldsymbol{\xi}') = \frac{1}{m} (\mathbf{x}_u - \alpha_0 \mathbf{1})^\top (W R_\eta(\boldsymbol{\xi}') W^\top)^{-1} (\mathbf{x}_u - \alpha_0 \mathbf{1})$$

Hence, we obtain the final concentrated log-likelihood as,

$$L(\boldsymbol{\xi}') = -\frac{m}{2} \log(2\pi) - \frac{m}{2} - \frac{m}{2} \log(\hat{\sigma}^2(\boldsymbol{\xi}')) - \frac{1}{2} \log|Q(\boldsymbol{\xi}')|$$

Then, we use optimisation algorithm such as Newton's Method to optimise with respect to $\boldsymbol{\xi}'$

Restricted Maximum Likelihood

MLE method typically have a downward bias for variance-related estimates. One way to mitigate is to use restricted maximum likelihood (REML).

Mathematically, we just need to find $K \in \mathbb{R}^{m \times (m-1)}$ such that $\mathbf{1}_m \in \ker K^\top$, and w.l.o.g assume K is orthonormal. Then, define $\mathbf{x}_u^* = K^\top \mathbf{x}_u$, we obtain that the expectation on the new variable would be zero,

$$\mathbb{E}[\mathbf{x}_u^*] = K^\top \mathbb{E}[\mathbf{x}_u] = \alpha_0 K^\top \mathbf{1} = 0.$$

Now, using the same procedure as outlined by LaMotte (2007), it can be shown that such restricted log-likelihood function is independent of the choice of matrix K . Specifically, it can be written in the form of,

$$L(\boldsymbol{\xi}) = -\frac{m-1}{2} \log(2\pi) - \frac{1}{2} (\mathbf{x}_u - \hat{\alpha}_0 \mathbf{1}_m)^\top V_\eta(\boldsymbol{\xi})^{-1} (\mathbf{x}_u - \hat{\alpha}_0 \mathbf{1}_m) - \frac{1}{2} \log(\mathbf{1}_m^\top V_\eta(\boldsymbol{\xi})^{-1} \mathbf{1}_m) - \frac{1}{2} \log|V_\eta(\boldsymbol{\xi})| + \frac{1}{2} \log(m)$$

where $\hat{\alpha}_0 \triangleq (\mathbf{1}_m^\top V_\eta(\boldsymbol{\xi})^{-1} \mathbf{1}_m)^{-1} \mathbf{1}_m^\top V_\eta(\boldsymbol{\xi})^{-1} \mathbf{x}_u$. The precise expression is an algebraic artifact but appears to be the same as above. Then, the estimation of the parameters $\boldsymbol{\xi}'$ can simply be done by maximising the similar concentrated likelihood.

Simulation Results

Across all simulations, both MLE and REML recover the true parameters reasonably well. REML shows *systematically smaller bias* for the covariance parameters σ^2 and ϕ , but its slightly larger SD means the RMSE can be similar or even a bit larger, suggesting the error is mainly variance-driven.

The range parameter ϕ is consistently the hardest to estimate, with the largest SD/RMSE across settings. When extending REML simulations, the estimators largely stabilise; increasing number of simulations, R , mainly reduces Monte Carlo noise in the reported metrics rather than the estimator's inherent sampling variability.

MLE Monte Carlo summaries




Figure 3: Monte Carlo summaries for MLE estimators across num. of simulations (R) (100,300,500): bias (dashed line = 0), standard deviation, and RMSE for σ^2 , ϕ , and α_0 .

REML Monte Carlo summaries




Figure 4: Monte Carlo summaries for the REML estimator as R increases (up to $R = 1000$): bias (dashed line at 0), standard deviation, and RMSE for σ^2 , ϕ , and α_0 .

Kriging

Now, in order to the result at correct resolution, we first need to perform kriging to obtain the best linear unbiased predictor (BLUP) with respect to MSE based on the estimated model.

Ultimately, we would like to perform kriging at resolution the same as $y(\cdot)$ at location $\{\mathbf{s}_1, \dots, \mathbf{s}_n\}$ with resolution $k \times k$. But, we will first focus on performing kriging at BAU level.




Figure 5: Kriging to the new locations and resolution.

As shown in Figure 5, we first need to krig for every BAU in 5×5 neighbourhood at new locations. This essentially needs to solve the following optimisation problem.

$$\begin{aligned} &\text{minimise} \quad \mathbb{E} \left[\left(x(\mathbf{v}_0) - \left(\lambda_0 + \sum_{i=1}^m \lambda_i x_u(\mathbf{u}_i) \right) \right)^2 \right] \\ &\text{subject to} \quad \mathbb{E}[x(\mathbf{v}_0)] = \mathbb{E}[\hat{x}(\mathbf{v}_0)] \end{aligned}$$

Then, using KKT condition from Boyd and Vandenberghe (2004), the solution is:

$$\hat{x}(\mathbf{v}_0) = \hat{\alpha}_0 + \mathbf{c}^\top V_\eta^{-1}(\boldsymbol{\xi}) (\mathbf{x}_u - \hat{\alpha}_0 \mathbf{1}),$$

However, not to forget that we actually need the BLUP for the new resolution, $k \times k$, i.e. solve the following problem,

$$\begin{aligned} &\text{minimise} \quad \mathbb{E} \left[\left(x_{k \times k}(\mathbf{s}_0) - \left(\lambda_0 + \sum_{i=1}^m \lambda_i x_{3 \times 3}(\mathbf{u}_i) \right) \right)^2 \right] \\ &\text{subject to} \quad \mathbb{E}[x(\mathbf{v}_0)] = \mathbb{E}[\hat{x}(\mathbf{v}_0)] \end{aligned}$$

Then, applying KKT conditions, we show that the “average predictor” is the optimal predictor.

$$\bar{x}_{k \times k}(\mathbf{s}_0) = \frac{1}{k^2} \sum_{i=1}^{k^2} \hat{x}_{\text{BAU}}(\mathbf{v}_j)$$

Acknowledgements

I would like to sincerely thank my supervisor, Dr. Tingjin Chu, for his invaluable guidance and feedback throughout this project, and for his openness in discussing ideas and answering my questions. I am particularly grateful to the School for offering this special opportunity as an external student.

References

- X. Zheng, N. Cressie, D. A. Clarke, M. A. McGeoch, and A. Zammit-Mangion, “Spatial-statistical downscaling with uncertainty quantification in biodiversity modelling,” *Methods in Ecology and Evolution*, vol. 16, no. 4, pp. 837–853, 2025.
- L. R. LaMotte, “A direct derivation of the reml likelihood function,” *Statistical Papers*, vol. 48, no. 2, pp. 321–327, 2007.
- S. Boyd and L. Vandenberghe, *Convex Optimization*. Cambridge University Press, 2004.