
Name: Ishaan Malhotra

Table of Contents

Roll no. : 1610110152	1
Instructor: Prof. Vijay Chakka	1
Lab 4	1
Aim: Properties of Laplacian Eigenvalue Decomposition	1
Question 1	1
Question 2	2
Question 3	3

Roll no. : 1610110152

Instructor: Prof. Vijay Chakka

Lab 4

Aim: Properties of Laplacian Eigenvalue Decomposition

```
clc
clear all
close all
```

Question 1

Part a)

```
A = randAdjMatrix(5);
LA1 = laplacianMat(A);
[V1, D1] = eig(LA1);
```

% Part b)

```
complete = [[0 1 1 1];[1 0 1 1];[1 1 0 1];[1 1 1 0]];
LA2 = laplacianMat(complete);
[V2, D2] = eig(LA2);
```

% Part c)


```
regular = [[0 1 1 1];[1 0 1 1];[1 1 0 1];[1 1 1 0]];
LA3 = laplacianMat(regular);
[V3, D3] = eig(regular);
[V4, D4] = eig(LA3);
```

```
% Part d) and e)
bipartite = [[0 0 0 1 1 1 1];[0 0 0 1 1 1 1];[0 0 0 1 1 1 1];[1 1 1 0
0 0 0];[1 1 1 0 0 0 0];[1 1 1 0 0 0 0];[1 1 1 0 0 0 0]];
LA4 = normLapMat(bipartite);
[V5, D5] = eig(LA4);
```

Question 2

```
U = load('A1.mat');
lambda1 = [1,0,0,0,0,0,0,0];
lambda2 = [0,1,0,0,0,0,0,0];
lambda3 = [0,0,1,1,0,0,0,0];
lambda4 = [0,0,0,0,1,1,0,0];
lambda5 = [0,0,0,0,1,1,1,1];
lambda6 = [1,1,1,1,1,1,1,1];
lambdas = [lambda1;lambda2;lambda3;lambda4;lambda5;lambda6].';

figure;
for i=1:6
    eigV = diag(lambdas(:,i).');
    L = U.A1*eigV*U.A1.';
    sum = 0;
    for k=1:length(L)
        sum = sum + L(1,k);
    end
    sum = round(1000*sum)/1000;
    D = diag(diag(L));
    A = D - L;
    ran2D = rand(length(A),2);
    subplot(3,2,i);
    if sum == 0
        plot2DGraph(A,ran2D);
    end
end
```



Question 3

```

U2 = load('A2.mat');
U3 = load('A3.mat');
U4 = load('A4.mat');
U5 = load('A5.mat');
U6 = load('A6.mat');

figure;
for i=1:6
    eigV = diag(lambdas(:,i).');
    L = U2.A2*eigV*U2.A2.';
    sum = 0;
    for k=1:length(L)
        sum = sum + L(1,k);
    end
    sum = round(1000*sum)/1000;
    D = diag(diag(L));
    A = D - L;
    ran2D = rand(length(A),2);
    subplot(3,2,i);
    if sum == 0
        plot2DGraph(A,ran2D);
    end
end
end

```

```
figure;
for i=1:6
    eigV = diag(lambdas(:,i).');
    L = U2.A2*eigV*U3.A3.';
    sum = 0;
    for k=1:length(L)
        sum = sum + L(1,k);
    end
    sum = round(1000*sum)/1000;
    D = diag(diag(L));
    A = D - L;
    ran2D = rand(length(A),2);
    subplot(3,2,i);
    if sum == 0
        plot2DGraph(A,ran2D);
    end
end
```

```
figure;
for i=1:6
    eigV = diag(lambdas(:,i).');
    L = U2.A2*eigV*U4.A4.';
    sum = 0;
    for k=1:length(L)
        sum = sum + L(1,k);
    end
    sum = round(1000*sum)/1000;
    D = diag(diag(L));
    A = D - L;
    ran2D = rand(length(A),2);
    subplot(3,2,i);
    if sum == 0
        plot2DGraph(A,ran2D);
    end
end
```

```
figure;
for i=1:6
    eigV = diag(lambdas(:,i).');
    L = U2.A2*eigV*U5.A5.';
    sum = 0;
    for k=1:length(L)
        sum = sum + L(1,k);
    end
    sum = round(1000*sum)/1000;
    D = diag(diag(L));
    A = D - L;
    ran2D = rand(length(A),2);
    subplot(3,2,i);
    if sum == 0
        plot2DGraph(A,ran2D);
    end
end
```

```
figure;
for i=1:6
    eigV = diag(lambdas(:,i).');
    L = U2.A2*eigV*U6.A6.';
    sum = 0;
    for k=1:length(L)
        sum = sum + L(1,k);
    end
    sum = round(1000*sum)/1000;
    D = diag(diag(L));
    A = D - L;
    ran2D = rand(length(A),2);
    subplot(3,2,i);
    if sum == 0
        plot2DGraph(A,ran2D);
    end
end
```

```
%{
                                Observation in Q2 and Q3
```

We observe that for eigenvalue set Lambda2 to Lambda5, four graphs are formed while for other eigenvalue sets, no graph is formed. On inspecting the Graph Laplacian of the graph formed, we observe that to be a valid Laplacian Matrix, the sum of elements in a columns and rows should be equal to 0.

```
%}
```

Published with MATLAB® R2018a