

4 D'S FOR DEALING WITH CRAVINGS



DELAY

Set a time limit before you give in to smoking. Delay as long as you can.



DEEP BREATHING AND MEDITATION

Take 10 deep breaths to relax yourself.



DRINK WATER

Drinking water is a healthy alternative to sticking a cigarette in your mouth.



DO SOMETHING ELSE TO DISTRACT YOURSELF

Read, go for a walk, listen to music, watch TV or engage in any hobby.



ANTICIPATE CHALLENGES

Learn how to Deal with them

HEADACHES

Get enough sleep, eat regularly, undertake physical activity.

COUGHING

Stay hydrated, consume a spoonful of honey, warm tea, juices and avoid dairy.

INCREASED APPETITE

AND WEIGHT GAIN

Make healthy eating choices (fruits, vegetables etc.) and drink lots of water.

MOOD CHANGES (SADNESS, IRRITABILITY, FRUSTRATION, OR ANGER)

Try focusing on positive thoughts about quitting or filling your head with happy, uplifting music to drown out the negative thoughts and emotions.

RESTLESSNESS

Get up and move around.

DECREASED HEART RATE

Make healthy lifestyle choices.

DIFFICULTY

CONCENTRATING

Do yoga, meditation, and mental imagery.


FLU-LIKE SYMPTOMS

Rest and hydrate.

INSOMNIA

Count sheep, guided meditation, and self-guided imagery.

BENEFITS OF QUITTING



BENEFITS AT DIFFERENT AGES



AT ABOUT 30

Gain almost 10 years of life expectancy



AT ABOUT 40

Gain 9 years of life expectancy



AT ABOUT 50

Gain 6 years of life expectancy



AT ABOUT 60

Gain 3 years of life expectancy

“QUITTING SMOKING eliminates the risk of secondhand smoke exposure for children, protecting them from a multitude of health complications. It can also significantly improve fertility and reduce the likelihood of adverse pregnancy outcomes, including premature birth, low birth weight, and miscarriage.

OUR COLLABORATORS




Hôpitaux
Universitaires
Genève

Ref. PEN Manual (MoHP), 2075



**KHDC
NEPAL**



TOBACCO CESSATION GUIDE FOR PRIMARY CARE PROVIDERS

USE THIS GUIDE TO

- IDENTIFY AND REACH TOBACCO USERS.
- TRIGGER USERS TO MAKE QUIT ATTEMPT.
- HELP THOSE RECEIVING BRIEF ADVICE TO QUIT SUCCESSFULLY.

Primary care providers have the potential to engage with over 80% of all tobacco users, catalysing quit attempts in 40% of cases, and achieving a successful quit rate of 2-3%

5 A'S OF SMOKING CESSATION

01 ASK

SYSTEMATICALLY IDENTIFY ALL TOBACCO USERS AT EVERY VISIT

Ask in a friendly, non accusatory way.
Tobacco use status should be included in all medical notes.
Ask about second-hand tobacco smoke exposure as well.

02 ADVISE

PERSUADE ALL TOBACCO USERS THAT THEY NEED TO QUIT

Urge tobacco user to quit in clear, strong and personalized manner.

03 ASSESS

DETERMINE READINESS TO MAKE A QUIT ATTEMPT

Ask two questions in relation to 'importance' and 'self-efficacy'.

Would you like to be a non-tobacco user?	Yes	Unsure	No
Do you think you have chance of quitting successfully?	Yes	Unsure	No

* Any answer in the shaded area indicates that the tobacco user is NOT ready to quit; to promote motivation to quit **refer to 5Rs**

04 ASSIST

HELP THE PATIENT WITH A QUIT PLAN



S SET A QUIT DATE
within 2 weeks



T TELL FAMILY, FRIENDS, COWORKERS
about quitting and request support



A ANTICIPATE CHALLENGES
to upcoming quit attempt



R REMOVE TOBACCO PRODUCTS
from environment; make home smokefree

05 ARRANGE

SCHEDULE FOLLOW-UP CONTACTS OR REFERRAL TO SPECIALIST SUPPORT.


WHEN 1st < 1 week after quit date. | 2nd within 1 month.

HOW Phone call, personal visit, and/or email to follow up.

WHAT Identify problems, Remind of social support, Assess need for expert consultation, Schedule follow-up

APPROACH TO COUNSELLING

THIS SHOULD TAKE AROUND 3-5 MINUTES DURING CONSULTATION !!



MAD-TEA TO ASSIST AVOIDING EXPOSURE TO SECOND HAND SMOKE:

M Meet friends/relatives in smoke-free spaces.
A Ask family members and visitors to smoke outside.
D Declare home and personal spaces to be smoke free.
T Talk to family members and colleagues about the risks of second-hand smoke.
E Encourage family members, friends, and workmates to stop smoking.
A Advocate comprehensive smoke-free laws or regulations in workplaces and public places.

FOR THOSE WHO ABSTAINED :

Congratulate on success

FOR RELAPSE :

Remind it as a learning experience;
Review circumstances and elicit recommitment;
Refer for expert support.

5 R'S OF SMOKING CESSATION

1. RELEVANCE



Encourage the patient to indicate how quitting is personally relevant to him or her.

2. RISK



Encourage patient to identify potential negative consequences of tobacco use that are relevant to him or her.



TOBACCO KILLS UP TO HALF OF ITS USERS



TOBACCO USE PUTS YOUR FAMILY AT RISK.



5-15% OF INCOME IS SPENT ON TOBACCO.



SMOKER'S CHILDREN ARE MORE LIKELY TO SMOKE

3. REWARDS



Ask the patient to identify potential relevant benefits of stopping tobacco use.

4. ROADBLOCKS



Ask the patient to identify barriers or impediments to quitting and provide counseling that could address barriers.



FOCUS ON 'ROADBLOCK(S)' IF THE PATIENT THINKS HE/SHE CAN'T QUIT.



CRAVINGS/URGES USUALLY LASTS FOR 1-2 MINUTES



SUDDEN STOPPING MAY CAUSE SYMPTOMS THAT MAY LAST 2-4 WEEKS.

5. REPETITION



Repeat assessment of readiness to quit.
Motivational intervention should be repeated every time a patient who is not willing to quit visits the clinic setting.

CONTACT US

For any clinical queries / If anything in the document is difficult to understand



+977-9852072088 / 9802766889

www.khdcnepal.org

