

2D Image Transforms

16-385 Computer Vision (Kris Kitani)

Carnegie Mellon University




Extract features from an image ...



what do we do next?

Feature matching

(object recognition, 3D reconstruction, augmented reality, image stitching)



How do you compute the transformation?

Given a set of matched feature points

$$\{x_i, x'_i\} \leftarrow \text{set of point correspondences}$$

point in point in the
one image other image

and a transformation

$$x' = f(x; p)$$

transformation parameters
function

Find the best estimate of

p

What kind of transformation functions are there?

$$\boldsymbol{x}' = \boldsymbol{f}(\boldsymbol{x}; \boldsymbol{p})$$

2D Transformations



translation



rotation



aspect



affine




perspective



cylindrical

2D Planar Transformations



2D Planar Transformations



Scale

- Each component multiplied by a scalar
- Uniform scaling - same scalar for each component

2D Planar Transformations



Scale

Scale

$$x' = ax$$

$$y' = by$$

- Each component multiplied by a scalar
- Uniform scaling - same scalar for each component

2D Planar Transformations




Scale

Scale

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \underbrace{\begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}}_{\text{scaling matrix } S} \begin{bmatrix} x \\ y \end{bmatrix}$$

- Each component multiplied by a scalar
- Uniform scaling - same scalar for each component

2D Planar Transformations



2D Planar Transformations



Shear

$$x' = x + a \cdot y$$

$$y' = b \cdot x + y$$


2D Planar Transformations




Shear

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 1 & a \\ b & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$


2D Planar Transformations




2D Planar Transformations



2D Planar Transformations





Polar coordinates...

$$x = r \cos(\phi)$$

$$y = r \sin(\phi)$$

$$x' = r \cos(\phi + \theta)$$

$$y' = r \sin(\phi + \theta)$$

Trig Identity...

$$x' = r \cos(\phi) \cos(\theta) - r \sin(\phi) \sin(\theta)$$


$$y' = r \sin(\phi) \cos(\theta) + r \cos(\phi) \sin(\theta)$$

Substitute...

$$x' = x \cos(\theta) - y \sin(\theta)$$

$$y' = x \sin(\theta) + y \cos(\theta)$$


2D Planar Transformations



2D **linear** transformation

(can be written in matrix form)


$$x' = f(x; p)$$



$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \mathbf{M} \begin{bmatrix} x \\ y \end{bmatrix}$$

parameters p

point x



Scale

$$\mathbf{M} = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix}$$

Flip across y

$$\mathbf{M} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

Rotate

$$\mathbf{M} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

Flip across origin

$$\mathbf{M} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$


Shear

$$\mathbf{M} = \begin{bmatrix} 1 & s_x \\ s_y & 1 \end{bmatrix}$$


Identity

$$\mathbf{M} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

How do you represent translation with a 2 x 2 matrix?



How do you represent translation with a 2 x 2 matrix?




Q: How can we represent translation in matrix form?

$$x' = x + t_x$$

$$y' = y + t_y$$

Homogeneous Coordinates



Represent 2D point with a 3D vector

Q: How can we represent translation in matrix form?

$$x' = x + t_x$$




$$y' = y + t_y$$

A: append 3rd element and append 3rd column & row

$$\begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \quad \mathbf{M} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$

Homogeneous Coordinates

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix}$$




A 2D point in an image can be represented as a 3D vector

$$\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix} \iff \mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

where $x = \frac{x_1}{x_3}$ $y = \frac{x_2}{x_3}$

Why?

Think of a point on the image plane in 3D



You can think of a conversion to homogenous coordinates as a conversion of a **point** to a **ray**

Conversion:

- 2D point → homogeneous point

append 1 as 3rd coordinate

$$\begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

- homogeneous point → 2D point

divide by 3rd coordinate

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow \begin{bmatrix} x/w \\ y/w \end{bmatrix}$$

Special Properties

- Scale invariant

$$\begin{bmatrix} x & y & w \end{bmatrix}^T = \lambda \begin{bmatrix} x & y & w \end{bmatrix}^T$$

- Point at infinity

$$\begin{bmatrix} x & y & 0 \end{bmatrix}$$

coz $x/0$ and $y/0$ is undefined

- Undefined

$$\begin{bmatrix} 0 & 0 & 0 \end{bmatrix}$$

$0/0$ is undefined

Basic 2D transformations as 3x3 matrices

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Translate

$$\begin{bmatrix} \mathbf{x}' \\ \mathbf{y}' \\ 1 \end{bmatrix} = \begin{bmatrix} \mathbf{s}_x & 0 & 0 \\ 0 & \mathbf{s}_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \\ 1 \end{bmatrix}$$

Scale

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \Theta & -\sin \Theta & 0 \\ \sin \Theta & \cos \Theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Rotate

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & \beta_x & 0 \\ \beta_y & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Shear

Matrix Composition

Transformations can be combined by matrix multiplication

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \left(\begin{bmatrix} 1 & 0 & tx \\ 0 & 1 & ty \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \Theta & -\sin \Theta & 0 \\ \sin \Theta & \cos \Theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} sx & 0 & 0 \\ 0 & sy & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

$\mathbf{p}' = \quad T(t_x, t_y) \quad R(\Theta) \quad S(s_x, s_y) \quad \mathbf{p}$

Does the order of multiplication matter?

2D transformations




Figure 1: Basic set of 2D planar transformations

Name	Matrix	# D.O.F.
translation	$\left[\begin{array}{c c} \mathbf{I} & \mathbf{t} \end{array} \right]_{2 \times 3}$	2
rigid (Euclidean)	$\left[\begin{array}{c c} \mathbf{R} & \mathbf{t} \end{array} \right]_{2 \times 3}$	3
similarity	$\left[\begin{array}{c c} s\mathbf{R} & \mathbf{t} \end{array} \right]_{2 \times 3}$	4
affine	$\left[\begin{array}{c} \mathbf{A} \end{array} \right]_{2 \times 3}$	6
projective	$\left[\begin{array}{c} \tilde{\mathbf{H}} \end{array} \right]_{3 \times 3}$	8

Affine Transformation


Affine transformations are combinations of

- Linear transformations, and
- Translations

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

Properties of affine transformations:

- Origin does not necessarily map to origin
- Lines map to lines
- Parallel lines remain parallel
- Ratios are preserved
- Closed under composition (affine times affine is affine)



Will the last coordinate w ever change?

Coming soon...

Projective Transform

Projective transformations are combos of

- Affine transformations, and
- Projective warps

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

Properties of projective transformations:

- Origin does not necessarily map to origin
- Lines map to lines
- Parallel lines do not necessarily remain parallel
- Ratios are not preserved
- Closed under composition
- Models change of basis
- Projective matrix is defined up to a scale (8 DOF)

