

# A MONTE CARLO SIMULATION OF THE STARS OF THE GALAXY

IMF: CHABRIER SINGLE 2005

**Names:** David Anríquez  
Isidora Morales

Teaching Lucas Gajardo  
assistants: Barbara Gutierrez

**Teacher:** Alejandra Rojas

# TABLE OF CONTENT

01 INTRODUCTION

01 IMF:  
CHABRIER S 2005

02 FLOWCHART

03 RESULTS

04 CONCLUTIONS

05 REFERENCES

# INTRODUCTION

**Monte Carlo statistical model**

**Simulated stars using IMF of  
Chabrier 2005**

**Diverse sample of stars:  
100, 1.000, 10.000, 100.000 and  
1 million**


**Study of stars on the MS and  
the object they will become  
upon leaving it**


# Initial Mass Function (IMF)

## Chabrier S 2005

- Focused on individual stars
- It arises with the aim of properly characterizing brown dwarfs.
- Its mass distribution is more constant than that of the other IMFs.
- The log-normal form for low masses is well supported observationally

$$\xi(\log m) = 0.093 \times \exp \left[ \frac{-(\log m - \log 0.2^2)}{2 \times (0.55)^2} \right], \quad m \leq 1M_{\odot}$$
$$= 0.041 m^{-1.35}, \quad m \geq 1M_{\odot}$$







# FLOWCHART

# OUR RESULTS

# RESIDUAL MASSES




**100 Stars**




**1.000 Stars**

# RESIDUAL MASSES




10.000 Stars



100.000 Stars





# RESIDUAL MASSES





1.000.000 Stars

Kalirai et al. (2008)





Raithel et al. (2018)


# FINAL MASSES DISTRIBUTION





# FINAL MASSES DISTRIBUTION





# FINAL MASSES DISTRIBUTION




# AGE DISTRIBUTION



# AGE DISTRIBUTION




# AGE DISTRIBUTION



# COMPARISON WITH THE MW

---





# COMPARISON WITH THE MW

| Number of stars | BH fraction | NS fraction | WD fraction | MS fraction |
|-----------------|-------------|-------------|-------------|-------------|
| 100             | 15.00 %     | 8.00%       | 31.00%      | 46.00%      |
| 1.000           | 12.30%      | 11.40%      | 28.10%      | 48.20%      |
| 10.000          | 11.16%      | 10.09       | 29.99%      | 48.76%      |
| 100.000         | 11.24%      | 10.06%      | 29.68%      | 49.02%      |
| 1.000.000       | 11.22 %     | 9.90%       | 29.54%      | 49.33%      |

# CONCLUSIONS

Using the Monte Carlo statistical model, it was possible to generate various analyses for the stellar distribution based on the initial mass function of Chabrier (2005).

It was possible to observe the impact this IMF has on the stellar mass distribution, which is the objective for which it was created.

The results obtained indicate that it is indeed a good regime for low-mass stars, such as brown dwarfs, but not for more massive stars, ultimately predicting a much higher value than expected for BHs in the Milky Way. This value is superior to 10% in all cases.

# REFERENCES

- How Many Stars in the Milky Way? | NASA Blueshift. (2015, 22 julio). <https://asd.gsfc.nasa.gov/blueshift/index.php/2015/07/22/how-many-stars-in-the-milky-way/>
- Samland, M. (1998). Modeling the Evolution of Disk Galaxies. II. Yields of Massive Stars. *The Astrophysical Journal*, 496(1), 155-171. <https://doi.org/10.1086/305368>
- Sartore, N., Ripamonti, E., Treves, A., & Turolla, R. (2009). Galactic neutron stars. *Astronomy And Astrophysics*, 510, A23. <https://doi.org/10.1051/0004-6361/200912222>
- Tremblay, P., Bédard, A., O'Brien, M. W., Munday, J., Elms, A. K., Pietro Gentilo Fusillo, N., & Sahu, S. (2024). The Gaia white dwarf revolution. *New Astronomy Reviews*, 99, 101705. <https://doi.org/10.1016/j.newar.2024.101705>
- Kalirai, J. S., Hansen, B. M. S., Kelson, D. D., Reitzel, D. B., Rich, R. M., Richer, H. B., Kalirai, J. S., Hansen, B. M. S., Kelson, D. D., Reitzel, D. B., Rich, R. M., & Richer, H. B. (2008). The Initial-Final Mass Relation: Direct Constraints at the Low-Mass End. *The Astrophysical Journal*, 676(1), 594-609. <https://doi.org/10.1086/527028>
- Raithel, C. A., Sukhbold, T., & Özel, F. (2018). Confronting Models of Massive Star Evolution and Explosions with Remnant Mass Measurements. *The Astrophysical Journal*, 856(1), 35. <https://doi.org/10.3847/1538-4357/aab09b>
- Chabrier, G. (2007). The Initial Mass Function: From Salpeter 1955 to 2005. En *Astrophysics and space science library* (pp. 41-50). [https://doi.org/10.1007/978-1-4020-3407-7\\_5](https://doi.org/10.1007/978-1-4020-3407-7_5)

Project 4

THANK YOU!

Questions?