

A MONTE CARLO SIMULATION OF THE STARS OF THE GALAXY

IMF: CHABRIER SINGLE 2005

Names: David Anríquez
Isidora Morales

Teaching assistants: Lucas Gajardo
Barbara Gutierrez

Teacher: Alejandra Rojas

TABLE OF CONTENT

01 INTRODUCTION

01 IMF: CHABRIER S 2005

02 FLOWCHART

03 RESULTS

04 CONCLUTIONS

05 REFERENCES

INTRODUCTION

Monte Carlo statistical model

Simulated stars using IMF of Chabrier 2005

Diverse sample of stars: 100, 1.000, 10.000, 100.000 and 1 million

Study of stars on the MS and the object they will become upon leaving it

Initial Mass Function (IMF)

Chabrier S 2005

- Focused on individual stars
- It arises with the aim of properly characterizing brown dwarfs.
- Its mass distribution is more constant than that of the other IMFs.
- The log-normal form for low masses is well supported observationally

$$\xi(log\, m) = 0.093 \times \exp\left[\frac{-(\log m - \log 0.2^2)}{2 \times (0.55)^2}\right], \quad m \leq 1 M_\odot$$

$$= 0.041\, m^{-1.35}, \quad m \geq 1 M_\odot$$





FLOWCHART

Project 4 AST-305

OUR RESULTS

Project 4 AST-305

RESIDUAL MASSES



100 Stars



1.000 Stars

RESIDUAL MASSES



10.000 Stars



100.000 Stars

RESIDUAL MASSES



1.000.000 Stars



Kalirai et al. (2008)



Raithel et al. (2018)

FINAL MASSES DISTRIBUTION





FINAL MASSES DISTRIBUTION





FINAL MASSES DISTRIBUTION



AGE DISTRIBUTION





AGE DISTRIBUTION





AGE DISTRIBUTION



COMPARISON WITH THE MW



COMPARISON WITH THE MW

Number of stars	BH fraction	NS fraction	WD fraction	MS fraction
100	15.00 %	8.00%	31.00%	46.00%
1.000	12.30%	11.40%	28.10%	48.20%
10.000	11.16%	10.09	29.99%	48.76%
100.000	11.24%	10.06%	29.68%	49.02%
1.000.000	11.22 %	9.90%	29.54%	49.33%

CONCLUTIONS

Using the Monte Carlo statistical model, it was possible to generate various analyses for the stellar distribution based on the initial mass function of Chabrier (2005).

It was possible to observe the impact this IMF has on the stellar mass distribution, which is the objective for which it was created.

The results obtained indicate that it is indeed a good regime for low-mass stars, such as brown dwarfs, but not for more massive stars, ultimately predicting a much higher value than expected for BHs in the Milky Way. This value is superior to 10% in all cases.

Proyect 4 AST-305

REFERENCES

- How Many Stars in the Milky Way? | NASA Blueshift. (2015, 22 julio). https://asd.gsfc.nasa.gov/blueshift/index.php/2015/07/22/how-many-stars-in-the-milky-way/
- Samland, M. (1998). Modeling the Evolution of Disk Galaxies. II. Yields of Massive Stars. The Astrophysical Journal, 496(1), 155-171. https://doi.org/10.1086/305368
- Sartore, N., Ripamonti, E., Treves, A., & Turolla, R. (2009). Galactic neutron stars. Astronomy And Astrophysics, 510, A23. https://doi.org/10.1051/0004-6361/200912222
- Tremblay, P., Bédard, A., O'Brien, M. W., Munday, J., Elms, A. K., Pietro Gentillo Fusillo, N., & Sahu, S. (2024). The Gaia white dwarf revolution. New Astronomy Reviews, 99, 101705. https://doi.org/10.1016/j.newar.2024.101705
- Kalirai, J. S., Hansen, B. M. S., Kelson, D. D., Reitzel, D. B., Rich, R. M., Richer, H. B., Kalirai, J. S., Hansen, B. M. S., Kelson, D. D., Reitzel, D. B., Rich, R. M., & Richer, H. B. (2008). The Initial-Final Mass Relation: Direct Constraints at the Low-Mass End. The Astrophysical Journal, 676(1), 594-609. https://doi.org/10.1086/527028
- Raithel, C. A., Sukhbold, T., & Özel, F. (2018). Confronting Models of Massive Star Evolution and Explosions with Remnant Mass Measurements. The Astrophysical Journal, 856(1), 35. https://doi.org/10.3847/1538-4357/aab09b
- Chabrier, G. (2007). The Initial Mass Function: From Salpeter 1955 to 2005. En Astrophysics and space science library (pp. 41-50). https://doi.org/10.1007/978-1-4020-3407-7_5

SCAN ME

Project 4

THANK YOU!

Questions?