

# Direct extraction of path weak values from interferograms without auxiliary qubits

# Introduction to weak values

## References:

Complex weak values in quantum measurement

Richard Jozsa

The significance of the imaginary part of the weak value

J. Dressel and A. N. Jordan

# Introduction to weak values

Weak value:

$$A_w = \frac{\langle \psi_f | \hat{A} | \psi_{in} \rangle}{\langle \psi_f | \psi_{in} \rangle}$$

# Introduction to weak values

Weak value:

$$A_w = \frac{\langle \psi_f | \hat{A} | \psi_{in} \rangle}{\langle \psi_f | \psi_{in} \rangle}$$

- Observable:  $\hat{A}$
- Pre-selected state:  $|\psi_{in}\rangle$
- Post-selected State:  $|\psi_f\rangle$

# Introduction to weak values

Weak value:

$$A_w = \frac{\langle \psi_f | \hat{A} | \psi_{in} \rangle}{\langle \psi_f | \psi_{in} \rangle}$$

- Observable:  $\hat{A}$
- Pre-selected state:  $|\psi_{in}\rangle$
- Post-selected State:  $|\psi_f\rangle$
- Complex number
- Not bounded by eigenvalues

# Introduction to weak values

## Von Neumann measurement

$$|\phi\rangle \otimes |\psi_{\text{in}}\rangle$$

# Introduction to weak values

## Von Neumann measurement

$$|\phi\rangle \otimes |\psi_{\text{in}}\rangle$$

|            |  
Detector   System

# Introduction to weak values

## Von Neumann measurement

$$|\phi\rangle \otimes |\psi_{\text{in}}\rangle = |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

|  
Detector    System

# Introduction to weak values

## Von Neumann measurement

$$|\phi\rangle \otimes |\psi_{\text{in}}\rangle = |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

$$\hat{H}_I(t) = g(t)\hat{p} \otimes \hat{A}$$

|

Interaction  
Hamiltonian

# Introduction to weak values

## Von Neumann measurement

$$|\phi\rangle \otimes |\psi_{\text{in}}\rangle = |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

$$\hat{H}_I(t) = g(t)\hat{p} \otimes \hat{A} = g(t)\hat{p}_d \hat{A}_s$$

|

Interaction  
Hamiltonian

# Introduction to weak values

## Von Neumann measurement

$$|\phi\rangle \otimes |\psi_{\text{in}}\rangle = |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

$$\hat{H}_I(t) = g(t)\hat{p} \otimes \hat{A} = g(t)\hat{p}_d \hat{A}_s$$

$$e^{-i \int_{t_1}^{t_2} \hat{H}_I(t) dt} |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

# Introduction to weak values

## Von Neumann measurement

$$|\phi\rangle \otimes |\psi_{\text{in}}\rangle = |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

$$\hat{H}_I(t) = g(t)\hat{p} \otimes \hat{A} = g(t)\hat{p}_d \hat{A}_s$$

$$e^{-i \int_{t_1}^{t_2} \hat{H}_I(t) dt} |\phi\rangle_d |\psi_{\text{in}}\rangle_s = e^{-i G \hat{p}_d \hat{A}_s} |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

# Introduction to weak values

## Von Neumann measurement

$$|\phi\rangle \otimes |\psi_{\text{in}}\rangle = |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

$$\hat{H}_I(t) = g(t)\hat{p} \otimes \hat{A} = g(t)\hat{p}_d \hat{A}_s$$

$$e^{-i \int_{t_1}^{t_2} \hat{H}_I(t) dt} |\phi\rangle_d |\psi_{\text{in}}\rangle_s = e^{-i G \hat{p}_d \hat{A}_s} |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

$$\int_{t_1}^{t_2} g(t) dt = G$$

# Introduction to weak values

## Von Neumann measurement

$$\sum_a \langle a | \psi_{\text{in}} \rangle |a\rangle = \sum_a c_a |a\rangle$$

# Introduction to weak values

## Von Neumann measurement

$$\sum_a \langle a | \psi_{\text{in}} \rangle |a\rangle = \sum_a c_a |a\rangle : \hat{A}_s |a\rangle = a |a\rangle$$

# Introduction to weak values

## Von Neumann measurement

$$\sum_a \langle a | \psi_{\text{in}} \rangle |a\rangle = \sum_a c_a |a\rangle : \hat{A}_s |a\rangle = a |a\rangle$$

$$e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

# Introduction to weak values

## Von Neumann measurement

$$\sum_a \langle a | \psi_{\text{in}} \rangle |a\rangle = \sum_a c_a |a\rangle : \hat{A}_s |a\rangle = a |a\rangle$$

$$e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{\text{in}}\rangle_s = \sum_a e^{-iG a \hat{p}_d} |\phi\rangle_d c_a |a\rangle$$

# Introduction to weak values

## Von Neumann measurement

$$\sum_a \langle a | \psi_{\text{in}} \rangle |a\rangle = \sum_a c_a |a\rangle : \hat{A}_s |a\rangle = a |a\rangle$$

$$e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{\text{in}}\rangle_s = \sum_a [e^{-iG_a\hat{p}_d}] |\phi\rangle_d c_a |a\rangle$$

Translation operator

# Introduction to weak values

## Von Neumann measurement

$$\sum_a \langle x |_d e^{-iG_a \hat{p}_d} |\phi \rangle_d c_a |a \rangle$$

a  
|

Position  
projection

# Introduction to weak values

## Von Neumann measurement


$$\sum_a \langle x|_d e^{-iG a \hat{p}_d} |\phi\rangle_d c_a |a\rangle = \sum_a \phi(x + G a) c_a |a\rangle$$

Position  
projection

# Introduction to weak values

## Stern-Gerlach:


Position    Spin



# Introduction to weak values

## Stern-Gerlach:

Position      Spin




$$\hat{H}_I(t) = g(t)\hat{p}_z \otimes \hat{\sigma}_z$$

# Introduction to weak values

## Stern-Gerlach:

Position      Spin



$$\hat{H}_I(t) = g(t)\hat{p}_z \otimes \hat{\sigma}_z$$

# Introduction to weak values

**Weak measurement:**

$$e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{\text{in}}\rangle_s$$

# Introduction to weak values

**Weak measurement:**

$$\langle \psi_f |_s e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{in}\rangle_s$$

# Introduction to weak values

**Weak measurement:**

$$\langle \psi_f |_s e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{in}\rangle_s \underset{G \ll 1}{\approx}$$

# Introduction to weak values

## Weak measurement:

$$\langle \psi_f |_s e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{in}\rangle_s \underset{G \ll 1}{\approx} \langle \psi_f |_s \left( \hat{\mathbb{I}} - iG\hat{p}_d\hat{A}_s \right) |\phi\rangle_d |\psi_{in}\rangle_s$$

# Introduction to weak values

## Weak measurement:

$$\begin{aligned} \langle \psi_f |_s e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{in}\rangle_s &\approx \langle \psi_f |_s \left( \hat{\mathbb{I}} - iG\hat{p}_d\hat{A}_s \right) |\phi\rangle_d |\psi_{in}\rangle_s \\ &= \langle \psi_f | \psi_{in} \rangle \left( \hat{\mathbb{I}} - iG \frac{\langle \psi_f | \hat{A} | \psi_{in} \rangle}{\langle \psi_f | \psi_{in} \rangle} \hat{p}_d \right) |\phi\rangle_d \end{aligned}$$

# Introduction to weak values

## Weak measurement:

$$\begin{aligned} \langle \psi_f |_s e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{in}\rangle_s &\approx \langle \psi_f |_s \left( \hat{\mathbb{I}} - iG\hat{p}_d\hat{A}_s \right) |\phi\rangle_d |\psi_{in}\rangle_s \\ &= \langle \psi_f | \psi_{in} \rangle \left( \hat{\mathbb{I}} - iG \frac{\langle \psi_f | \hat{A} | \psi_{in} \rangle}{\langle \psi_f | \psi_{in} \rangle} \hat{p}_d \right) |\phi\rangle_d \end{aligned}$$

# Introduction to weak values

## Weak measurement:

$$\begin{aligned} \langle \psi_f |_s e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{in}\rangle_s &\approx \langle \psi_f |_s \left( \hat{\mathbb{I}} - iG\hat{p}_d\hat{A}_s \right) |\phi\rangle_d |\psi_{in}\rangle_s \\ &= \langle \psi_f | \psi_{in} \rangle \left( \hat{\mathbb{I}} - iG A_w \hat{p}_d \right) |\phi\rangle_d \end{aligned}$$

# Introduction to weak values

## Weak measurement:

$$\begin{aligned} \langle \psi_f |_s e^{-iG\hat{p}_d\hat{A}_s} |\phi\rangle_d |\psi_{in}\rangle_s &\approx \langle \psi_f |_s \left( \hat{\mathbb{I}} - iG\hat{p}_d\hat{A}_s \right) |\phi\rangle_d |\psi_{in}\rangle_s \\ &= \langle \psi_f | \psi_{in} \rangle \left( \hat{\mathbb{I}} - iG A_w \hat{p}_d \right) |\phi\rangle_d \\ &\approx \langle \psi_f | \psi_{in} \rangle \left( e^{-iG A_w \hat{p}_d} \right) |\phi\rangle_d \end{aligned}$$

# Introduction to weak values

## Weak measurement:

$$\langle \psi_f | \psi_{in} \rangle \langle x |_d \left( e^{-iGA_w \hat{p}_d} \right) |\phi\rangle_d \approx \langle \psi_f | \psi_{in} \rangle \phi(x - GA_w)$$

# Introduction to weak values

## Weak measurement:

$$\langle \psi_f | \psi_{in} \rangle \langle x |_d (e^{-iGA_w \hat{p}_d}) |\phi \rangle_d \approx \langle \psi_f | \psi_{in} \rangle \phi(x - GA_w)$$

**The sense in which a “weak measurement” of a spin- $\frac{1}{2}$  particle’s spin component yields a value 100**

I. M. Duck and P. M. Stevenson

# Introduction to weak values

## Weak measurement:

$$\langle \psi_f | \psi_{in} \rangle \langle x |_d \left( e^{-iG A_w \hat{p}_d} \right) |\phi\rangle_d \approx \langle \psi_f | \psi_{in} \rangle \phi(x - G A_w)$$

$$A_w = A_w^{\Re} + i A_w^{\Im}$$

# Introduction to weak values

## Weak measurement:

$$\langle \psi_f | \psi_{in} \rangle \langle x |_d (e^{-iG A_w \hat{p}_d}) |\phi \rangle_d \approx \langle \psi_f | \psi_{in} \rangle \phi(x - G A_w)$$

$$A_w = A_w^{\Re} + i A_w^{\Im}$$

For a real valued detector function  $\phi(x)$

# Introduction to weak values

## Weak measurement:

$$\langle \psi_f | \psi_{\text{in}} \rangle \langle x |_d \left( e^{-iG A_w \hat{p}_d} \right) |\phi\rangle_d \approx \langle \psi_f | \psi_{\text{in}} \rangle \phi(x - G A_w)$$

$$A_w = A_w^{\Re} + i A_w^{\Im}$$

For a real valued detector function  $\phi(x)$

$$\langle \hat{x} \rangle_f = \langle \hat{x} \rangle_i + G A_w^{\Re}$$

$$\langle \hat{p} \rangle_f = \langle \hat{p} \rangle_i + G A_w^{\Im} \text{Var}(\hat{p})$$

# IS THE WEAK VALUE ALWAYS WEAK?

# Is the weak value always weak?

**Expectation value:**

$$\langle \psi_{\text{in}} | \hat{A} | \psi_{\text{in}} \rangle$$

# Is the weak value always weak?

Expectation value:

$$\langle \psi_{\text{in}} | \hat{A} | \psi_{\text{in}} \rangle$$
$$\hat{\mathbb{I}} = \sum_k | \psi_f^k \rangle \langle \psi_f^k |$$

# Is the weak value always weak?

**Expectation value:**

$$\langle \psi_{\text{in}} | \hat{A} | \psi_{\text{in}} \rangle = \sum_k \langle \psi_{\text{in}} | \psi_f^k \rangle \langle \psi_f^k | \hat{A} | \psi_{\text{in}} \rangle$$
$$\hat{\mathbb{I}} = \sum_k | \psi_f^k \rangle \langle \psi_f^k |$$

# Is the weak value always weak?

**Expectation value:**

$$\langle \psi_{\text{in}} | \hat{A} | \psi_{\text{in}} \rangle = \sum_k \langle \psi_{\text{in}} | \psi_f^k \rangle \langle \psi_f^k | \hat{A} | \psi_{\text{in}} \rangle$$

# Is the weak value always weak?

**Expectation value:**

$$\begin{aligned}\langle \psi_{\text{in}} | \hat{A} | \psi_{\text{in}} \rangle &= \sum_k \langle \psi_{\text{in}} | \psi_{\text{f}}^k \rangle \langle \psi_{\text{f}}^k | \hat{A} | \psi_{\text{in}} \rangle \\ &= \sum_k | \langle \psi_{\text{f}} | \psi_{\text{in}}^k \rangle |^2 \frac{\langle \psi_{\text{f}}^k | \hat{A} | \psi_{\text{in}} \rangle}{\langle \psi_{\text{f}}^k | \psi_{\text{in}} \rangle}\end{aligned}$$

# Is the weak value always weak?


**Expectation value:**

$$\begin{aligned}\langle \psi_{\text{in}} | \hat{A} | \psi_{\text{in}} \rangle &= \sum_k \langle \psi_{\text{in}} | \psi_f^k \rangle \langle \psi_f^k | \hat{A} | \psi_{\text{in}} \rangle \\ &= \sum_k | \langle \psi_f | \psi_{\text{in}}^k \rangle |^2 \frac{\langle \psi_f^k | \hat{A} | \psi_{\text{in}} \rangle}{\langle \psi_f^k | \psi_{\text{in}} \rangle} \\ &= \sum_k | \langle \psi_f | \psi_{\text{in}}^k \rangle |^2 A_w^k\end{aligned}$$

# CAN WEAK VALUES DESCRIBE INTERFEROMETRY?

# Weak values and interferometry

## Standard interferometry formalism



# Weak values and interferometry

## Standard interferometry formalism

Measured intensity

$$I_{\pm,1}(\chi, \chi_0) = \left| \langle \pm | (e^{-i\chi_0} a_1 |1\rangle + e^{i\chi} a_2 |2\rangle) \right|^2 = \frac{1}{2} \pm a_1 a_2 \cos(\chi + \chi_0)$$