

Lecture 5: Noncoercive problems: Babuška theorem

Pr. Ismail Merabet

Univ. of K-M-Ouargla

November 9, 2025

Contents

- 1 Introduction
- 2 Examples of noncoercive problems
- 3 Babuška theorem
- 4 Discretization of non-coercive problems

Introduction

So far we have treated **coercive problems**. This means that in the linear variational problem, we are trying to solve,

$$\begin{cases} \text{find } u \in V \text{ such that} \\ a(u, v) = F(v), \text{ for all } v \in V \end{cases} \quad (1)$$

the bilinear form $a(u, v)$ **satisfies**

$$a(v, v) \geq \alpha \|v\|_V^2, \quad \forall v \in V$$

for some $\alpha > 0$.

Recall that the best constant α satisfying the definition is given by

$$\alpha := \inf_{\substack{v \in V \\ v \neq 0}} \frac{a(v, v)}{\|v\|_V^2}$$

Introduction

So far we have treated **coercive problems**. This means that in the linear variational problem, we are trying to solve,

$$\begin{cases} \text{find } u \in V \text{ such that} \\ a(u, v) = F(v), \text{ for all } v \in V \end{cases} \quad (1)$$

the bilinear form $a(u, v)$ **satisfies**

$$a(v, v) \geq \alpha \|v\|_V^2, \quad \forall v \in V$$

for some $\alpha > 0$.

Recall that the best constant α satisfying the definition is given by

$$\alpha := \inf_{\substack{v \in V \\ v \neq 0}} \frac{a(v, v)}{\|v\|_V^2}$$

Introduction

So far we have treated **coercive problems**. This means that in the linear variational problem, we are trying to solve,

$$\begin{cases} \text{find } u \in V \text{ such that} \\ a(u, v) = F(v), \text{ for all } v \in V \end{cases} \quad (1)$$

the bilinear form $a(u, v)$ **satisfies**

$$a(v, v) \geq \alpha \|v\|_V^2, \quad \forall v \in V$$

for some $\alpha > 0$.

Recall that the best constant α satisfying the definition is given by

$$\alpha := \inf_{\substack{v \in V \\ v \neq 0}} \frac{a(v, v)}{\|v\|_V^2}$$

We now consider **noncoercive problems**, one for which no such $\alpha > 0$ exists. We will develop more general (necessary and sufficient) criteria for the well-posedness of the linear variational problem, known as the inf-sup or Babuška conditions.

For coercive problems, well-posedness is inherited for $V_h \subset V$. This is **not true for noncoercive problems**. Well-posedness is not inherited for an arbitrary $V_h \subset V$. One must prove the stability of each candidate discretisation individually.

We now consider **noncoercive problems**, one for which no such $\alpha > 0$ exists. We will develop more general **(necessary and sufficient)** criteria for the well-posedness of **the linear variational problem**, known as **the inf-sup or Babuška conditions**.

For coercive problems, well-posedness is inherited for $V_h \subset V$. This is **not true for noncoercive problems**. Well-posedness **is not inherited for an arbitrary $V_h \subset V$** . One must prove the stability of each candidate discretisation individually.

We now consider **noncoercive problems**, one for which no such $\alpha > 0$ exists. We will develop more general **(necessary and sufficient)** criteria for the well-posedness of **the linear variational problem**, known as **the inf-sup or Babuška conditions**.

For coercive problems, well-posedness is inherited for $V_h \subset V$. This is **not true for noncoercive problems**. Well-posedness **is not inherited for an arbitrary $V_h \subset V$** . One must prove the stability of each candidate discretisation individually.

We now consider **noncoercive problems**, one for which no such $\alpha > 0$ exists. We will develop more general (**necessary and sufficient**) criteria for the well-posedness of **the linear variational problem**, known as **the inf-sup or Babuška conditions**.

For coercive problems, well-posedness is inherited for $V_h \subset V$. This is **not true for noncoercive problems**. Well-posedness **is not inherited for an arbitrary $V_h \subset V$** . One must prove the stability of each candidate discretisation individually.

We now consider **noncoercive problems**, one for which no such $\alpha > 0$ exists. We will develop more general (**necessary and sufficient**) criteria for the well-posedness of **the linear variational problem**, known as **the inf-sup or Babuška conditions**.

For coercive problems, well-posedness is inherited for $V_h \subset V$. This is **not true for noncoercive problems**. Well-posedness **is not inherited for an arbitrary $V_h \subset V$** . One must prove the stability of each candidate discretisation individually.

Helmholtz equation

As an example of non-coercive, we can show that the Helmholtz equation

$$\begin{aligned} -\Delta u - k^2 u &= f \text{ in } \Omega \\ u &= 0 \text{ on } \partial\Omega \end{aligned}$$

is well-posed if k^2 is not an eigenvalue of the Dirichlet Laplacian, **but is not coercive for k large enough**. For k^2 to be an eigenvalue of the Dirichlet Laplacian, it means that there exists $u \neq 0$ such that

$$\begin{aligned} -\Delta u &= k^2 u \text{ in } \Omega \\ u &= 0 \quad \text{on } \partial\Omega \end{aligned}$$

i.e. $-\Delta - k^2 I$ has a nontrivial kernel.

Helmholtz equation

As an example of non-coercive, we can show that the Helmholtz equation

$$\begin{aligned} -\Delta u - k^2 u &= f \text{ in } \Omega \\ u &= 0 \text{ on } \partial\Omega \end{aligned}$$

is well-posed if k^2 is not an eigenvalue of the Dirichlet Laplacian, **but is not coercive for k large enough**. For k^2 to be an eigenvalue of the Dirichlet Laplacian, it means that there exists $u \neq 0$ such that

$$\begin{aligned} -\Delta u &= k^2 u \text{ in } \Omega \\ u &= 0 \quad \text{on } \partial\Omega \end{aligned}$$

i.e. $-\Delta - k^2 I$ has a nontrivial kernel.

Helmholtz equation

As an example of non-coercive, we can show that the Helmholtz equation

$$\begin{aligned}-\Delta u - k^2 u &= f \text{ in } \Omega \\ u &= 0 \text{ on } \partial\Omega\end{aligned}$$

is well-posed if k^2 is not an eigenvalue of the Dirichlet Laplacian, **but is not coercive for k large enough**. For k^2 to be an eigenvalue of the Dirichlet Laplacian, it means that there exists $u \neq 0$ such that

$$\begin{aligned}-\Delta u &= k^2 u \text{ in } \Omega \\ u &= 0 \quad \text{on } \partial\Omega\end{aligned}$$

i.e. $-\Delta - k^2 I$ has a nontrivial kernel.

Helmholtz equation

As an example of non-coercive, we can show that the Helmholtz equation

$$\begin{aligned} -\Delta u - k^2 u &= f \text{ in } \Omega \\ u &= 0 \text{ on } \partial\Omega \end{aligned}$$

is well-posed if k^2 is not an eigenvalue of the Dirichlet Laplacian, **but is not coercive for k large enough**. For k^2 to be an eigenvalue of the Dirichlet Laplacian, it means that there exists $u \neq 0$ such that

$$\begin{aligned} -\Delta u &= k^2 u \text{ in } \Omega \\ u &= 0 \quad \text{on } \partial\Omega \end{aligned}$$

i.e. $-\Delta - k^2 I$ has a nontrivial kernel.

Suppose we want to accurately determine the flux in the Poisson equation.

We can solve the mixed formulation:

find $\sigma : \Omega \rightarrow \mathbb{R}^n, u : \Omega \rightarrow \mathbb{R}$ such that

$$\begin{aligned}\sigma &= -\nabla u && \text{in } \Omega, \\ \operatorname{div} \sigma &= f && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega.\end{aligned}$$

Mixed Laplacian

Suppose we want to accurately determine the flux in the Poisson equation.

We can solve the mixed formulation:

find $\sigma : \Omega \rightarrow \mathbb{R}^n, u : \Omega \rightarrow \mathbb{R}$ such that

$$\begin{aligned}\sigma &= -\nabla u && \text{in } \Omega, \\ \operatorname{div} \sigma &= f && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega.\end{aligned}$$

Let's multiply the first equation by a vector-valued test function τ , and the second by a scalar-valued function v :

$$\begin{aligned}\int_{\Omega} \sigma \cdot \tau \, dx + \int_{\Omega} \nabla u \cdot \tau &= 0 \\ \int_{\Omega} \operatorname{div}(\sigma) v \, dx &= \int_{\Omega} f v \, dx\end{aligned}$$

Since σ needs to have a divergence, and we want τ and σ to come from the same space, let's integrate by parts in the first equation. For symmetry, I'll negate the second equation:

$$\begin{aligned}\int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} u \operatorname{div}(\tau) + \int_{\partial\Omega} u \tau \cdot n \, ds &= 0 \\ - \int_{\Omega} \operatorname{div}(\sigma) v \, dx &= - \int_{\Omega} f v \, dx\end{aligned}$$

mixed Laplacian

Let's multiply the first equation by a vector-valued test function τ , and the second by a scalar-valued function v :

$$\begin{aligned}\int_{\Omega} \sigma \cdot \tau \, dx + \int_{\Omega} \nabla u \cdot \tau &= 0 \\ \int_{\Omega} \operatorname{div}(\sigma) v \, dx &= \int_{\Omega} f v \, dx\end{aligned}$$

Since σ needs to have a divergence, and we want τ and σ to come from the same space, let's integrate by parts in the first equation. For symmetry, I'll negate the second equation:

$$\begin{aligned}\int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} u \operatorname{div}(\tau) + \int_{\partial\Omega} u \tau \cdot n \, ds &= 0 \\ - \int_{\Omega} \operatorname{div}(\sigma) v \, dx &= - \int_{\Omega} f v \, dx\end{aligned}$$

Let's multiply the first equation by a vector-valued test function τ , and the second by a scalar-valued function v :

$$\begin{aligned}\int_{\Omega} \sigma \cdot \tau \, dx + \int_{\Omega} \nabla u \cdot \tau &= 0 \\ \int_{\Omega} \operatorname{div}(\sigma) v \, dx &= \int_{\Omega} f v \, dx\end{aligned}$$

Since σ needs to have a divergence, and we want τ and σ to come from the same space, let's integrate by parts in the first equation. For symmetry, I'll negate the second equation:

$$\begin{aligned}\int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} u \operatorname{div}(\tau) + \int_{\partial\Omega} u \tau \cdot n \, ds &= 0 \\ - \int_{\Omega} \operatorname{div}(\sigma) v \, dx &= - \int_{\Omega} f v \, dx\end{aligned}$$

What function spaces do we need to make sense of

$$\begin{cases} \int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} u \operatorname{div} \tau = 0 \\ - \int_{\Omega} \operatorname{div} \sigma v \, dx = - \int_{\Omega} f v \, dx \end{cases}$$

We don't need any derivatives on u or v , so $u \in L^2(\Omega)$. For σ and τ , we need $\sigma \in L^2(\Omega; \mathbb{R}^n)$ and for $\operatorname{div} \sigma \in L^2(\Omega)$. This is the space $H(\operatorname{div}, \Omega)$:

$$H(\operatorname{div}, \Omega) = \{ \sigma \in L^2(\Omega; \mathbb{R}^n) : \operatorname{div}(\sigma) \in L^2(\Omega) \}$$

Its inner product is

$$(u, \tau)_{H(\operatorname{div}, \Omega)} = \int_{\Omega} u \cdot \tau + \operatorname{div} u \operatorname{div} \tau \, dx$$

What function spaces do we need to make sense of

$$\begin{cases} \int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} u \operatorname{div} \tau = 0 \\ - \int_{\Omega} \operatorname{div} \sigma v \, dx = - \int_{\Omega} f v \, dx \end{cases}$$

We don't need any derivatives on u or v , so $u \in L^2(\Omega)$. For σ and τ , we need $\sigma \in L^2(\Omega; \mathbb{R}^n)$ and for $\operatorname{div} \sigma \in L^2(\Omega)$. This is the space $H(\operatorname{div}, \Omega)$:

$$H(\operatorname{div}, \Omega) = \{ \sigma \in L^2(\Omega; \mathbb{R}^n) : \operatorname{div}(\sigma) \in L^2(\Omega) \}$$

Its inner product is

$$(u, \tau)_{H(\operatorname{div}, \Omega)} = \int_{\Omega} u \cdot \tau + \operatorname{div} u \operatorname{div} \tau \, dx$$

Mixed Laplacian

A nice property of variational problems is that we can add the two equations together. The problem is the same as:

$$\begin{cases} \text{Find } (\sigma, u) \in H(\operatorname{div}, \Omega) \times L^2(\Omega) \text{ such that} \\ \int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} \operatorname{div}(\tau)u - \int_{\Omega} \operatorname{div}(\sigma)v \, dx = - \int_{\Omega} fv \, dx \end{cases}$$

for all $(\tau, v) \in H(\operatorname{div}, \Omega) \times L^2(\Omega)$.

We define:

$$B(\sigma, u; \tau, v) := \int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} \operatorname{div}(\tau)u - \int_{\Omega} \operatorname{div}(\sigma)v \, dx$$

Lax-Milgram certainly won't apply:

$$B(0, u; 0, u) = 0 \text{ for all } u \in L^2(\Omega)$$

Mixed Laplacian

A nice property of variational problems is that we can add the two equations together. The problem is the same as:

$$\begin{cases} \text{Find } (\sigma, u) \in H(\operatorname{div}, \Omega) \times L^2(\Omega) \text{ such that} \\ \int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} \operatorname{div}(\tau)u - \int_{\Omega} \operatorname{div}(\sigma)v \, dx = - \int_{\Omega} fv \, dx \end{cases}$$

for all $(\tau, v) \in H(\operatorname{div}, \Omega) \times L^2(\Omega)$.

We define:

$$B(\sigma, u; \tau, v) := \int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} \operatorname{div}(\tau)u - \int_{\Omega} \operatorname{div}(\sigma)v \, dx$$

Lax-Milgram certainly won't apply:

$$B(0, u; 0, u) = 0 \text{ for all } u \in L^2(\Omega)$$

Mixed Laplacian

A nice property of variational problems is that we can add the two equations together. The problem is the same as:

$$\begin{cases} \text{Find } (\sigma, u) \in H(\operatorname{div}, \Omega) \times L^2(\Omega) \text{ such that} \\ \int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} \operatorname{div}(\tau)u - \int_{\Omega} \operatorname{div}(\sigma)v \, dx = - \int_{\Omega} fv \, dx \end{cases}$$

for all $(\tau, v) \in H(\operatorname{div}, \Omega) \times L^2(\Omega)$.

We define:

$$B(\sigma, u; \tau, v) := \int_{\Omega} \sigma \cdot \tau \, dx - \int_{\Omega} \operatorname{div}(\tau)u - \int_{\Omega} \operatorname{div}(\sigma)v \, dx$$

Lax-Milgram certainly won't apply:

$$B(0, u; 0, u) = 0 \text{ for all } u \in L^2(\Omega)$$

Theorem (Babuška theorem)

Let V and W be two Hilbert spaces. Let $a : V \times W \rightarrow \mathbb{R}$ be a bilinear form for which there exist constants $C < \infty, \gamma > 0, \gamma' > 0$ such that:

1 $|a(v, w)| \leq C \|v\|_V \|w\|_W, \quad \text{for all } v \in V, w \in W,$

2 $\inf_{\substack{v \in V \\ v \neq 0}} \sup_{\substack{w \in W \\ w \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma > 0,$

3 $\inf_{\substack{w \in W \\ w \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma' > 0.$

Then for any $F \in W'$ there exists exactly one solution $u \in V$ such that:

$$a(u, w) = F(w), \quad \text{for all } w \in W.$$

Furthermore, the problem is stable, i.e.,

$$\|u\|_V \leq \frac{\|F\|_{W'}}{\gamma}$$

Theorem (Babuška theorem)

Let V and W be two Hilbert spaces. Let $a : V \times W \rightarrow \mathbb{R}$ be a bilinear form for which there exist constants $C < \infty, \gamma > 0, \gamma' > 0$ such that:

① $|a(v, w)| \leq C \|v\|_V \|w\|_W, \quad \text{for all } v \in V, w \in W,$

② $\inf_{\substack{v \in V \\ v \neq 0}} \sup_{\substack{w \in W \\ w \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma > 0,$

③ $\inf_{\substack{w \in W \\ w \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma' > 0.$

Then for any $F \in W'$ there exists exactly one solution $u \in V$ such that:

$$a(u, w) = F(w), \quad \text{for all } w \in W.$$

Furthermore, the problem is stable, i.e.,

$$\|u\|_V \leq \frac{\|F\|_{W'}}{\gamma}$$

Theorem (Babuška theorem)

Let V and W be two Hilbert spaces. Let $a : V \times W \rightarrow \mathbb{R}$ be a bilinear form for which there exist constants $C < \infty, \gamma > 0, \gamma' > 0$ such that:

$$\textcircled{1} \quad |a(v, w)| \leq C \|v\|_V \|w\|_W, \quad \text{for all } v \in V, w \in W,$$

$$\textcircled{2} \quad \inf_{\substack{v \in V \\ v \neq 0}} \sup_{\substack{w \in W \\ w \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma > 0,$$

$$\textcircled{3} \quad \inf_{\substack{w \in W \\ w \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma' > 0.$$

Then for any $F \in W'$ there exists exactly one solution $u \in V$ such that:

$$a(u, w) = F(w), \quad \text{for all } w \in W.$$

Furthermore, the problem is stable, i.e.,

$$\|u\|_V \leq \frac{\|F\|_{W'}}{\gamma}$$

Theorem (Babuška theorem)

Let V and W be two Hilbert spaces. Let $a : V \times W \rightarrow \mathbb{R}$ be a bilinear form for which there exist constants $C < \infty, \gamma > 0, \gamma' > 0$ such that:

① $|a(v, w)| \leq C \|v\|_V \|w\|_W, \quad \text{for all } v \in V, w \in W,$

② $\inf_{\substack{v \in V \\ v \neq 0}} \sup_{\substack{w \in W \\ w \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma > 0,$

③ $\inf_{\substack{w \in W \\ w \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma' > 0.$

Then for any $F \in W'$ there exists exactly one solution $u \in V$ such that:

$$a(u, w) = F(w), \quad \text{for all } w \in W.$$

Furthermore, the problem is stable, i.e.,

$$\|u\|_V \leq \frac{\|F\|_{W'}}{\gamma}$$

Theorem (Babuška theorem)

Let V and W be two Hilbert spaces. Let $a : V \times W \rightarrow \mathbb{R}$ be a bilinear form for which there exist constants $C < \infty, \gamma > 0, \gamma' > 0$ such that:

$$\textcircled{1} \quad |a(v, w)| \leq C \|v\|_V \|w\|_W, \quad \text{for all } v \in V, w \in W,$$

$$\textcircled{2} \quad \inf_{\substack{v \in V \\ v \neq 0}} \sup_{\substack{w \in W \\ w \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma > 0,$$

$$\textcircled{3} \quad \inf_{\substack{w \in W \\ w \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(v, w)}{\|v\|_V \|w\|_W} \geq \gamma' > 0.$$

Then for any $F \in W'$ there exists exactly one solution $u \in V$ such that:

$$a(u, w) = F(w), \quad \text{for all } w \in W.$$

Furthermore, the problem is stable, i.e.,

$$\|u\|_V \leq \frac{\|F\|_{W'}}{\gamma}$$

As a first example of how to manipulate inf-sup conditions, let's show that a coercive problem satisfies the inf-sup conditions. Suppose $a(u, v)$ satisfies

$$\alpha \|u\|_V^2 \leq a(u, u) \quad \text{for all } u \in V$$

Dividing both sides of the inequality by $\|u\|_V$ for $u \neq 0$, we have

$$\begin{aligned} \alpha \|u\|_V &\leq \frac{a(u, u)}{\|u\|_V} \\ &\leq \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|v\|_V} \end{aligned}$$

Infimising over $u \neq 0$, we have

$$0 < \alpha \leq \inf_{\substack{u \in V \\ u \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|u\|_V \|v\|_V}$$

So the coercivity constant α works for γ (and γ').

As a first example of how to manipulate inf-sup conditions, let's show that a coercive problem satisfies the inf-sup conditions. Suppose $a(u, v)$ satisfies

$$\alpha \|u\|_V^2 \leq a(u, u) \quad \text{for all } u \in V$$

Dividing both sides of the inequality by $\|u\|_V$ for $u \neq 0$, we have

$$\begin{aligned} \alpha \|u\|_V &\leq \frac{a(u, u)}{\|u\|_V} \\ &\leq \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|v\|_V} \end{aligned}$$

Infimising over $u \neq 0$, we have

$$0 < \alpha \leq \inf_{\substack{u \in V \\ u \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|u\|_V \|v\|_V}$$

So the coercivity constant α works for γ (and γ').

As a first example of how to manipulate inf-sup conditions, let's show that a coercive problem satisfies the inf-sup conditions. Suppose $a(u, v)$ satisfies

$$\alpha \|u\|_V^2 \leq a(u, u) \quad \text{for all } u \in V$$

Dividing both sides of the inequality by $\|u\|_V$ for $u \neq 0$, we have

$$\begin{aligned} \alpha \|u\|_V &\leq \frac{a(u, u)}{\|u\|_V} \\ &\leq \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|v\|_V} \end{aligned}$$

Infimising over $u \neq 0$, we have

$$0 < \alpha \leq \inf_{\substack{u \in V \\ u \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|u\|_V \|v\|_V}$$

So the coercivity constant α works for γ (and γ').

As a first example of how to manipulate inf-sup conditions, let's show that a coercive problem satisfies the inf-sup conditions. Suppose $a(u, v)$ satisfies

$$\alpha \|u\|_V^2 \leq a(u, u) \quad \text{for all } u \in V$$

Dividing both sides of the inequality by $\|u\|_V$ for $u \neq 0$, we have

$$\begin{aligned} \alpha \|u\|_V &\leq \frac{a(u, u)}{\|u\|_V} \\ &\leq \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|v\|_V} \end{aligned}$$

Infimising over $u \neq 0$, we have

$$0 < \alpha \leq \inf_{\substack{u \in V \\ u \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|u\|_V \|v\|_V}$$

So the coercivity constant α works for γ (and γ').

As a first example of how to manipulate inf-sup conditions, let's show that a coercive problem satisfies the inf-sup conditions. Suppose $a(u, v)$ satisfies

$$\alpha \|u\|_V^2 \leq a(u, u) \quad \text{for all } u \in V$$

Dividing both sides of the inequality by $\|u\|_V$ for $u \neq 0$, we have

$$\begin{aligned} \alpha \|u\|_V &\leq \frac{a(u, u)}{\|u\|_V} \\ &\leq \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|v\|_V} \end{aligned}$$

Infimising over $u \neq 0$, we have

$$0 < \alpha \leq \inf_{\substack{u \in V \\ u \neq 0}} \sup_{\substack{v \in V \\ v \neq 0}} \frac{a(u, v)}{\|u\|_V \|v\|_V}$$

So the coercivity constant α works for γ (and γ').

Start with find $u \in V$ such that

$$a(u, v) = F(v), \text{ for all } v \in V,$$

and take the Galerkin approximation over closed $V_h \subset V$:
find $u_h \in V_h$ such that

$$a(u_h, v_h) = F(v_h) \text{ for all } v_h \in V_h.$$

Note that Galerkin orthogonality still holds.

Is the discrete problem well-posed?

Start with find $u \in V$ such that

$$a(u, v) = F(v), \text{ for all } v \in V,$$

and take the Galerkin approximation over closed $V_h \subset V$:
find $u_h \in V_h$ such that

$$a(u_h, v_h) = F(v_h) \text{ for all } v_h \in V_h.$$

Note that Galerkin orthogonality still holds.

Is the discrete problem well-posed?

Let's check the Babuška conditions.

- 1 Satisfaction of (1) is inherited.
- 2 What about (2)?

That is, does there exist $\tilde{\gamma}$ such that

$$\inf_{\substack{u_h \in V_h \\ u_h \neq 0}} \sup_{\substack{v_h \in V_h \\ v_h \neq 0}} \frac{a(u_h, v_h)}{\|u_h\|_V \|v_h\|_V} \geq \tilde{\gamma} > 0$$

with $\tilde{\gamma}$ independent of the mesh size h ?No! Examples later.

Remark

We don't need to check (3) in this case! The discrete system is square and finite-dimensional, so (2) \iff (3) by rank-nullity.)

Let's check the Babuška conditions.

- 1 Satisfaction of (1) is inherited.
- 2 What about (2)?

That is, does there exist $\tilde{\gamma}$ such that

$$\inf_{\substack{u_h \in V_h \\ u_h \neq 0}} \sup_{\substack{v_h \in V_h \\ v_h \neq 0}} \frac{a(u_h, v_h)}{\|u_h\|_V \|v_h\|_V} \geq \tilde{\gamma} > 0$$

with $\tilde{\gamma}$ independent of the mesh size h ?No! Examples later.

Remark

We don't need to check (3) in this case! The discrete system is square and finite-dimensional, so (2) \iff (3) by rank-nullity.)

Let's check the Babuška conditions.

- 1 Satisfaction of (1) is inherited.
- 2 What about (2)?

That is, does there exist $\tilde{\gamma}$ such that

$$\inf_{\substack{u_h \in V_h \\ u_h \neq 0}} \sup_{\substack{v_h \in V_h \\ v_h \neq 0}} \frac{a(u_h, v_h)}{\|u_h\|_V \|v_h\|_V} \geq \tilde{\gamma} > 0$$

with $\tilde{\gamma}$ independent of the mesh size h ? No! Examples later.

Remark

We don't need to check (3) in this case! The discrete system is square and finite-dimensional, so (2) \iff (3) by rank-nullity.)

Let's check the Babuška conditions.

- 1 Satisfaction of (1) is inherited.
- 2 What about (2)?

That is, does there exist $\tilde{\gamma}$ such that

$$\inf_{\substack{u_h \in V_h \\ u_h \neq 0}} \sup_{\substack{v_h \in V_h \\ v_h \neq 0}} \frac{a(u_h, v_h)}{\|u_h\|_V \|v_h\|_V} \geq \tilde{\gamma} > 0$$

with $\tilde{\gamma}$ independent of the mesh size h ?No! Examples later.

Remark

We don't need to check (3) in this case! The discrete system is square and finite-dimensional, so (2) \iff (3) by rank-nullity.)

Let's check the Babuška conditions.

- 1 Satisfaction of (1) is inherited.
- 2 What about (2)?

That is, does there exist $\tilde{\gamma}$ such that

$$\inf_{\substack{u_h \in V_h \\ u_h \neq 0}} \sup_{\substack{v_h \in V_h \\ v_h \neq 0}} \frac{a(u_h, v_h)}{\|u_h\|_V \|v_h\|_V} \geq \tilde{\gamma} > 0$$

with $\tilde{\gamma}$ independent of the mesh size h ?No! Examples later.

Remark

We don't need to check (3) in this case! The discrete system is square and finite-dimensional, so (2) \iff (3) by rank-nullity.)

Let's check the Babuška conditions.

- 1 Satisfaction of (1) is inherited.
- 2 What about (2)?

That is, does there exist $\tilde{\gamma}$ such that

$$\inf_{\substack{u_h \in V_h \\ u_h \neq 0}} \sup_{\substack{v_h \in V_h \\ v_h \neq 0}} \frac{a(u_h, v_h)}{\|u_h\|_V \|v_h\|_V} \geq \tilde{\gamma} > 0$$

with $\tilde{\gamma}$ independent of the mesh size h ?No! Examples later.

Remark

We don't need to check (3) in this case! The discrete system is square and finite-dimensional, so (2) \iff (3) by rank-nullity.)

Theorem

Assume we have a well-posed discretisation of a well-posed problem. Then

$$\|u - u_h\|_V \leq \left(1 + \frac{C}{\tilde{\gamma}}\right) \|u - v_h\|_V$$

For every $v_h \in V_h$, we have

$$\begin{aligned}
 \tilde{\gamma} \|v_h - u_h\|_V &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u_h, w_h)}{\|w_h\|_V} && \text{(discrete inf-sup)} \\
 &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h) + a(u - u_h, w_h)}{\|w_h\|_V} \\
 &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h)}{\|w_h\|_V} && \text{(Galerkin orth.)} \\
 &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{C \|v_h - u\|_V \|w_h\|_V}{\|w_h\|_V}
 \end{aligned}$$

For every $v_h \in V_h$, we have

$$\begin{aligned}\tilde{\gamma} \|v_h - u_h\|_V &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u_h, w_h)}{\|w_h\|_V} && \text{(discrete inf-sup)} \\ &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h) + a(u - u_h, w_h)}{\|w_h\|_V} \\ &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h)}{\|w_h\|_V} && \text{(Galerkin orth.)} \\ &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{C \|v_h - u\|_V \|w_h\|_V}{\|w_h\|_V}\end{aligned}$$

For every $v_h \in V_h$, we have

$$\begin{aligned}\tilde{\gamma} \|v_h - u_h\|_V &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u_h, w_h)}{\|w_h\|_V} && \text{(discrete inf-sup)} \\ &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h) + a(u - u_h, w_h)}{\|w_h\|_V} \\ &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h)}{\|w_h\|_V} && \text{(Galerkin orth.)} \\ &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{C \|v_h - u\|_V \|w_h\|_V}{\|w_h\|_V}\end{aligned}$$

For every $v_h \in V_h$, we have

$$\begin{aligned}
 \tilde{\gamma} \|v_h - u_h\|_V &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u_h, w_h)}{\|w_h\|_V} && \text{(discrete inf-sup)} \\
 &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h) + a(u - u_h, w_h)}{\|w_h\|_V} \\
 &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h)}{\|w_h\|_V} && \text{(Galerkin orth.)} \\
 &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{C \|v_h - u\|_V \|w_h\|_V}{\|w_h\|_V}
 \end{aligned}$$

For every $v_h \in V_h$, we have

$$\begin{aligned}
 \tilde{\gamma} \|v_h - u_h\|_V &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u_h, w_h)}{\|w_h\|_V} && \text{(discrete inf-sup)} \\
 &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h) + a(u - u_h, w_h)}{\|w_h\|_V} \\
 &= \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{a(v_h - u, w_h)}{\|w_h\|_V} && \text{(Galerkin orth.)} \\
 &\leq \sup_{\substack{w_h \in V_h \\ w_h \neq 0}} \frac{C \|v_h - u\|_V \|w_h\|_V}{\|w_h\|_V}
 \end{aligned}$$

Now apply the triangle inequality to $\|u - u_h\|_V$:

$$\begin{aligned}\|u - u_h\|_V &\leq \|u - v_h\|_V + \|v_h - u_h\|_V \\ &\leq \|u - v_h\|_V + \frac{C}{\tilde{\gamma}} \|u - v_h\|_V \\ &= \left(1 + \frac{C}{\tilde{\gamma}}\right) \|u - v_h\|_V\end{aligned}$$

As before, we can combine this with an approximation result and a regularity result to derive error estimates for finite element discretisations.

Now apply the triangle inequality to $\|u - u_h\|_V$:

$$\begin{aligned}\|u - u_h\|_V &\leq \|u - v_h\|_V + \|v_h - u_h\|_V \\ &\leq \|u - v_h\|_V + \frac{C}{\tilde{\gamma}} \|u - v_h\|_V \\ &= \left(1 + \frac{C}{\tilde{\gamma}}\right) \|u - v_h\|_V\end{aligned}$$

As before, we can combine this with an approximation result and a regularity result to derive error estimates for finite element discretisations.

Now apply the triangle inequality to $\|u - u_h\|_V$:

$$\begin{aligned}\|u - u_h\|_V &\leq \|u - v_h\|_V + \|v_h - u_h\|_V \\ &\leq \|u - v_h\|_V + \frac{C}{\tilde{\gamma}} \|u - v_h\|_V \\ &= \left(1 + \frac{C}{\tilde{\gamma}}\right) \|u - v_h\|_V\end{aligned}$$

As before, we can combine this with an approximation result and a regularity result to derive error estimates for finite element discretisations.

Now apply the triangle inequality to $\|u - u_h\|_V$:

$$\begin{aligned}\|u - u_h\|_V &\leq \|u - v_h\|_V + \|v_h - u_h\|_V \\ &\leq \|u - v_h\|_V + \frac{C}{\tilde{\gamma}} \|u - v_h\|_V \\ &= \left(1 + \frac{C}{\tilde{\gamma}}\right) \|u - v_h\|_V\end{aligned}$$

As before, we can combine this with an approximation result and a regularity result to derive error estimates for finite element discretisations.

Now apply the triangle inequality to $\|u - u_h\|_V$:

$$\begin{aligned}\|u - u_h\|_V &\leq \|u - v_h\|_V + \|v_h - u_h\|_V \\ &\leq \|u - v_h\|_V + \frac{C}{\tilde{\gamma}} \|u - v_h\|_V \\ &= \left(1 + \frac{C}{\tilde{\gamma}}\right) \|u - v_h\|_V\end{aligned}$$

As before, we can combine this with an approximation result and a regularity result to derive error estimates for finite element discretisations.

Now apply the triangle inequality to $\|u - u_h\|_V$:

$$\begin{aligned}\|u - u_h\|_V &\leq \|u - v_h\|_V + \|v_h - u_h\|_V \\ &\leq \|u - v_h\|_V + \frac{C}{\tilde{\gamma}} \|u - v_h\|_V \\ &= \left(1 + \frac{C}{\tilde{\gamma}}\right) \|u - v_h\|_V\end{aligned}$$

As before, we can combine this with an approximation result and a regularity result to derive error estimates for finite element discretisations.